Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
United States												
All races, both sexes	87,391	626	131	55	59	89	159	292	557	1,305	3,186	5,411
Male	44,398	388	80	31	36	44	90	170	336	819	2,038	3,653
Female	42,993	238	51	24	23	45	69	122	221	486	1,148	1,758
White, both sexes	75,864	390	93	39	32	66	106	224	408	1,047	2,617	4,403
Male	38,270	239	54	19	18	33	60	136	256	671	1,700	2,999
Female	37,594	151	39	20	14	33	46	88	152	376	917	1,404
Black, both sexes	9,088	213	30	14	23	18	50	50	98	179	406	812
Male	4,801	140	20	10	17	10	28	22	48	101	239	527
Female	4,287	73	10	4	6	8	22	28	50	78	167	285
American Indian 1/, both sexes	1,049	10	5	-	1	2	2	13	32	52	118	129
Male	564	3	3	-	1	-	2	8	19	25	66	76
Female	485	7	2	-	-	2	-	5	13	27	52	53
Asian or Pacific Islander 2/, both sexes	1,390	13	3	2	3	3	1	5	19	27	45	67
Male	763	6	3	2	-	1	-	4	13	22	33	51
Female	627	7	-	-	3	2	1	1	6	5	12	16
Alabama												
All races, both sexes	1,608	17	-	2	1	3	2	5	14	22	61	104
Male	797	11	-	2	-	1	2	1	7	12	37	70
Female	811	6	-	-	1	2	-	4	7	10	24	34
White, both sexes	1,255	5	-	1	-	-	1	3	9	15	41	73
Male	625	5	-	1	-	-	1	1	6	10	24	53
Female	630	-	-	-	-	-	-	2	3	5	17	20
Black, both sexes	349	12	-	1	1	3	1	2	5	7	18	31
Male	169	6	-	1	-	1	1	-	1	2	11	17
Female	180	6	-	-	1	2	-	2	4	5	7	14
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Alaska												
All races, both sexes	134	2	-	-	-	-	-	-	-	4	11	9
Male	62	1	-	-	-	-	-	-	-	1	4	2
Female	72	1	-	-	-	-	-	-	-	3	7	7
White, both sexes	94	1	-	-	-	-	-	-	-	4	7	5
Male	48	1	-	-	-	-	-	-	-	1	3	2
Female	46	-	-	-	-	-	-	-	-	3	4	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	36	1	-	-	-	-	-	-	-	-	3	4
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	-	-	-	-	3	4
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,807	18	5	1	-	1	2	9	16	42	82	123
Male	1,002	10	3	-	-	-	2	7	13	28	54	76
Female	805	8	2	1	-	1	-	2	3	14	28	47
White, both sexes	1,592	17	4	1	-	1	2	3	9	32	62	91
Male	882	9	3	-	-	-	2	2	9	23	40	59
Female	710	8	1	1	-	1	-	1	-	9	22	32
Black, both sexes	42	1	-	-	-	-	-	1	-	1	3	3
Male	22	1	-	-	-	-	-	1	-	-	2	1
Female	20	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	161	-	1	-	-	-	-	5	6	9	17	26
Male	89	-	-	-	-	-	-	4	3	5	12	13
Female	72	-	1	-	-	-	-	1	3	4	5	13
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	1	-	-	3
Male	9	-	-	-	-	-	-	-	1	-	-	3
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Arkansas												
All races, both sexes	958	6	3	-	1	-	3	1	8	14	23	48
Male	455	6	1	-	1	-	2	-	3	5	10	33
Female	503	-	2	-	-	-	1	1	5	9	13	15
White, both sexes	835	3	3	-	1	-	1	-	4	12	18	43
Male	400	3	1	-	1	-	-	-	2	4	8	31
Female	435	-	2	-	-	-	1	-	2	8	10	12
Black, both sexes	118	3	-	-	-	-	2	1	4	2	4	5
Male	52	3	-	-	-	-	2	-	1	1	1	2
Female	66	-	-	-	-	-	-	1	3	1	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9,277	72	17	7	12	14	14	47	75	194	410	671
Male	5,157	42	12	3	7	9	7	34	52	131	272	467
Female	4,120	30	5	4	5	5	7	13	23	63	138	204
White, both sexes	8,011	55	14	7	7	12	11	39	63	175	348	574
Male	4,456	31	10	3	5	8	5	29	42	120	234	396
Female	3,555	24	4	4	2	4	6	10	21	55	114	178
Black, both sexes	613	15	1	-	3	1	3	6	2	10	33	63
Male	342	10	1	-	2	1	2	4	2	5	19	44
Female	271	5	-	-	1	-	1	2	-	5	14	19
American Indian 1/, both sexes	102	-	1	-	-	-	-	1	4	3	9	9
Male	59	-	-	-	-	-	-	-	4	2	5	8
Female	43	-	1	-	-	-	-	1	-	1	4	1
Asian or Pacific Islander 2/, both sexes	551	2	1	-	2	1	-	1	6	6	20	25
Male	300	1	1	-	-	-	-	1	4	4	14	19
Female	251	1	-	-	2	1	-	-	2	2	6	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Colorado												
All races, both sexes	1,264	5	2	2	1	2	3	4	17	24	51	112
Male	672	4	2	-	1	2	1	3	11	12	37	78
Female	592	1	-	2	-	-	2	1	6	12	14	34
White, both sexes	1,204	4	1	2	1	2	3	4	16	22	50	105
Male	643	4	1	-	1	2	1	3	10	12	36	74
Female	561	-	-	2	-	-	2	1	6	10	14	31
Black, both sexes	36	1	1	-	-	-	-	-	-	1	1	5
Male	17	-	1	-	-	-	-	-	-	-	1	3
Female	19	1	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	1	1	-	2
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,151	9	2	1	1	1	3	2	3	13	34	44
Male	530	7	2	-	1	1	1	2	2	8	23	34
Female	621	2	-	1	-	-	2	-	1	5	11	10
White, both sexes	1,073	4	1	1	-	-	3	2	2	10	31	42
Male	492	3	1	-	-	-	1	2	2	7	21	33
Female	581	1	-	1	-	-	2	-	-	3	10	9
Black, both sexes	69	4	1	-	1	1	-	-	1	2	3	2
Male	34	3	1	-	1	1	-	-	-	1	2	1
Female	35	1	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Delaware												
All races, both sexes	228	-	-	-	1	-	-	1	-	2	8	14
Male	119	-	-	-	1	-	-	1	-	2	4	10
Female	109	-	-	-	-	-	-	-	-	-	4	4
White, both sexes	203	-	-	-	-	-	-	1	-	1	8	13
Male	108	-	-	-	-	-	-	1	-	1	4	9
Female	95	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	24	-	-	-	1	-	-	-	-	1	-	1
Male	10	-	-	-	1	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	198	2	2	-	-	-	-	1	1	1	9	14
Male	120	1	1	-	-	-	-	-	1	1	6	11
Female	78	1	1	-	-	-	-	1	-	-	3	3
White, both sexes	38	1	-	-	-	-	-	1	1	-	1	1
Male	22	-	-	-	-	-	-	-	1	-	1	1
Female	16	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	153	1	1	-	-	-	-	-	-	1	8	13
Male	92	1	-	-	-	-	-	-	-	1	5	10
Female	61	-	1	-	-	-	-	-	-	-	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Florida												
All races, both sexes	5,677	45	5	3	1	4	7	10	26	70	186	383
Male	2,989	29	1	2	1	3	3	4	13	39	116	251
Female	2,688	16	4	1	-	1	4	6	13	31	70	132
White, both sexes	5,140	20	4	1	1	4	2	7	20	59	163	329
Male	2,706	12	-	1	1	3	-	3	10	33	100	218
Female	2,434	8	4	-	-	1	2	4	10	26	63	111
Black, both sexes	498	24	-	2	-	-	5	3	6	10	21	54
Male	259	16	-	1	-	-	3	1	3	5	14	33
Female	239	8	-	1	-	-	2	2	3	5	7	21
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	1	-	-	-	-	-	-	1	2	-
Male	19	1	1	-	-	-	-	-	-	1	2	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,415	36	4	2	5	1	7	10	19	46	103	165
Male	1,242	25	4	-	2	-	3	5	12	28	63	111
Female	1,173	11	-	2	3	1	4	5	7	18	40	54
White, both sexes	1,788	12	2	2	2	-	6	8	9	28	74	107
Male	926	10	2	-	-	-	3	4	6	17	46	75
Female	862	2	-	2	2	-	3	4	3	11	28	32
Black, both sexes	604	22	1	-	3	1	1	2	10	16	27	55
Male	303	14	1	-	2	-	-	1	6	10	15	35
Female	301	8	-	-	1	1	1	1	4	6	12	20
American Indian 1/, both sexes	6	2	1	-	-	-	-	-	-	1	1	1
Male	3	1	1	-	-	-	-	-	-	-	1	-
Female	3	1	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Hawaii												
All races, both sexes	326	4	-	1	-	1	2	2	2	2	11	17
Male	178	2	-	1	-	-	-	2	1	2	6	12
Female	148	2	-	-	-	1	2	-	1	-	5	5
White, both sexes	115	1	-	-	-	-	2	1	1	-	2	7
Male	63	1	-	-	-	-	-	1	-	-	-	5
Female	52	-	-	-	-	-	2	-	1	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	207	3	-	1	-	1	-	1	1	2	9	10
Male	114	1	-	1	-	-	-	1	1	2	6	7
Female	93	2	-	-	-	1	-	-	-	-	3	3
Idaho												
All races, both sexes	408	5	-	-	-	1	-	2	2	6	12	22
Male	206	3	-	-	-	1	-	2	2	2	6	13
Female	202	2	-	-	-	-	-	-	-	4	6	9
White, both sexes	397	5	-	-	-	1	-	2	2	4	10	21
Male	202	3	-	-	-	1	-	2	2	2	6	13
Female	195	2	-	-	-	-	-	-	-	2	4	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	2	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	2	2	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Illinois												
All races, both sexes	3,585	16	2	4	2	2	9	21	28	47	145	214
Male	1,732	15	-	4	2	1	5	12	18	31	101	146
Female	1,853	1	2	-	-	1	4	9	10	16	44	68
White, both sexes	3,089	11	1	1	1	2	4	17	26	40	115	156
Male	1,455	10	-	1	1	1	4	11	17	26	82	104
Female	1,634	1	1	-	-	1	-	6	9	14	33	52
Black, both sexes	449	5	1	2	1	-	5	3	1	6	27	55
Male	252	5	-	2	1	-	1	1	1	4	17	40
Female	197	-	1	-	-	-	4	2	-	2	10	15
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	37	-	-	1	-	-	-	1	1	-	2	1
Male	20	-	-	1	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	1,834	22	5	2	2	-	5	4	14	25	54	108
Male	882	12	2	1	2	-	2	3	8	17	36	65
Female	952	10	3	1	-	-	3	1	6	8	18	43
White, both sexes	1,696	16	5	2	1	-	4	4	13	25	47	102
Male	810	8	2	1	1	-	1	3	7	17	31	62
Female	886	8	3	1	-	-	3	1	6	8	16	40
Black, both sexes	130	6	-	-	1	-	1	-	1	-	6	4
Male	69	4	-	-	1	-	1	-	1	-	4	3
Female	61	2	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Iowa												
All races, both sexes	880	2	1	-	1	1	1	2	6	8	16	45
Male	399	2	1	-	-	-	-	2	3	5	12	31
Female	481	-	-	-	1	1	1	-	3	3	4	14
White, both sexes	853	2	1	-	1	1	1	2	4	8	16	41
Male	387	2	1	-	-	-	-	2	2	5	12	29
Female	466	-	-	-	1	1	1	-	2	3	4	12
Black, both sexes	16	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	860	6	2	1	-	2	1	2	3	7	27	57
Male	422	4	1	1	-	1	1	1	2	4	22	39
Female	438	2	1	-	-	1	-	1	1	3	5	18
White, both sexes	809	4	1	1	-	2	1	2	3	7	24	50
Male	393	2	-	1	-	1	1	1	2	4	19	35
Female	416	2	1	-	-	1	-	1	1	3	5	15
Black, both sexes	31	2	1	-	-	-	-	-	-	-	1	5
Male	17	2	1	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	2	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Kentucky												
All races, both sexes	1,305	6	3	-	-	-	1	5	10	19	46	81
Male	669	5	1	-	-	-	1	2	4	16	33	58
Female	636	1	2	-	-	-	-	3	6	3	13	23
White, both sexes	1,227	4	3	-	-	-	1	4	9	19	41	80
Male	624	3	1	-	-	-	1	2	4	16	30	58
Female	603	1	2	-	-	-	-	2	5	3	11	22
Black, both sexes	72	2	-	-	-	-	-	1	1	-	5	1
Male	40	2	-	-	-	-	-	-	-	-	3	-
Female	32	-	-	-	-	-	-	1	1	-	2	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,394	8	2	1	2	3	4	7	17	20	59	98
Male	705	3	1	-	2	1	3	4	10	9	38	67
Female	689	5	1	1	-	2	1	3	7	11	21	31
White, both sexes	980	2	1	-	1	-	-	4	7	7	35	62
Male	483	1	-	-	1	-	-	3	5	2	22	41
Female	497	1	1	-	-	-	-	1	2	5	13	21
Black, both sexes	403	6	1	1	1	3	4	3	10	13	24	35
Male	217	2	1	-	1	1	3	1	5	7	16	25
Female	186	4	-	1	-	2	1	2	5	6	8	10
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Maine												
All races, both sexes	482	2	-	-	-	-	-	2	1	3	7	19
Male	248	1	-	-	-	-	-	1	1	1	5	11
Female	234	1	-	-	-	-	-	1	-	2	2	8
White, both sexes	479	2	-	-	-	-	-	2	1	3	7	19
Male	246	1	-	-	-	-	-	1	1	1	5	11
Female	233	1	-	-	-	-	-	1	-	2	2	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,512	10	-	2	-	1	3	7	5	25	50	88
Male	763	6	-	2	-	1	2	3	1	13	27	55
Female	749	4	-	-	-	-	1	4	4	12	23	33
White, both sexes	1,116	6	-	1	-	1	2	3	3	13	34	54
Male	547	4	-	1	-	1	2	-	1	7	20	35
Female	569	2	-	-	-	-	-	3	2	6	14	19
Black, both sexes	376	4	-	1	-	-	1	4	1	11	15	34
Male	209	2	-	1	-	-	-	3	-	6	6	20
Female	167	2	-	-	-	-	1	1	1	5	9	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Massachusetts												
All races, both sexes	2,122	8	1	1	1	1	1	3	11	24	72	109
Male	1,040	6	1	1	1	1	-	3	8	13	42	74
Female	1,082	2	-	-	-	-	1	-	3	11	30	35
White, both sexes	2,017	7	1	1	1	-	1	2	10	24	69	98
Male	983	6	1	1	1	-	-	2	7	13	40	67
Female	1,034	1	-	-	-	-	1	-	3	11	29	31
Black, both sexes	81	1	-	-	-	-	-	-	1	-	2	9
Male	43	-	-	-	-	-	-	-	1	-	2	5
Female	38	1	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	1	-	1	-	-	1	2
Male	14	-	-	-	-	1	-	1	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	3,088	21	7	-	2	3	8	6	28	53	118	189
Male	1,557	13	5	-	2	1	3	4	14	30	76	125
Female	1,531	8	2	-	-	2	5	2	14	23	42	64
White, both sexes	2,603	12	5	-	-	2	6	3	19	45	99	151
Male	1,277	7	4	-	-	-	3	2	9	26	62	102
Female	1,326	5	1	-	-	2	3	1	10	19	37	49
Black, both sexes	441	9	2	-	1	1	2	3	6	7	16	37
Male	253	6	1	-	1	1	-	2	3	3	12	22
Female	188	3	1	-	-	-	2	1	3	4	4	15
American Indian 1/, both sexes	24	-	-	-	1	-	-	-	2	1	3	1
Male	18	-	-	-	1	-	-	-	1	1	2	1
Female	6	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Minnesota												
All races, both sexes	1,295	8	-	-	1	2	3	1	11	14	42	67
Male	644	4	-	-	-	1	2	1	6	7	25	44
Female	651	4	-	-	1	1	1	-	5	7	17	23
White, both sexes	1,226	6	-	-	1	2	1	1	7	13	33	59
Male	601	4	-	-	-	1	1	1	5	6	19	38
Female	625	2	-	-	1	1	-	-	2	7	14	21
Black, both sexes	26	-	-	-	-	-	1	-	2	1	4	3
Male	16	-	-	-	-	-	1	-	-	1	3	2
Female	10	-	-	-	-	-	-	-	2	-	1	1
American Indian 1/, both sexes	29	1	-	-	-	-	-	-	2	-	5	4
Male	18	-	-	-	-	-	-	-	1	-	3	3
Female	11	1	-	-	-	-	-	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	1	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	953	10	5	2	-	-	8	3	17	21	29	76
Male	501	7	5	1	-	-	6	1	5	12	15	51
Female	452	3	-	1	-	-	2	2	12	9	14	25
White, both sexes	660	4	1	-	-	-	5	1	6	11	21	46
Male	347	2	1	-	-	-	4	-	4	7	11	33
Female	313	2	-	-	-	-	1	1	2	4	10	13
Black, both sexes	283	6	4	2	-	-	3	2	9	10	7	28
Male	152	5	4	1	-	-	2	1	1	5	4	17
Female	131	1	-	1	-	-	1	1	8	5	3	11
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	2	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	2	-	1	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Missouri												
All races, both sexes	1,797	10	4	1	-	3	2	3	9	16	52	105
Male	862	7	3	-	-	2	-	2	5	9	31	72
Female	935	3	1	1	-	1	2	1	4	7	21	33
White, both sexes	1,634	5	4	1	-	3	1	3	7	15	47	91
Male	780	5	3	-	-	2	-	2	4	8	29	61
Female	854	-	1	1	-	1	1	1	3	7	18	30
Black, both sexes	151	5	-	-	-	-	1	-	2	1	4	11
Male	76	2	-	-	-	-	-	-	1	1	1	9
Female	75	3	-	-	-	-	1	-	1	-	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	346	1	-	-	-	1	1	2	3	7	17	19
Male	177	-	-	-	-	-	1	-	1	3	13	12
Female	169	1	-	-	-	1	-	2	2	4	4	7
White, both sexes	301	-	-	-	-	1	1	-	-	3	10	15
Male	159	-	-	-	-	-	1	-	-	1	8	9
Female	142	-	-	-	-	1	-	-	-	2	2	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	44	1	-	-	-	-	-	2	3	4	7	3
Male	18	-	-	-	-	-	-	-	1	2	5	3
Female	26	1	-	-	-	-	-	2	2	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Nebraska												
All races, both sexes	579	3	2	1	2	-	1	1	4	3	31	28
Male	282	2	2	1	2	-	-	-	2	1	17	19
Female	297	1	-	-	-	-	1	1	2	2	14	9
White, both sexes	542	3	1	1	1	-	-	-	3	3	29	24
Male	261	2	1	1	1	-	-	-	2	1	16	17
Female	281	1	-	-	-	-	-	-	1	2	13	7
Black, both sexes	23	-	1	-	1	-	1	1	-	-	1	2
Male	14	-	1	-	1	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	1	1	-	-	-	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	768	1	1	-	-	2	3	3	6	18	48	64
Male	454	1	1	-	-	-	3	2	1	12	29	45
Female	314	-	-	-	-	2	-	1	5	6	19	19
White, both sexes	702	-	1	-	-	2	2	2	3	17	45	56
Male	417	-	1	-	-	-	2	2	1	11	26	38
Female	285	-	-	-	-	2	-	-	2	6	19	18
Black, both sexes	40	1	-	-	-	-	1	1	2	1	2	5
Male	21	1	-	-	-	-	1	-	-	1	2	5
Female	19	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
New Hampshire												
All races, both sexes	404	1	-	-	-	-	-	-	2	3	14	15
Male	193	-	-	-	-	-	-	-	2	2	10	7
Female	211	1	-	-	-	-	-	-	-	1	4	8
White, both sexes	402	1	-	-	-	-	-	-	2	3	13	15
Male	192	-	-	-	-	-	-	-	2	2	9	7
Female	210	1	-	-	-	-	-	-	-	1	4	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,384	16	4	2	2	3	6	4	14	40	87	131
Male	1,216	8	1	1	-	2	3	1	13	28	56	89
Female	1,168	8	3	1	2	1	3	3	1	12	31	42
White, both sexes	2,044	9	3	2	1	2	3	2	9	33	69	107
Male	1,045	4	1	1	-	1	2	-	8	24	44	74
Female	999	5	2	1	1	1	1	2	1	9	25	33
Black, both sexes	290	7	1	-	-	1	3	2	5	6	17	21
Male	141	4	-	-	-	1	1	1	5	3	11	12
Female	149	3	1	-	-	-	2	1	-	3	6	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	1	-	-	-	-	1	1	3
Male	28	-	-	-	-	-	-	-	-	1	1	3
Female	19	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
New Mexico												
All races, both sexes	717	5	3	-	1	2	2	7	9	18	46	62
Male	415	2	3	-	-	1	2	6	6	9	30	49
Female	302	3	-	-	1	1	-	1	3	9	16	13
White, both sexes	585	3	2	-	-	2	2	5	4	12	30	46
Male	344	2	2	-	-	1	2	4	2	8	23	37
Female	241	1	-	-	-	1	-	1	2	4	7	9
Black, both sexes	9	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	120	2	1	-	-	-	-	2	4	6	15	16
Male	67	-	1	-	-	-	-	2	3	1	7	12
Female	53	2	-	-	-	-	-	-	1	5	8	4
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,611	26	9	1	4	6	7	11	16	52	119	216
Male	2,268	21	6	1	3	3	6	5	13	37	83	159
Female	2,343	5	3	-	1	3	1	6	3	15	36	57
White, both sexes	3,881	16	7	-	1	5	5	9	14	31	101	167
Male	1,873	13	4	-	-	3	4	4	11	21	70	128
Female	2,008	3	3	-	1	2	1	5	3	10	31	39
Black, both sexes	611	10	2	1	3	1	2	1	2	14	17	39
Male	323	8	2	1	3	-	2	-	2	10	13	25
Female	288	2	-	-	-	1	-	1	-	4	4	14
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	1	1	4
Male	8	-	-	-	-	-	-	-	-	1	-	2
Female	10	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	1	-	6	-	6
Male	64	-	-	-	-	-	-	1	-	5	-	4
Female	37	-	-	-	-	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
North Carolina												
All races, both sexes	2,710	34	5	-	2	4	6	9	19	34	84	149
Male	1,343	21	-	-	1	2	3	7	9	23	50	103
Female	1,367	13	5	-	1	2	3	2	10	11	34	46
White, both sexes	2,126	18	2	-	1	3	4	9	12	22	55	116
Male	1,025	9	-	-	-	1	1	7	6	15	32	78
Female	1,101	9	2	-	1	2	3	2	6	7	23	38
Black, both sexes	537	16	3	-	1	1	1	-	6	11	26	33
Male	294	12	-	-	1	1	1	-	3	7	16	25
Female	243	4	3	-	-	-	-	-	3	4	10	8
American Indian 1/, both sexes	35	-	-	-	-	-	1	-	1	1	3	-
Male	20	-	-	-	-	-	1	-	-	1	2	-
Female	15	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	237	1	2	-	-	1	1	1	1	4	10	12
Male	121	1	2	-	-	1	1	1	1	2	6	6
Female	116	-	-	-	-	-	-	-	-	2	4	6
White, both sexes	211	1	2	-	-	1	1	1	-	2	6	11
Male	106	1	2	-	-	1	1	1	-	1	4	5
Female	105	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	2	4	1
Male	15	-	-	-	-	-	-	-	1	1	2	1
Female	11	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Ohio												
All races, both sexes	3,822	18	3	2	4	3	5	13	16	57	125	217
Male	1,871	13	2	1	2	1	3	6	7	32	79	143
Female	1,951	5	1	1	2	2	2	7	9	25	46	74
White, both sexes	3,406	13	2	2	4	1	5	10	14	47	102	173
Male	1,643	9	1	1	2	-	3	5	7	26	63	111
Female	1,763	4	1	1	2	1	2	5	7	21	39	62
Black, both sexes	397	5	1	-	-	2	-	3	2	6	20	44
Male	219	4	1	-	-	1	-	1	-	4	13	32
Female	178	1	-	-	-	1	-	2	2	2	7	12
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	3	2	-
Male	5	-	-	-	-	-	-	-	-	2	2	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	1,353	7	3	1	1	3	4	7	4	17	71	81
Male	651	4	1	-	1	-	2	1	2	14	38	48
Female	702	3	2	1	-	3	2	6	2	3	33	33
White, both sexes	1,151	6	2	-	1	1	4	4	4	12	48	59
Male	549	3	-	-	1	-	2	-	2	11	29	38
Female	602	3	2	-	-	1	2	4	2	1	19	21
Black, both sexes	67	1	1	1	-	-	-	2	-	1	5	8
Male	33	1	1	-	-	-	-	1	-	1	2	4
Female	34	-	-	1	-	-	-	1	-	-	3	4
American Indian 1/, both sexes	130	-	-	-	-	2	-	1	-	4	18	14
Male	67	-	-	-	-	-	-	-	-	2	7	6
Female	63	-	-	-	-	2	-	1	-	2	11	8
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Oregon												
All races, both sexes	1,192	8	1	-	1	2	1	3	8	20	33	85
Male	592	4	1	-	1	-	1	2	4	12	20	49
Female	600	4	-	-	-	2	-	1	4	8	13	36
White, both sexes	1,146	7	1	-	1	2	1	2	8	20	30	78
Male	568	4	1	-	1	-	1	1	4	12	18	45
Female	578	3	-	-	-	2	-	1	4	8	12	33
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	-	-	1	6
Male	12	-	-	-	-	-	-	1	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	3
Asian or Pacific Islander 2/, both sexes	18	1	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	1	-	-	-	-	-	-	-	-	1	-
Pennsylvania												
All races, both sexes	4,455	14	4	1	1	5	8	8	18	48	123	235
Male	2,086	12	4	1	1	2	5	3	12	32	76	156
Female	2,369	2	-	-	-	3	3	5	6	16	47	79
White, both sexes	4,042	9	3	1	-	5	2	6	14	42	112	187
Male	1,891	8	3	1	-	2	2	2	10	29	73	127
Female	2,151	1	-	-	-	3	-	4	4	13	39	60
Black, both sexes	395	5	1	-	1	-	6	2	4	6	11	47
Male	188	4	1	-	1	-	3	1	2	3	3	29
Female	207	1	-	-	-	-	3	1	2	3	8	18
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Rhode Island												
All races, both sexes	378	2	1	1	-	-	-	2	-	5	9	13
Male	168	2	1	1	-	-	-	1	-	5	3	7
Female	210	-	-	-	-	-	-	1	-	-	6	6
White, both sexes	363	1	1	1	-	-	-	2	-	4	9	13
Male	159	1	1	1	-	-	-	1	-	4	3	7
Female	204	-	-	-	-	-	-	1	-	-	6	6
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,499	13	3	3	2	-	2	4	8	20	58	102
Male	760	8	1	1	-	-	1	3	5	8	33	65
Female	739	5	2	2	2	-	1	1	3	12	25	37
White, both sexes	1,111	5	-	2	1	-	-	3	4	12	38	72
Male	565	3	-	-	-	-	-	2	3	5	25	46
Female	546	2	-	2	1	-	-	1	1	7	13	26
Black, both sexes	379	8	3	1	1	-	2	1	4	8	20	29
Male	189	5	1	1	-	-	1	1	2	3	8	18
Female	190	3	2	-	1	-	1	-	2	5	12	11
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
South Dakota												
All races, both sexes	280	2	-	-	-	-	1	1	3	9	10	20
Male	142	2	-	-	-	-	-	1	3	4	7	15
Female	138	-	-	-	-	-	1	-	-	5	3	5
White, both sexes	229	1	-	-	-	-	1	-	1	3	3	11
Male	112	1	-	-	-	-	-	-	1	1	2	8
Female	117	-	-	-	-	-	1	-	-	2	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	50	1	-	-	-	-	-	1	2	6	7	8
Male	29	1	-	-	-	-	-	1	2	3	5	6
Female	21	-	-	-	-	-	-	-	-	3	2	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,944	11	2	-	1	-	1	7	12	24	74	112
Male	986	5	1	-	-	-	-	6	8	15	50	73
Female	958	6	1	-	1	-	1	1	4	9	24	39
White, both sexes	1,703	3	2	-	1	-	1	6	9	20	63	92
Male	855	2	1	-	-	-	-	5	5	13	44	59
Female	848	1	1	-	1	-	1	1	4	7	19	33
Black, both sexes	232	6	-	-	-	-	-	1	2	4	11	20
Male	126	3	-	-	-	-	-	1	2	2	6	14
Female	106	3	-	-	-	-	-	-	-	2	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	2	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	3	2	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Texas												
All races, both sexes	6,213	69	8	8	4	7	14	28	41	107	285	477
Male	3,328	31	3	4	2	4	9	15	30	80	193	338
Female	2,885	38	5	4	2	3	5	13	11	27	92	139
White, both sexes	5,566	56	7	6	2	4	10	26	32	93	260	418
Male	2,979	24	3	2	1	2	6	15	22	72	178	302
Female	2,587	32	4	4	1	2	4	11	10	21	82	116
Black, both sexes	576	12	1	2	2	3	4	2	7	12	19	54
Male	308	6	-	2	1	2	3	-	6	6	11	32
Female	268	6	1	-	1	1	1	2	1	6	8	22
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	3	1
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	58	1	-	-	-	-	-	-	2	2	3	4
Male	35	1	-	-	-	-	-	-	2	2	2	4
Female	23	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	474	2	-	-	-	-	-	3	2	9	7	29
Male	244	1	-	-	-	-	-	1	2	9	5	19
Female	230	1	-	-	-	-	-	2	-	-	2	10
White, both sexes	458	2	-	-	-	-	-	3	2	7	7	26
Male	232	1	-	-	-	-	-	1	2	7	5	16
Female	226	1	-	-	-	-	-	2	-	-	2	10
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	3
Male	5	-	-	-	-	-	-	-	-	1	-	3
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Vermont												
All races, both sexes	193	-	-	-	-	1	-	1	-	2	5	8
Male	84	-	-	-	-	1	-	-	-	1	4	5
Female	109	-	-	-	-	-	-	1	-	1	1	3
White, both sexes	191	-	-	-	-	1	-	1	-	2	5	8
Male	83	-	-	-	-	1	-	-	-	1	4	5
Female	108	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,009	12	2	-	-	2	-	6	13	34	69	119
Male	1,020	6	1	-	-	1	-	3	4	19	45	85
Female	989	6	1	-	-	1	-	3	9	15	24	34
White, both sexes	1,592	6	1	-	-	2	-	3	11	25	55	90
Male	809	2	-	-	-	1	-	2	4	13	39	66
Female	783	4	1	-	-	1	-	1	7	12	16	24
Black, both sexes	397	6	1	-	-	-	-	3	1	8	14	29
Male	206	4	1	-	-	-	-	1	-	6	6	19
Female	191	2	-	-	-	-	-	2	1	2	8	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Washington												
All races, both sexes	1,666	16	3	1	-	1	2	5	8	22	54	98
Male	813	11	2	1	-	-	1	3	4	15	31	66
Female	853	5	1	-	-	1	1	2	4	7	23	32
White, both sexes	1,531	10	3	1	-	1	2	5	6	16	48	83
Male	735	7	2	1	-	-	1	3	2	9	28	55
Female	796	3	1	-	-	1	1	2	4	7	20	28
Black, both sexes	44	3	-	-	-	-	-	-	-	1	2	7
Male	27	2	-	-	-	-	-	-	-	1	1	6
Female	17	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	50	2	-	-	-	-	-	-	2	3	4	7
Male	28	1	-	-	-	-	-	-	2	3	2	4
Female	22	1	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, both sexes	41	1	-	-	-	-	-	-	-	2	-	1
Male	23	1	-	-	-	-	-	-	-	2	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	708	4	1	1	-	-	1	3	-	7	26	36
Male	343	2	-	1	-	-	1	2	-	5	17	25
Female	365	2	1	-	-	-	-	1	-	2	9	11
White, both sexes	684	3	1	1	-	-	1	3	-	7	22	34
Male	334	1	-	1	-	-	1	2	-	5	16	25
Female	350	2	1	-	-	-	-	1	-	2	6	9
Black, both sexes	24	1	-	-	-	-	-	-	-	-	4	2
Male	9	1	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
Wisconsin												
All races, both sexes	1,704	9	2	-	-	-	4	2	5	21	57	84
Male	797	4	2	-	-	-	2	-	3	12	40	59
Female	907	5	-	-	-	-	2	2	2	9	17	25
White, both sexes	1,588	7	-	-	-	-	3	2	3	19	49	67
Male	743	4	-	-	-	-	1	-	2	11	34	47
Female	845	3	-	-	-	-	2	2	1	8	15	20
Black, both sexes	73	2	1	-	-	-	-	-	-	-	4	14
Male	31	-	1	-	-	-	-	-	-	-	3	11
Female	42	2	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	33	-	1	-	-	-	1	-	1	1	4	3
Male	17	-	1	-	-	-	1	-	-	-	3	1
Female	16	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	187	1	-	-	-	-	-	1	2	2	6	17
Male	91	1	-	-	-	-	-	1	2	1	2	5
Female	96	-	-	-	-	-	-	-	-	1	4	12
White, both sexes	175	1	-	-	-	-	-	1	2	-	5	15
Male	85	1	-	-	-	-	-	1	2	-	2	4
Female	90	-	-	-	-	-	-	-	-	-	3	11
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	2	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
United States												
All races, both sexes	7,012	7,238	6,602	6,654	7,859	9,687	11,261	9,880	6,542	2,365	413	8
Male	4,906	4,868	4,116	3,707	4,060	4,563	4,635	3,395	1,860	515	80	8
Female	2,106	2,370	2,486	2,947	3,799	5,124	6,626	6,485	4,682	1,850	333	-
White, both sexes	5,718	6,086	5,571	5,683	6,892	8,616	10,206	9,115	6,046	2,136	363	7
Male	4,061	4,126	3,491	3,163	3,577	4,063	4,215	3,134	1,728	457	63	7
Female	1,657	1,960	2,080	2,520	3,315	4,553	5,991	5,981	4,318	1,679	300	-
Black, both sexes	1,043	962	819	766	764	821	819	585	391	182	42	1
Male	678	614	495	441	391	383	309	183	93	39	12	1
Female	365	348	324	325	373	438	510	402	298	143	30	-
American Indian 1/, both sexes	148	112	106	80	65	57	56	32	18	9	2	-
Male	91	74	70	35	25	24	21	11	6	3	1	-
Female	57	38	36	45	40	33	35	21	12	6	1	-
Asian or Pacific Islander 2/, both sexes	103	78	106	125	138	193	180	148	87	38	6	-
Male	76	54	60	68	67	93	90	67	33	16	4	-
Female	27	24	46	57	71	100	90	81	54	22	2	-
Alabama												
All races, both sexes	132	140	128	147	147	182	207	162	92	33	7	-
Male	89	95	80	79	69	82	86	45	17	9	3	-
Female	43	45	48	68	78	100	121	117	75	24	4	-
White, both sexes	93	106	103	116	120	154	174	137	77	24	3	-
Male	61	72	66	62	57	69	74	42	14	7	-	-
Female	32	34	37	54	63	85	100	95	63	17	3	-
Black, both sexes	39	33	25	30	27	28	33	25	15	9	4	-
Male	28	22	14	17	12	13	12	3	3	2	3	-
Female	11	11	11	13	15	15	21	22	12	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Alaska												
All races, both sexes	16	11	14	9	14	11	16	10	5	1	1	-
Male	10	8	11	3	9	5	4	3	-	1	-	-
Female	6	3	3	6	5	6	12	7	5	-	1	-
White, both sexes	12	8	10	4	12	7	12	6	4	1	1	-
Male	7	6	8	2	8	4	4	1	-	1	-	-
Female	5	2	2	2	4	3	8	5	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	4	5	1	4	3	4	1	-	-	-
Male	2	2	3	1	-	1	-	2	-	-	-	-
Female	1	1	1	4	1	3	3	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	184	159	164	147	176	165	196	156	119	33	8	1
Male	139	105	94	93	101	78	96	52	40	9	1	1
Female	45	54	70	54	75	87	100	104	79	24	7	-
White, both sexes	151	142	143	134	160	150	181	154	115	31	8	1
Male	116	95	84	87	95	72	87	51	39	7	1	1
Female	35	47	59	47	65	78	94	103	76	24	7	-
Black, both sexes	3	4	7	4	4	5	4	-	2	-	-	-
Male	1	2	4	1	2	2	4	-	1	-	-	-
Female	2	2	3	3	2	3	-	-	1	-	-	-
American Indian 1/, both sexes	29	13	13	7	11	7	11	2	2	2	-	-
Male	21	8	6	4	3	2	5	1	-	2	-	-
Female	8	5	7	3	8	5	6	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	3	-	-	-	-	-	-
Male	1	-	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Arkansas												
All races, both sexes	77	84	70	74	105	99	127	112	73	24	6	-
Male	53	52	37	45	55	41	49	39	19	3	1	-
Female	24	32	33	29	50	58	78	73	54	21	5	-
White, both sexes	66	69	61	63	93	88	117	105	66	19	3	-
Male	45	47	34	38	52	32	45	38	17	2	-	-
Female	21	22	27	25	41	56	72	67	49	17	3	-
Black, both sexes	10	14	9	9	12	11	10	7	7	5	3	-
Male	8	5	3	5	3	9	4	1	2	1	1	-
Female	2	9	6	4	9	2	6	6	5	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	909	903	761	746	756	895	1,078	900	552	207	34	3
Male	635	599	514	419	418	458	482	345	188	51	9	3
Female	274	304	247	327	338	437	596	555	364	156	25	-
White, both sexes	775	794	654	633	647	756	937	805	491	180	31	3
Male	550	537	449	352	357	386	414	312	162	45	6	3
Female	225	257	205	281	290	370	523	493	329	135	25	-
Black, both sexes	81	71	58	57	50	49	54	28	19	8	1	-
Male	49	38	34	36	31	25	22	8	7	1	1	-
Female	32	33	24	21	19	24	32	20	12	7	-	-
American Indian 1/, both sexes	13	15	12	10	9	5	5	3	2	1	-	-
Male	8	9	10	3	5	3	2	-	-	-	-	-
Female	5	6	2	7	4	2	3	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	40	23	37	46	50	85	82	64	40	18	2	-
Male	28	15	21	28	25	44	44	25	19	5	2	-
Female	12	8	16	18	25	41	38	39	21	13	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Colorado												
All races, both sexes	120	127	90	92	95	150	119	132	74	38	4	-
Male	80	83	55	53	47	71	53	46	22	11	-	-
Female	40	44	35	39	48	79	66	86	52	27	4	-
White, both sexes	111	124	79	81	94	144	116	132	72	37	4	-
Male	72	81	50	49	46	70	53	46	21	11	-	-
Female	39	43	29	32	48	74	63	86	51	26	4	-
Black, both sexes	8	-	7	4	-	3	3	-	1	1	-	-
Male	7	-	3	2	-	-	-	-	-	-	-	-
Female	1	-	4	2	-	3	3	-	1	1	-	-
American Indian 1/, both sexes	1	3	3	3	1	2	-	-	1	-	-	-
Male	1	2	2	-	1	1	-	-	1	-	-	-
Female	-	1	1	3	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	4	-	1	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	68	80	79	66	78	145	155	174	123	63	7	-
Male	36	57	46	38	37	68	59	68	30	8	2	-
Female	32	23	33	28	41	77	96	106	93	55	5	-
White, both sexes	57	73	67	62	75	136	147	171	120	63	6	-
Male	32	53	38	36	35	64	56	68	30	8	2	-
Female	25	20	29	26	40	72	91	103	90	55	4	-
Black, both sexes	11	6	10	4	3	8	7	3	1	-	1	-
Male	4	3	6	2	2	4	3	-	-	-	-	-
Female	7	3	4	2	1	4	4	3	1	-	1	-
American Indian 1/, both sexes	-	1	1	-	-	1	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Delaware												
All races, both sexes	16	28	16	14	21	28	31	30	14	3	1	-
Male	10	20	12	7	8	13	16	11	4	-	-	-
Female	6	8	4	7	13	15	15	19	10	3	1	-
White, both sexes	13	25	15	11	19	26	28	26	13	3	1	-
Male	8	18	12	6	8	13	16	9	3	-	-	-
Female	5	7	3	5	11	13	12	17	10	3	1	-
Black, both sexes	3	3	1	3	2	2	3	4	-	-	-	-
Male	2	2	-	1	-	-	-	2	-	-	-	-
Female	1	1	1	2	2	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	31	15	12	16	13	24	19	10	7	1	-
Male	16	20	10	9	12	8	9	11	2	2	-	-
Female	4	11	5	3	4	5	15	8	8	5	1	-
White, both sexes	3	3	1	3	2	3	5	4	5	3	1	-
Male	2	2	1	3	2	2	3	2	2	-	-	-
Female	1	1	-	-	-	1	2	2	3	3	1	-
Black, both sexes	16	28	13	9	14	10	17	13	5	4	-	-
Male	13	18	8	6	10	6	4	8	-	2	-	-
Female	3	10	5	3	4	4	13	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	2	1	-	-	-	-
Male	1	-	1	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Florida												
All races, both sexes	451	553	459	432	495	673	699	616	394	140	25	-
Male	322	384	294	251	262	324	310	226	109	37	8	-
Female	129	169	165	181	233	349	389	390	285	103	17	-
White, both sexes	394	494	409	390	454	628	656	579	375	129	22	-
Male	284	344	269	226	245	303	294	212	104	37	7	-
Female	110	150	140	164	209	325	362	367	271	92	15	-
Black, both sexes	50	55	48	39	38	42	38	34	16	10	3	-
Male	32	36	24	23	16	21	15	12	3	-	1	-
Female	18	19	24	16	22	21	23	22	13	10	2	-
American Indian 1/, both sexes	3	1	-	-	1	-	1	-	-	-	-	-
Male	3	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	2	3	2	3	4	3	3	1	-	-
Male	3	3	1	2	1	-	-	2	2	-	-	-
Female	1	-	1	1	1	3	4	1	1	1	-	-
Georgia												
All races, both sexes	237	247	199	201	224	249	268	220	123	41	8	-
Male	153	164	117	119	121	111	102	65	29	7	1	-
Female	84	83	82	82	103	138	166	155	94	34	7	-
White, both sexes	161	176	149	149	175	197	209	182	102	32	6	-
Male	110	120	86	91	94	88	85	58	26	4	1	-
Female	51	56	63	58	81	109	124	124	76	28	5	-
Black, both sexes	74	71	48	51	46	49	58	38	20	9	2	-
Male	42	44	30	28	25	21	16	7	3	3	-	-
Female	32	27	18	23	21	28	42	31	17	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	1	3	3	1	-	1	-	-	-
Male	1	-	1	-	2	2	1	-	-	-	-	-
Female	1	-	1	1	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Hawaii												
All races, both sexes	26	23	29	20	31	33	44	40	24	13	1	-
Male	18	15	19	15	16	17	21	14	9	7	1	-
Female	8	8	10	5	15	16	23	26	15	6	-	-
White, both sexes	11	13	16	10	8	11	16	6	4	6	-	-
Male	7	9	10	9	3	7	6	2	2	1	-	-
Female	4	4	6	1	5	4	10	4	2	5	-	-
Black, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	9	13	9	22	22	28	34	19	7	1	-
Male	11	6	9	5	13	10	15	12	7	6	1	-
Female	4	3	4	4	9	12	13	22	12	1	-	-
Idaho												
All races, both sexes	31	31	31	28	35	38	63	61	29	11	-	-
Male	20	23	18	18	17	17	25	26	10	3	-	-
Female	11	8	13	10	18	21	38	35	19	8	-	-
White, both sexes	30	30	30	28	33	37	63	61	29	11	-	-
Male	19	23	18	18	15	16	25	26	10	3	-	-
Female	11	7	12	10	18	21	38	35	19	8	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Illinois												
All races, both sexes	236	273	250	240	305	406	498	463	301	101	22	-
Male	156	181	156	143	150	170	189	146	86	13	7	-
Female	80	92	94	97	155	236	309	317	215	88	15	-
White, both sexes	185	211	203	203	274	361	459	434	273	94	18	-
Male	122	135	127	117	138	153	176	135	79	12	4	-
Female	63	76	76	86	136	208	283	299	194	82	14	-
Black, both sexes	47	58	46	33	25	38	33	26	27	7	3	-
Male	33	42	28	23	10	13	12	8	7	1	3	-
Female	14	16	18	10	15	25	21	18	20	6	-	-
American Indian 1/, both sexes	4	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	2	6	7	6	3	1	-	1	-
Male	-	4	1	1	2	4	1	3	-	-	-	-
Female	-	-	-	1	4	3	5	-	1	-	1	-
Indiana												
All races, both sexes	150	133	142	148	164	213	237	218	140	41	7	-
Male	101	92	78	72	78	95	98	70	35	13	2	-
Female	49	41	64	76	86	118	139	148	105	28	5	-
White, both sexes	121	122	129	139	155	196	225	210	133	41	6	-
Male	79	86	71	68	75	89	93	67	34	13	2	-
Female	42	36	58	71	80	107	132	143	99	28	4	-
Black, both sexes	29	11	12	8	9	14	12	8	7	-	1	-
Male	22	6	6	4	3	5	5	3	1	-	-	-
Female	7	5	6	4	6	9	7	5	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Iowa												
All races, both sexes	57	36	52	61	68	113	135	122	113	33	7	-
Male	41	24	30	31	31	57	53	36	34	6	-	-
Female	16	12	22	30	37	56	82	86	79	27	7	-
White, both sexes	51	33	50	60	66	109	134	121	113	32	7	-
Male	36	23	30	30	31	56	53	35	34	6	-	-
Female	15	10	20	30	35	53	81	86	79	26	7	-
Black, both sexes	3	1	2	-	1	3	1	1	-	1	-	-
Male	3	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	2	-	1	2	1	-	-	1	-	-
American Indian 1/, both sexes	3	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	57	63	45	61	54	98	120	118	99	30	7	-
Male	40	36	31	40	24	49	47	46	23	7	3	-
Female	17	27	14	21	30	49	73	72	76	23	4	-
White, both sexes	56	59	40	52	50	93	115	115	98	29	7	-
Male	39	35	28	34	22	47	45	45	23	6	3	-
Female	17	24	12	18	28	46	70	70	75	23	4	-
Black, both sexes	-	4	3	4	3	2	3	1	1	1	-	-
Male	-	1	2	3	1	2	1	-	-	1	-	-
Female	-	3	1	1	2	-	2	1	1	-	-	-
American Indian 1/, both sexes	1	-	1	3	1	3	-	1	-	-	-	-
Male	1	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	3	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	2	1	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Kentucky												
All races, both sexes	104	101	114	115	115	144	176	143	94	24	4	-
Male	78	67	61	62	60	70	62	53	29	5	2	-
Female	26	34	53	53	55	74	114	90	65	19	2	-
White, both sexes	93	95	108	108	107	135	170	134	88	24	4	-
Male	67	62	57	59	56	64	59	50	28	5	2	-
Female	26	33	51	49	51	71	111	84	60	19	2	-
Black, both sexes	10	4	5	7	8	9	5	8	6	-	-	-
Male	10	3	3	3	4	6	2	3	1	-	-	-
Female	-	1	2	4	4	3	3	5	5	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	121	113	123	124	134	159	158	135	73	30	3	-
Male	85	81	77	60	67	76	61	38	14	7	1	-
Female	36	32	46	64	67	83	97	97	59	23	2	-
White, both sexes	80	75	96	84	96	119	122	107	58	21	3	-
Male	60	54	59	41	47	57	47	28	11	3	1	-
Female	20	21	37	43	49	62	75	79	47	18	2	-
Black, both sexes	41	35	27	40	37	39	33	26	15	9	-	-
Male	25	24	18	19	20	18	14	10	3	4	-	-
Female	16	11	9	21	17	21	19	16	12	5	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	3	1	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Maine												
All races, both sexes	36	31	36	30	33	64	74	75	48	17	4	-
Male	32	22	22	14	15	37	28	37	12	6	3	-
Female	4	9	14	16	18	27	46	38	36	11	1	-
White, both sexes	35	31	35	30	33	64	74	74	48	17	4	-
Male	31	22	22	14	15	37	28	36	12	6	3	-
Female	4	9	13	16	18	27	46	38	36	11	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	113	131	128	119	125	170	204	181	104	39	6	1
Male	87	92	83	70	53	84	87	59	30	5	2	1
Female	26	39	45	49	72	86	117	122	74	34	4	-
White, both sexes	69	88	76	84	100	133	169	158	86	32	4	-
Male	52	61	46	47	43	68	76	53	25	4	1	-
Female	17	27	30	37	57	65	93	105	61	28	3	-
Black, both sexes	41	42	49	32	24	34	35	21	17	7	2	1
Male	33	31	35	22	10	16	11	5	5	1	1	1
Female	8	11	14	10	14	18	24	16	12	6	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	3	1	3	-	2	-	-	-	-
Male	2	-	2	1	-	-	-	1	-	-	-	-
Female	-	1	1	2	1	3	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Massachusetts												
All races, both sexes	156	146	125	127	190	254	310	273	205	89	15	-
Male	111	88	72	73	104	127	125	98	69	21	2	-
Female	45	58	53	54	86	127	185	175	136	68	13	-
White, both sexes	150	134	117	122	174	235	302	269	201	84	15	-
Male	106	81	68	70	99	115	121	97	68	18	2	-
Female	44	53	49	52	75	120	181	172	133	66	13	-
Black, both sexes	4	12	8	5	12	12	7	4	2	2	-	-
Male	4	7	4	3	5	7	3	1	-	1	-	-
Female	-	5	4	2	7	5	4	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	4	7	1	-	2	3	-	-
Male	1	-	-	-	-	5	1	-	1	2	-	-
Female	1	-	-	-	4	2	-	-	1	1	-	-
Michigan												
All races, both sexes	272	267	231	222	299	321	389	336	229	74	13	-
Male	187	190	142	122	155	146	146	121	61	12	2	-
Female	85	77	89	100	144	175	243	215	168	62	11	-
White, both sexes	210	204	194	179	255	281	349	304	204	68	13	-
Male	144	147	116	93	131	128	127	111	53	10	2	-
Female	66	57	78	86	124	153	222	193	151	58	11	-
Black, both sexes	56	58	31	39	40	40	34	28	25	6	-	-
Male	37	39	22	28	21	18	19	8	8	2	-	-
Female	19	19	9	11	19	22	15	20	17	4	-	-
American Indian 1/, both sexes	5	3	5	1	1	-	1	-	-	-	-	-
Male	5	2	4	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	3	3	-	5	4	-	-	-	-
Male	1	2	-	1	2	-	-	2	-	-	-	-
Female	-	-	1	2	1	-	5	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Minnesota												
All races, both sexes	83	82	83	103	102	111	181	175	164	52	10	-
Male	62	55	53	49	58	57	91	62	52	13	2	-
Female	21	27	30	54	44	54	90	113	112	39	8	-
White, both sexes	75	75	75	100	98	104	178	173	163	52	10	-
Male	56	51	45	49	56	54	88	61	51	13	2	-
Female	19	24	30	51	42	50	90	112	112	39	8	-
Black, both sexes	4	4	3	1	-	1	2	-	-	-	-	-
Male	3	1	3	-	-	-	2	-	-	-	-	-
Female	1	3	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	5	1	2	4	-	1	-	-	-	-
Male	2	1	5	-	1	2	-	-	-	-	-	-
Female	1	-	-	1	1	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	1	2	2	1	1	1	-	-	-
Male	1	2	-	-	1	1	1	1	1	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	82	93	67	75	97	110	103	87	37	29	2	-
Male	64	68	39	41	52	50	38	25	13	8	-	-
Female	18	25	28	34	45	60	65	62	24	21	2	-
White, both sexes	49	69	51	48	76	81	83	63	29	16	-	-
Male	38	51	30	27	39	38	31	17	10	4	-	-
Female	11	18	21	21	37	43	52	46	19	12	-	-
Black, both sexes	31	24	16	26	20	28	20	24	8	13	2	-
Male	25	17	9	14	13	12	7	8	3	4	-	-
Female	6	7	7	12	7	16	13	16	5	9	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Missouri												
All races, both sexes	123	105	116	124	184	213	262	233	167	52	13	-
Male	90	67	60	59	103	110	113	74	40	13	2	-
Female	33	38	56	65	81	103	149	159	127	39	11	-
White, both sexes	104	88	96	111	169	197	248	227	154	51	12	-
Male	79	55	54	54	94	99	107	71	39	13	1	-
Female	25	33	42	57	75	98	141	156	115	38	11	-
Black, both sexes	18	17	18	12	14	14	13	6	13	1	1	-
Male	11	12	6	5	8	10	5	3	1	-	1	-
Female	7	5	12	7	6	4	8	3	12	1	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	37	26	15	28	25	37	40	42	36	9	-	-
Male	24	15	12	15	11	22	14	20	13	1	-	-
Female	13	11	3	13	14	15	26	22	23	8	-	-
White, both sexes	29	23	14	24	25	34	37	41	35	9	-	-
Male	22	14	11	13	11	21	14	20	13	1	-	-
Female	7	9	3	11	14	13	23	21	22	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	3	1	4	-	3	3	1	1	-	-	-
Male	2	1	1	2	-	1	-	-	-	-	-	-
Female	6	2	-	2	-	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Nebraska												
All races, both sexes	32	37	33	38	43	66	87	73	53	39	2	-
Male	26	26	19	22	24	34	41	23	12	9	-	-
Female	6	11	14	16	19	32	46	50	41	30	2	-
White, both sexes	28	30	31	35	40	63	85	73	52	38	2	-
Male	23	21	19	21	21	31	41	23	11	9	-	-
Female	5	9	12	14	19	32	44	50	41	29	2	-
Black, both sexes	1	3	2	2	3	2	1	-	1	1	-	-
Male	1	3	-	-	3	2	-	-	1	-	-	-
Female	-	-	2	2	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	3	4	-	1	-	1	-	-	-	-	-	-
Male	2	2	-	1	-	1	-	-	-	-	-	-
Female	1	2	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Nevada												
All races, both sexes	67	79	84	73	74	79	80	53	26	5	2	-
Male	42	53	59	46	40	44	42	25	5	2	2	-
Female	25	26	25	27	34	35	38	28	21	3	-	-
White, both sexes	62	72	77	65	69	72	74	51	25	5	2	-
Male	40	49	55	40	38	42	38	25	5	2	2	-
Female	22	23	22	25	31	30	36	26	20	3	-	-
Black, both sexes	4	5	4	3	3	4	3	1	-	-	-	-
Male	1	2	3	2	-	2	1	-	-	-	-	-
Female	3	3	1	1	3	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	1	2	3	1	1	-	-	-
Male	1	2	1	3	1	-	3	-	-	-	-	-
Female	-	-	1	1	-	2	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
New Hampshire												
All races, both sexes	27	24	25	25	34	58	69	57	31	16	3	-
Male	21	13	13	15	14	25	30	27	11	3	-	-
Female	6	11	12	10	20	33	39	30	20	13	3	-
White, both sexes	27	23	25	25	34	58	69	57	31	16	3	-
Male	21	13	13	15	14	25	30	27	11	3	-	-
Female	6	10	12	10	20	33	39	30	20	13	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	182	174	163	165	236	263	338	283	192	69	10	-
Male	127	115	104	94	127	123	153	92	60	16	3	-
Female	55	59	59	71	109	140	185	191	132	53	7	-
White, both sexes	142	149	134	136	202	233	302	260	176	63	7	-
Male	102	101	87	78	111	111	135	87	57	14	3	-
Female	40	48	47	58	91	122	167	173	119	49	4	-
Black, both sexes	36	24	25	20	27	24	31	20	13	4	3	-
Male	22	13	14	8	13	9	16	4	3	1	-	-
Female	14	11	11	12	14	15	15	16	10	3	3	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	4	9	6	5	4	3	3	2	-	-
Male	3	1	3	8	2	2	2	1	-	1	-	-
Female	1	-	1	1	4	3	2	2	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
New Mexico												
All races, both sexes	75	67	50	60	63	64	82	52	33	16	-	-
Male	56	40	36	37	36	32	40	18	7	5	-	-
Female	19	27	14	23	27	32	42	34	26	11	-	-
White, both sexes	55	56	40	51	51	58	76	48	31	13	-	-
Male	45	33	31	32	33	29	35	16	5	4	-	-
Female	10	23	9	19	18	29	41	32	26	9	-	-
Black, both sexes	1	-	1	1	-	-	2	2	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	1	-	1	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	19	11	9	8	11	6	4	2	2	2	-	-
Male	11	7	5	4	3	3	3	2	2	1	-	-
Female	8	4	4	4	8	3	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
New York												
All races, both sexes	317	323	351	365	456	577	594	598	390	147	26	-
Male	222	218	224	201	209	273	243	196	120	24	1	-
Female	95	105	127	164	247	304	351	402	270	123	25	-
White, both sexes	234	266	272	297	383	491	521	543	359	135	24	-
Male	168	180	169	162	171	238	213	175	115	23	1	-
Female	66	86	103	135	212	253	308	368	244	112	23	-
Black, both sexes	75	52	65	57	65	65	60	41	27	10	2	-
Male	47	35	45	32	36	22	22	13	4	1	-	-
Female	28	17	20	25	29	43	38	28	23	9	2	-
American Indian 1/, both sexes	1	2	2	2	1	-	3	1	-	-	-	-
Male	1	2	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	2	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	12	9	7	21	10	13	4	2	-	-
Male	6	1	9	7	2	13	7	8	1	-	-	-
Female	1	2	3	2	5	8	3	5	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
North Carolina												
All races, both sexes	218	226	227	236	278	312	335	278	189	58	7	-
Male	149	145	145	148	128	152	120	74	54	9	-	-
Female	69	81	82	88	150	160	215	204	135	49	7	-
White, both sexes	154	168	173	181	223	265	274	235	160	45	6	-
Male	115	110	106	113	101	123	97	61	45	5	-	-
Female	39	58	67	68	122	142	177	174	115	40	6	-
Black, both sexes	54	54	46	50	53	44	57	39	28	13	1	-
Male	29	33	33	33	27	28	22	11	8	4	-	-
Female	25	21	13	17	26	16	35	28	20	9	1	-
American Indian 1/, both sexes	9	3	5	5	1	-	3	2	1	-	-	-
Male	5	1	4	2	-	-	1	2	1	-	-	-
Female	4	2	1	3	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	-	1	3	1	2	-	-	-	-
Male	-	1	2	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	1	2	1	2	-	-	-	-
North Dakota												
All races, both sexes	12	17	13	15	15	21	25	33	34	17	2	-
Male	9	13	9	11	7	10	16	11	12	2	-	-
Female	3	4	4	4	8	11	9	22	22	15	2	-
White, both sexes	9	15	11	14	13	20	21	31	34	16	2	-
Male	6	12	8	10	7	10	12	11	12	2	-	-
Female	3	3	3	4	6	10	9	20	22	14	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	2	2	1	2	1	4	2	-	1	-	-
Male	3	1	1	1	-	-	4	-	-	-	-	-
Female	-	1	1	-	2	1	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Ohio												
All races, both sexes	303	283	274	291	358	444	535	472	286	105	8	-
Male	208	187	164	157	187	200	212	166	76	24	1	-
Female	95	96	110	134	171	244	323	306	210	81	7	-
White, both sexes	257	248	240	258	324	402	496	448	264	89	7	-
Male	174	166	141	139	168	181	198	154	71	22	1	-
Female	83	82	99	119	156	221	298	294	193	67	6	-
Black, both sexes	45	35	32	31	33	39	37	24	21	16	1	-
Male	33	21	22	17	19	18	14	12	5	2	-	-
Female	12	14	10	14	14	21	23	12	16	14	1	-
American Indian 1/, both sexes	1	-	1	-	1	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	2	2	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	2	2	-	1	-	-	-
Oklahoma												
All races, both sexes	97	139	115	104	118	133	159	135	109	39	6	-
Male	66	92	67	56	52	64	59	49	29	6	-	-
Female	31	47	48	48	66	69	100	86	80	33	6	-
White, both sexes	82	107	97	84	106	128	144	123	99	35	5	-
Male	59	69	55	45	46	62	54	43	25	5	-	-
Female	23	38	42	39	60	66	90	80	74	30	5	-
Black, both sexes	4	8	5	8	2	2	7	4	5	2	1	-
Male	3	4	2	5	1	1	1	2	3	1	-	-
Female	1	4	3	3	1	1	6	2	2	1	1	-
American Indian 1/, both sexes	11	24	12	12	10	3	6	8	3	2	-	-
Male	4	19	9	6	5	1	3	4	1	-	-	-
Female	7	5	3	6	5	2	3	4	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	2	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Oregon												
All races, both sexes	91	102	95	84	89	116	152	162	93	41	5	-
Male	65	71	59	41	42	61	58	60	31	10	-	-
Female	26	31	36	43	47	55	94	102	62	31	5	-
White, both sexes	89	97	89	79	87	111	148	158	92	41	5	-
Male	65	68	54	39	42	59	57	57	30	10	-	-
Female	24	29	35	40	45	52	91	101	62	31	5	-
Black, both sexes	-	-	-	-	-	3	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	2	2	2	3	-	1	2	-	-	-	-	-
Male	-	2	2	2	-	1	-	-	-	-	-	-
Female	2	-	-	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	4	2	2	1	1	2	-	-	-	-
Male	-	1	3	-	-	-	1	2	-	-	-	-
Female	-	2	1	2	2	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	264	292	237	282	415	584	710	639	401	131	33	2
Male	186	193	144	156	228	260	284	192	111	19	7	2
Female	78	99	93	126	187	324	426	447	290	112	26	-
White, both sexes	218	252	198	252	390	536	669	611	380	122	31	2
Male	154	169	128	141	218	242	269	184	103	17	7	2
Female	64	83	70	111	172	294	400	427	277	105	24	-
Black, both sexes	43	40	36	27	25	47	38	27	19	8	2	-
Male	29	24	16	14	10	18	14	7	7	2	-	-
Female	14	16	20	13	15	29	24	20	12	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	3	3	-	1	3	1	2	1	-	-
Male	3	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	3	2	-	1	2	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Rhode Island												
All races, both sexes	28	24	20	22	36	49	68	42	40	13	3	-
Male	16	17	11	12	15	22	30	15	8	1	1	-
Female	12	7	9	10	21	27	38	27	32	12	2	-
White, both sexes	25	22	18	20	35	46	68	42	40	13	3	-
Male	14	15	11	11	15	20	30	15	8	1	1	-
Female	11	7	7	9	20	26	38	27	32	12	2	-
Black, both sexes	-	2	-	1	1	2	-	-	-	-	-	-
Male	-	2	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	140	134	132	125	164	159	177	139	81	29	4	-
Male	104	89	80	68	88	71	61	45	23	6	-	-
Female	36	45	52	57	76	88	116	94	58	23	4	-
White, both sexes	94	94	97	89	128	127	144	111	64	23	3	-
Male	72	63	62	47	64	58	52	39	19	5	-	-
Female	22	31	35	42	64	69	92	72	45	18	3	-
Black, both sexes	45	39	33	35	36	31	32	28	17	5	1	-
Male	31	25	18	20	24	13	8	6	4	-	-	-
Female	14	14	15	15	12	18	24	22	13	5	1	-
American Indian 1/, both sexes	1	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
South Dakota												
All races, both sexes	17	13	16	13	13	35	45	42	20	17	3	-
Male	12	8	12	7	5	17	23	16	6	4	-	-
Female	5	5	4	6	8	18	22	26	14	13	3	-
White, both sexes	13	9	10	12	10	32	42	41	20	17	3	-
Male	9	6	8	6	5	16	23	16	6	4	-	-
Female	4	3	2	6	5	16	19	25	14	13	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	4	6	1	3	3	3	1	-	-	-	-
Male	3	2	4	1	-	1	-	-	-	-	-	-
Female	1	2	2	-	3	2	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	154	175	168	152	173	222	253	218	137	40	8	-
Male	116	128	115	83	82	99	95	71	33	5	1	-
Female	38	47	53	69	91	123	158	147	104	35	7	-
White, both sexes	128	155	144	129	149	203	233	200	127	32	6	-
Male	97	112	96	71	70	89	87	67	32	4	1	-
Female	31	43	48	58	79	114	146	133	95	28	5	-
Black, both sexes	25	17	24	23	23	19	20	18	9	8	2	-
Male	18	13	19	12	12	10	8	4	1	1	-	-
Female	7	4	5	11	11	9	12	14	8	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	-	1	-	-	-	-	-	-	-
Male	1	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Texas												
All races, both sexes	599	582	522	504	591	646	653	527	378	127	36	-
Male	435	381	337	267	331	273	285	175	102	30	3	-
Female	164	201	185	237	260	373	368	352	276	97	33	-
White, both sexes	525	508	466	452	531	583	597	486	359	114	31	-
Male	393	336	298	239	297	242	262	159	98	26	2	-
Female	132	172	168	213	234	341	335	327	261	88	29	-
Black, both sexes	68	69	49	44	50	60	49	39	17	11	2	-
Male	38	43	35	26	27	28	20	15	4	3	-	-
Female	30	26	14	18	23	32	29	24	13	8	2	-
American Indian 1/, both sexes	3	1	2	2	-	-	-	-	-	-	1	-
Male	2	-	1	1	-	-	-	-	-	-	-	-
Female	1	1	1	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	4	5	6	10	3	7	2	2	2	2	-
Male	2	2	3	1	7	3	3	1	-	1	1	-
Female	1	2	2	5	3	-	4	1	2	1	1	-
Utah												
All races, both sexes	26	30	41	43	50	49	78	57	40	6	1	1
Male	21	21	24	25	25	23	36	18	9	3	1	1
Female	5	9	17	18	25	26	42	39	31	3	-	-
White, both sexes	26	29	40	41	49	46	76	56	40	6	1	1
Male	21	20	23	25	24	22	34	17	9	3	1	1
Female	5	9	17	16	25	24	42	39	31	3	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	1	1	-	-	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Vermont												
All races, both sexes	10	10	12	11	14	24	27	25	23	17	3	-
Male	5	8	7	4	9	10	10	10	6	3	1	-
Female	5	2	5	7	5	14	17	15	17	14	2	-
White, both sexes	10	10	12	10	14	23	27	25	23	17	3	-
Male	5	8	7	4	9	9	10	10	6	3	1	-
Female	5	2	5	6	5	14	17	15	17	14	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	152	154	153	161	190	221	287	235	145	49	5	-
Male	101	109	98	88	97	104	109	96	41	11	2	-
Female	51	45	55	73	93	117	178	139	104	38	3	-
White, both sexes	105	112	115	122	142	187	237	209	127	40	3	-
Male	76	79	74	66	75	89	91	84	37	10	1	-
Female	29	33	41	56	67	98	146	125	90	30	2	-
Black, both sexes	45	41	36	36	46	30	47	25	18	9	2	-
Male	24	29	23	22	21	15	17	12	4	1	1	-
Female	21	12	13	14	25	15	30	13	14	8	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	3	2	4	3	1	-	-	-	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	3	1	4	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Washington												
All races, both sexes	112	146	117	121	146	169	229	187	159	58	12	-
Male	70	100	73	55	71	79	93	69	48	19	2	-
Female	42	46	44	66	75	90	136	118	111	39	10	-
White, both sexes	97	129	104	115	132	159	221	177	153	58	11	-
Male	59	87	68	53	64	74	91	63	48	19	1	-
Female	38	42	36	62	68	85	130	114	105	39	10	-
Black, both sexes	6	6	2	4	2	3	3	2	3	-	-	-
Male	4	6	-	-	1	3	1	2	-	-	-	-
Female	2	-	2	4	1	-	2	-	3	-	-	-
American Indian 1/, both sexes	4	6	10	-	4	2	2	1	2	-	1	-
Male	3	4	5	-	2	1	-	-	-	-	1	-
Female	1	2	5	-	2	1	2	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	1	2	8	5	3	7	1	-	-	-
Male	4	3	-	2	4	1	1	4	-	-	-	-
Female	1	2	1	-	4	4	2	3	1	-	-	-
West Virginia												
All races, both sexes	44	63	52	65	65	95	108	73	45	14	5	-
Male	23	51	27	39	32	50	37	19	9	3	-	-
Female	21	12	25	26	33	45	71	54	36	11	5	-
White, both sexes	43	60	50	63	64	93	108	69	45	13	4	-
Male	22	49	27	38	32	49	37	17	9	3	-	-
Female	21	11	23	25	32	44	71	52	36	10	4	-
Black, both sexes	1	3	2	2	1	2	-	4	-	1	1	-
Male	1	2	-	1	-	1	-	2	-	-	-	-
Female	-	1	2	1	1	1	-	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Wisconsin												
All races, both sexes	114	110	105	119	138	185	239	244	179	74	13	-
Male	73	77	54	67	71	88	87	84	51	22	1	-
Female	41	33	51	52	67	97	152	160	128	52	12	-
White, both sexes	95	97	92	111	134	180	231	240	174	72	12	-
Male	64	67	50	65	71	85	85	83	51	22	1	-
Female	31	30	42	46	63	95	146	157	123	50	11	-
Black, both sexes	10	7	10	3	4	2	6	4	4	1	1	-
Male	4	5	2	1	-	1	2	1	-	-	-	-
Female	6	2	8	2	4	1	4	3	4	1	1	-
American Indian 1/, both sexes	7	3	2	4	-	3	2	-	-	1	-	-
Male	3	3	2	1	-	2	-	-	-	-	-	-
Female	4	-	-	3	-	1	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	1	-	-	-	-	1	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	17	17	15	15	12	22	17	23	16	4	-	-
Male	14	10	12	8	7	6	7	8	6	1	-	-
Female	3	7	3	7	5	16	10	15	10	3	-	-
White, both sexes	16	14	15	14	12	21	17	23	15	4	-	-
Male	13	9	12	8	7	5	7	8	5	1	-	-
Female	3	5	3	6	5	16	10	15	10	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	-	1	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
United States												
All races, both sexes	241	2	-	2	1	-	2	1	3	1	9	4
Male	116	1	-	1	1	-	2	-	3	1	6	3
Female	125	1	-	1	-	-	-	1	-	-	3	1
White, both sexes	212	2	-	-	1	-	2	-	3	1	6	2
Male	100	1	-	-	1	-	2	-	3	1	3	2
Female	112	1	-	-	-	-	-	-	-	-	3	-
Black, both sexes	24	-	-	2	-	-	-	1	-	-	3	2
Male	11	-	-	1	-	-	-	-	-	-	3	1
Female	13	-	-	1	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Arkansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	27	1	-	-	-	-	-	-	2	-	1	1
Male	15	1	-	-	-	-	-	-	2	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	24	1	-	-	-	-	-	-	2	-	1	1
Male	13	1	-	-	-	-	-	-	2	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Colorado												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Florida												
All races, both sexes	13	-	-	1	-	-	-	-	1	-	1	-
Male	8	-	-	1	-	-	-	-	1	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Illinois												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Iowa												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Kentucky												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Massachusetts												
All races, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Missouri												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	12	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
North Carolina												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Ohio												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Oregon												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	16	-	-	-	-	-	1	-	-	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Texas												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Washington												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Wisconsin												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
United States												
All races, both sexes	17	10	10	13	21	29	35	36	25	14	6	-
Male	9	6	4	6	15	16	17	13	7	3	2	-
Female	8	4	6	7	6	13	18	23	18	11	4	-
White, both sexes	9	9	9	12	20	26	30	36	24	14	6	-
Male	6	5	4	6	14	14	13	13	7	3	2	-
Female	3	4	5	6	6	12	17	23	17	11	4	-
Black, both sexes	7	1	1	1	-	2	3	-	1	-	-	-
Male	2	1	-	-	-	1	2	-	-	-	-	-
Female	5	-	1	1	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	2	-	-	-	-	-
Male	1	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	-	1	1	-	-	1	-	-	-
Male	2	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Arkansas												
All races, both sexes	-	-	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	1	-	3	2	3	4	2	4	1	1	-
Male	-	1	-	1	2	2	1	2	1	1	-	-
Female	1	-	-	2	-	1	3	-	3	-	1	-
White, both sexes	-	-	-	3	2	3	3	2	4	1	1	-
Male	-	-	-	1	2	2	-	2	1	1	-	-
Female	-	-	-	2	-	1	3	-	3	-	1	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Colorado												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Delaware												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Florida												
All races, both sexes	1	-	-	1	1	1	-	4	-	2	-	-
Male	-	-	-	1	1	1	-	2	-	-	-	-
Female	1	-	-	-	-	-	-	2	-	2	-	-
White, both sexes	-	-	-	1	1	1	-	4	-	2	-	-
Male	-	-	-	1	1	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	1	-	1	-	1	1	1	-	-	-
Male	1	-	1	-	1	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	1	1	-	1	-	-	1	1	-	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Illinois												
All races, both sexes	-	-	-	-	1	2	2	2	2	-	1	-
Male	-	-	-	-	-	-	1	-	2	-	1	-
Female	-	-	-	-	1	2	1	2	-	-	-	-
White, both sexes	-	-	-	-	1	2	2	2	2	-	1	-
Male	-	-	-	-	-	-	1	-	2	-	1	-
Female	-	-	-	-	1	2	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Iowa												
All races, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Kentucky												
All races, both sexes	-	-	-	-	-	1	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	1	3	-	1	-	-	-
Male	-	-	-	-	-	1	3	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Massachusetts												
All races, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	1	-	2	-	1	3	1	-	-	-
Male	-	-	-	-	2	-	1	-	1	-	-	-
Female	-	1	1	-	-	-	-	3	-	-	-	-
White, both sexes	-	1	1	-	1	-	1	3	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	1	1	-	-	-	-	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Minnesota												
All races, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Missouri												
All races, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Nebraska												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	2	-	-	1	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	1	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	3	-	1	-	1	1	-	1	1	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	2	-	2	-	-	-	1	1	-	-	1	-
White, both sexes	1	-	2	-	1	-	1	1	-	1	1	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	1	-	1	-	-	-	1	1	-	-	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
North Carolina												
All races, both sexes	-	1	1	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Ohio												
All races, both sexes	1	1	-	-	-	-	-	2	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Oregon												
All races, both sexes	-	2	-	1	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	2	-	1	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	1	1	5	1	3	3	1	-	-
Male	-	-	-	1	1	4	-	-	1	1	-	-
Female	-	-	-	-	-	1	1	3	2	-	-	-
White, both sexes	-	-	-	1	1	4	1	3	3	1	-	-
Male	-	-	-	1	1	4	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	3	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Rhode Island												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	1	1	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	1	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	1	2	-	-	-	-
Male	2	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	2	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Texas												
All races, both sexes	2	-	3	1	5	2	4	4	1	3	-	-
Male	1	-	1	1	3	2	1	3	-	-	-	-
Female	1	-	2	-	2	-	3	1	1	3	-	-
White, both sexes	1	-	3	1	5	2	4	4	1	3	-	-
Male	1	-	1	1	3	2	1	3	-	-	-	-
Female	-	-	2	-	2	-	3	1	1	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	-	1	-	-	2	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	2	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Washington												
All races, both sexes	-	-	-	-	1	1	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
White, both sexes	-	-	-	-	1	1	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Wisconsin												
All races, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
United States												
All races, both sexes	87,150	624	131	53	58	89	157	291	554	1,304	3,177	5,407
Male	44,282	387	80	30	35	44	88	170	333	818	2,032	3,650
Female	42,868	237	51	23	23	45	69	121	221	486	1,145	1,757
White, both sexes	75,652	388	93	39	31	66	104	224	405	1,046	2,611	4,401
Male	38,170	238	54	19	17	33	58	136	253	670	1,697	2,997
Female	37,482	150	39	20	14	33	46	88	152	376	914	1,404
Black, both sexes	9,064	213	30	12	23	18	50	49	98	179	403	810
Male	4,790	140	20	9	17	10	28	22	48	101	236	526
Female	4,274	73	10	3	6	8	22	27	50	78	167	284
American Indian 1/, both sexes	1,049	10	5	-	1	2	2	13	32	52	118	129
Male	564	3	3	-	1	-	2	8	19	25	66	76
Female	485	7	2	-	-	2	-	5	13	27	52	53
Asian or Pacific Islander 2/, both sexes	1,385	13	3	2	3	3	1	5	19	27	45	67
Male	758	6	3	2	-	1	-	4	13	22	33	51
Female	627	7	-	-	3	2	1	1	6	5	12	16
Alabama												
All races, both sexes	1,606	17	-	2	1	3	2	5	14	22	61	104
Male	797	11	-	2	-	1	2	1	7	12	37	70
Female	809	6	-	-	1	2	-	4	7	10	24	34
White, both sexes	1,253	5	-	1	-	-	1	3	9	15	41	73
Male	625	5	-	1	-	-	1	1	6	10	24	53
Female	628	-	-	-	-	-	-	2	3	5	17	20
Black, both sexes	349	12	-	1	1	3	1	2	5	7	18	31
Male	169	6	-	1	-	1	1	-	1	2	11	17
Female	180	6	-	-	1	2	-	2	4	5	7	14
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Alaska												
All races, both sexes	134	2	-	-	-	-	-	-	-	4	11	9
Male	62	1	-	-	-	-	-	-	-	1	4	2
Female	72	1	-	-	-	-	-	-	-	3	7	7
White, both sexes	94	1	-	-	-	-	-	-	-	4	7	5
Male	48	1	-	-	-	-	-	-	-	1	3	2
Female	46	-	-	-	-	-	-	-	-	3	4	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	36	1	-	-	-	-	-	-	-	-	3	4
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	-	-	-	-	3	4
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,802	18	5	1	-	1	2	9	16	42	82	123
Male	998	10	3	-	-	-	2	7	13	28	54	76
Female	804	8	2	1	-	1	-	2	3	14	28	47
White, both sexes	1,589	17	4	1	-	1	2	3	9	32	62	91
Male	880	9	3	-	-	-	2	2	9	23	40	59
Female	709	8	1	1	-	1	-	1	-	9	22	32
Black, both sexes	42	1	-	-	-	-	-	1	-	1	3	3
Male	22	1	-	-	-	-	-	1	-	-	2	1
Female	20	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	161	-	1	-	-	-	-	5	6	9	17	26
Male	89	-	-	-	-	-	-	4	3	5	12	13
Female	72	-	1	-	-	-	-	1	3	4	5	13
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	3
Male	7	-	-	-	-	-	-	-	1	-	-	3
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Arkansas												
All races, both sexes	955	6	3	-	1	-	3	1	8	14	23	48
Male	453	6	1	-	1	-	2	-	3	5	10	33
Female	502	-	2	-	-	-	1	1	5	9	13	15
White, both sexes	832	3	3	-	1	-	1	-	4	12	18	43
Male	398	3	1	-	1	-	-	-	2	4	8	31
Female	434	-	2	-	-	-	1	-	2	8	10	12
Black, both sexes	118	3	-	-	-	-	2	1	4	2	4	5
Male	52	3	-	-	-	-	2	-	1	1	1	2
Female	66	-	-	-	-	-	-	1	3	1	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9,250	71	17	7	12	14	14	47	73	194	409	670
Male	5,142	41	12	3	7	9	7	34	50	131	272	466
Female	4,108	30	5	4	5	5	7	13	23	63	137	204
White, both sexes	7,987	54	14	7	7	12	11	39	61	175	347	573
Male	4,443	30	10	3	5	8	5	29	40	120	234	395
Female	3,544	24	4	4	2	4	6	10	21	55	113	178
Black, both sexes	611	15	1	-	3	1	3	6	2	10	33	63
Male	341	10	1	-	2	1	2	4	2	5	19	44
Female	270	5	-	-	1	-	1	2	-	5	14	19
American Indian 1/, both sexes	102	-	1	-	-	-	-	1	4	3	9	9
Male	59	-	-	-	-	-	-	-	4	2	5	8
Female	43	-	1	-	-	-	-	1	-	1	4	1
Asian or Pacific Islander 2/, both sexes	550	2	1	-	2	1	-	1	6	6	20	25
Male	299	1	1	-	-	-	-	1	4	4	14	19
Female	251	1	-	-	2	1	-	-	2	2	6	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Colorado												
All races, both sexes	1,262	5	2	2	1	2	3	4	17	24	50	112
Male	671	4	2	-	1	2	1	3	11	12	37	78
Female	591	1	-	2	-	-	2	1	6	12	13	34
White, both sexes	1,202	4	1	2	1	2	3	4	16	22	49	105
Male	642	4	1	-	1	2	1	3	10	12	36	74
Female	560	-	-	2	-	-	2	1	6	10	13	31
Black, both sexes	36	1	1	-	-	-	-	-	-	1	1	5
Male	17	-	1	-	-	-	-	-	-	-	1	3
Female	19	1	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	1	1	-	2
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,150	9	2	1	1	1	3	2	3	13	34	44
Male	530	7	2	-	1	1	1	2	2	8	23	34
Female	620	2	-	1	-	-	2	-	1	5	11	10
White, both sexes	1,072	4	1	1	-	-	3	2	2	10	31	42
Male	492	3	1	-	-	-	1	2	2	7	21	33
Female	580	1	-	1	-	-	2	-	-	3	10	9
Black, both sexes	69	4	1	-	1	1	-	-	1	2	3	2
Male	34	3	1	-	1	1	-	-	-	1	2	1
Female	35	1	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Delaware												
All races, both sexes	227	-	-	-	1	-	-	1	-	2	8	14
Male	119	-	-	-	1	-	-	1	-	2	4	10
Female	108	-	-	-	-	-	-	-	-	-	4	4
White, both sexes	202	-	-	-	-	-	-	1	-	1	8	13
Male	108	-	-	-	-	-	-	1	-	1	4	9
Female	94	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	24	-	-	-	1	-	-	-	-	1	-	1
Male	10	-	-	-	1	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	198	2	2	-	-	-	-	1	1	1	9	14
Male	120	1	1	-	-	-	-	-	1	1	6	11
Female	78	1	1	-	-	-	-	1	-	-	3	3
White, both sexes	38	1	-	-	-	-	-	1	1	-	1	1
Male	22	-	-	-	-	-	-	-	1	-	1	1
Female	16	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	153	1	1	-	-	-	-	-	-	1	8	13
Male	92	1	-	-	-	-	-	-	-	1	5	10
Female	61	-	1	-	-	-	-	-	-	-	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Florida												
All races, both sexes	5,664	45	5	2	1	4	7	10	25	70	185	383
Male	2,981	29	1	1	1	3	3	4	12	39	115	251
Female	2,683	16	4	1	-	1	4	6	13	31	70	132
White, both sexes	5,129	20	4	1	1	4	2	7	19	59	162	329
Male	2,699	12	-	1	1	3	-	3	9	33	99	218
Female	2,430	8	4	-	-	1	2	4	10	26	63	111
Black, both sexes	496	24	-	1	-	-	5	3	6	10	21	54
Male	258	16	-	-	-	-	3	1	3	5	14	33
Female	238	8	-	1	-	-	2	2	3	5	7	21
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	1	-	-	-	-	-	-	1	2	-
Male	19	1	1	-	-	-	-	-	-	1	2	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,407	36	4	2	5	1	7	10	19	46	102	165
Male	1,236	25	4	-	2	-	3	5	12	28	62	111
Female	1,171	11	-	2	3	1	4	5	7	18	40	54
White, both sexes	1,782	12	2	2	2	-	6	8	9	28	74	107
Male	922	10	2	-	-	-	3	4	6	17	46	75
Female	860	2	-	2	2	-	3	4	3	11	28	32
Black, both sexes	602	22	1	-	3	1	1	2	10	16	26	55
Male	301	14	1	-	2	-	-	1	6	10	14	35
Female	301	8	-	-	1	1	1	1	4	6	12	20
American Indian 1/, both sexes	6	2	1	-	-	-	-	-	-	1	1	1
Male	3	1	1	-	-	-	-	-	-	-	1	-
Female	3	1	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Hawaii												
All races, both sexes	325	4	-	1	-	1	2	2	2	2	11	17
Male	177	2	-	1	-	-	-	2	1	2	6	12
Female	148	2	-	-	-	1	2	-	1	-	5	5
White, both sexes	115	1	-	-	-	-	2	1	1	-	2	7
Male	63	1	-	-	-	-	-	1	-	-	-	5
Female	52	-	-	-	-	-	2	-	1	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	206	3	-	1	-	1	-	1	1	2	9	10
Male	113	1	-	1	-	-	-	1	1	2	6	7
Female	93	2	-	-	-	1	-	-	-	-	3	3
Idaho												
All races, both sexes	406	5	-	-	-	1	-	2	2	6	12	22
Male	206	3	-	-	-	1	-	2	2	2	6	13
Female	200	2	-	-	-	-	-	-	-	4	6	9
White, both sexes	395	5	-	-	-	1	-	2	2	4	10	21
Male	202	3	-	-	-	1	-	2	2	2	6	13
Female	193	2	-	-	-	-	-	-	-	2	4	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	2	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	2	2	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Illinois												
All races, both sexes	3,574	16	2	4	2	2	9	21	28	47	145	213
Male	1,728	15	-	4	2	1	5	12	18	31	101	146
Female	1,846	1	2	-	-	1	4	9	10	16	44	67
White, both sexes	3,079	11	1	1	1	2	4	17	26	40	115	156
Male	1,451	10	-	1	1	1	4	11	17	26	82	104
Female	1,628	1	1	-	-	1	-	6	9	14	33	52
Black, both sexes	448	5	1	2	1	-	5	3	1	6	27	54
Male	252	5	-	2	1	-	1	1	1	4	17	40
Female	196	-	1	-	-	-	4	2	-	2	10	14
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	37	-	-	1	-	-	-	1	1	-	2	1
Male	20	-	-	1	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	1,832	22	5	2	2	-	5	4	14	25	54	108
Male	881	12	2	1	2	-	2	3	8	17	36	65
Female	951	10	3	1	-	-	3	1	6	8	18	43
White, both sexes	1,694	16	5	2	1	-	4	4	13	25	47	102
Male	809	8	2	1	1	-	1	3	7	17	31	62
Female	885	8	3	1	-	-	3	1	6	8	16	40
Black, both sexes	130	6	-	-	1	-	1	-	1	-	6	4
Male	69	4	-	-	1	-	1	-	1	-	4	3
Female	61	2	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Iowa												
All races, both sexes	876	2	1	-	1	1	1	2	6	8	16	45
Male	397	2	1	-	-	-	-	2	3	5	12	31
Female	479	-	-	-	1	1	1	-	3	3	4	14
White, both sexes	849	2	1	-	1	1	1	2	4	8	16	41
Male	385	2	1	-	-	-	-	2	2	5	12	29
Female	464	-	-	-	1	1	1	-	2	3	4	12
Black, both sexes	16	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	857	6	2	1	-	2	-	2	3	7	27	57
Male	419	4	1	1	-	1	-	1	2	4	22	39
Female	438	2	1	-	-	1	-	1	1	3	5	18
White, both sexes	806	4	1	1	-	2	-	2	3	7	24	50
Male	390	2	-	1	-	1	-	1	2	4	19	35
Female	416	2	1	-	-	1	-	1	1	3	5	15
Black, both sexes	31	2	1	-	-	-	-	-	-	-	1	5
Male	17	2	1	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	2	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Kentucky												
All races, both sexes	1,300	6	3	-	-	-	1	5	10	19	46	81
Male	668	5	1	-	-	-	1	2	4	16	33	58
Female	632	1	2	-	-	-	-	3	6	3	13	23
White, both sexes	1,222	4	3	-	-	-	1	4	9	19	41	80
Male	623	3	1	-	-	-	1	2	4	16	30	58
Female	599	1	2	-	-	-	-	2	5	3	11	22
Black, both sexes	72	2	-	-	-	-	-	1	1	-	5	1
Male	40	2	-	-	-	-	-	-	-	-	3	-
Female	32	-	-	-	-	-	-	1	1	-	2	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,389	8	2	1	2	3	4	7	17	20	59	98
Male	701	3	1	-	2	1	3	4	10	9	38	67
Female	688	5	1	1	-	2	1	3	7	11	21	31
White, both sexes	978	2	1	-	1	-	-	4	7	7	35	62
Male	481	1	-	-	1	-	-	3	5	2	22	41
Female	497	1	1	-	-	-	-	1	2	5	13	21
Black, both sexes	400	6	1	1	1	3	4	3	10	13	24	35
Male	215	2	1	-	1	1	3	1	5	7	16	25
Female	185	4	-	1	-	2	1	2	5	6	8	10
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Maine												
All races, both sexes	481	2	-	-	-	-	-	2	1	3	7	19
Male	248	1	-	-	-	-	-	1	1	1	5	11
Female	233	1	-	-	-	-	-	1	-	2	2	8
White, both sexes	478	2	-	-	-	-	-	2	1	3	7	19
Male	246	1	-	-	-	-	-	1	1	1	5	11
Female	232	1	-	-	-	-	-	1	-	2	2	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,508	10	-	2	-	1	3	7	5	25	49	88
Male	761	6	-	2	-	1	2	3	1	13	26	55
Female	747	4	-	-	-	-	1	4	4	12	23	33
White, both sexes	1,114	6	-	1	-	1	2	3	3	13	34	54
Male	546	4	-	1	-	1	2	-	1	7	20	35
Female	568	2	-	-	-	-	-	3	2	6	14	19
Black, both sexes	374	4	-	1	-	-	1	4	1	11	14	34
Male	208	2	-	1	-	-	-	3	-	6	5	20
Female	166	2	-	-	-	-	1	1	1	5	9	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Massachusetts												
All races, both sexes	2,118	8	1	1	-	1	1	3	11	24	72	109
Male	1,038	6	1	1	-	1	-	3	8	13	42	74
Female	1,080	2	-	-	-	-	1	-	3	11	30	35
White, both sexes	2,013	7	1	1	-	-	1	2	10	24	69	98
Male	981	6	1	1	-	-	-	2	7	13	40	67
Female	1,032	1	-	-	-	-	1	-	3	11	29	31
Black, both sexes	81	1	-	-	-	-	-	-	1	-	2	9
Male	43	-	-	-	-	-	-	-	1	-	2	5
Female	38	1	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	1	-	1	-	-	1	2
Male	14	-	-	-	-	1	-	1	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	3,078	20	7	-	2	3	8	6	28	53	118	189
Male	1,553	13	5	-	2	1	3	4	14	30	76	125
Female	1,525	7	2	-	-	2	5	2	14	23	42	64
White, both sexes	2,594	11	5	-	-	2	6	3	19	45	99	151
Male	1,274	7	4	-	-	-	3	2	9	26	62	102
Female	1,320	4	1	-	-	2	3	1	10	19	37	49
Black, both sexes	441	9	2	-	1	1	2	3	6	7	16	37
Male	253	6	1	-	1	1	-	2	3	3	12	22
Female	188	3	1	-	-	-	2	1	3	4	4	15
American Indian 1/, both sexes	24	-	-	-	1	-	-	-	2	1	3	1
Male	18	-	-	-	1	-	-	-	1	1	2	1
Female	6	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Minnesota												
All races, both sexes	1,291	8	-	-	1	2	3	1	11	14	42	67
Male	641	4	-	-	-	1	2	1	6	7	25	44
Female	650	4	-	-	1	1	1	-	5	7	17	23
White, both sexes	1,222	6	-	-	1	2	1	1	7	13	33	59
Male	598	4	-	-	-	1	1	1	5	6	19	38
Female	624	2	-	-	1	1	-	-	2	7	14	21
Black, both sexes	26	-	-	-	-	-	1	-	2	1	4	3
Male	16	-	-	-	-	-	1	-	-	1	3	2
Female	10	-	-	-	-	-	-	-	2	-	1	1
American Indian 1/, both sexes	29	1	-	-	-	-	-	-	2	-	5	4
Male	18	-	-	-	-	-	-	-	1	-	3	3
Female	11	1	-	-	-	-	-	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	1	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	951	10	5	1	-	-	8	3	17	21	29	76
Male	501	7	5	1	-	-	6	1	5	12	15	51
Female	450	3	-	-	-	-	2	2	12	9	14	25
White, both sexes	659	4	1	-	-	-	5	1	6	11	21	46
Male	347	2	1	-	-	-	4	-	4	7	11	33
Female	312	2	-	-	-	-	1	1	2	4	10	13
Black, both sexes	282	6	4	1	-	-	3	2	9	10	7	28
Male	152	5	4	1	-	-	2	1	1	5	4	17
Female	130	1	-	-	-	-	1	1	8	5	3	11
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	2	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	2	-	1	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Missouri												
All races, both sexes	1,792	10	4	1	-	3	2	3	9	16	51	104
Male	859	7	3	-	-	2	-	2	5	9	30	71
Female	933	3	1	1	-	1	2	1	4	7	21	33
White, both sexes	1,630	5	4	1	-	3	1	3	7	15	46	91
Male	778	5	3	-	-	2	-	2	4	8	28	61
Female	852	-	1	1	-	1	1	1	3	7	18	30
Black, both sexes	150	5	-	-	-	-	1	-	2	1	4	10
Male	75	2	-	-	-	-	-	-	1	1	1	8
Female	75	3	-	-	-	-	1	-	1	-	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	343	1	-	-	-	1	1	2	3	7	17	19
Male	175	-	-	-	-	-	1	-	1	3	13	12
Female	168	1	-	-	-	1	-	2	2	4	4	7
White, both sexes	298	-	-	-	-	1	1	-	-	3	10	15
Male	157	-	-	-	-	-	1	-	-	1	8	9
Female	141	-	-	-	-	1	-	-	-	2	2	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	44	1	-	-	-	-	-	2	3	4	7	3
Male	18	-	-	-	-	-	-	-	1	2	5	3
Female	26	1	-	-	-	-	-	2	2	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Nebraska												
All races, both sexes	577	3	2	1	2	-	1	1	4	3	31	28
Male	280	2	2	1	2	-	-	-	2	1	17	19
Female	297	1	-	-	-	-	1	1	2	2	14	9
White, both sexes	540	3	1	1	1	-	-	-	3	3	29	24
Male	259	2	1	1	1	-	-	-	2	1	16	17
Female	281	1	-	-	-	-	-	-	1	2	13	7
Black, both sexes	23	-	1	-	1	-	1	1	-	-	1	2
Male	14	-	1	-	1	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	1	1	-	-	-	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	768	1	1	-	-	2	3	3	6	18	48	64
Male	454	1	1	-	-	-	3	2	1	12	29	45
Female	314	-	-	-	-	2	-	1	5	6	19	19
White, both sexes	702	-	1	-	-	2	2	2	3	17	45	56
Male	417	-	1	-	-	-	2	2	1	11	26	38
Female	285	-	-	-	-	2	-	-	2	6	19	18
Black, both sexes	40	1	-	-	-	-	1	1	2	1	2	5
Male	21	1	-	-	-	-	1	-	-	1	2	5
Female	19	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
New Hampshire												
All races, both sexes	403	1	-	-	-	-	-	-	2	3	14	15
Male	193	-	-	-	-	-	-	-	2	2	10	7
Female	210	1	-	-	-	-	-	-	-	1	4	8
White, both sexes	401	1	-	-	-	-	-	-	2	3	13	15
Male	192	-	-	-	-	-	-	-	2	2	9	7
Female	209	1	-	-	-	-	-	-	-	1	4	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,379	16	4	2	2	3	6	4	14	40	86	131
Male	1,214	8	1	1	-	2	3	1	13	28	56	89
Female	1,165	8	3	1	2	1	3	3	1	12	30	42
White, both sexes	2,039	9	3	2	1	2	3	2	9	33	68	107
Male	1,043	4	1	1	-	1	2	-	8	24	44	74
Female	996	5	2	1	1	1	1	2	1	9	24	33
Black, both sexes	290	7	1	-	-	1	3	2	5	6	17	21
Male	141	4	-	-	-	1	1	1	5	3	11	12
Female	149	3	1	-	-	-	2	1	-	3	6	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	1	-	-	-	-	1	1	3
Male	28	-	-	-	-	-	-	-	-	1	1	3
Female	19	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
New Mexico												
All races, both sexes	717	5	3	-	1	2	2	7	9	18	46	62
Male	415	2	3	-	-	1	2	6	6	9	30	49
Female	302	3	-	-	1	1	-	1	3	9	16	13
White, both sexes	585	3	2	-	-	2	2	5	4	12	30	46
Male	344	2	2	-	-	1	2	4	2	8	23	37
Female	241	1	-	-	-	1	-	1	2	4	7	9
Black, both sexes	9	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	120	2	1	-	-	-	-	2	4	6	15	16
Male	67	-	1	-	-	-	-	2	3	1	7	12
Female	53	2	-	-	-	-	-	-	1	5	8	4
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,599	26	9	1	4	6	7	10	16	52	119	215
Male	2,264	21	6	1	3	3	6	5	13	37	83	158
Female	2,335	5	3	-	1	3	1	5	3	15	36	57
White, both sexes	3,872	16	7	-	1	5	5	9	14	31	101	166
Male	1,869	13	4	-	-	3	4	4	11	21	70	127
Female	2,003	3	3	-	1	2	1	5	3	10	31	39
Black, both sexes	608	10	2	1	3	1	2	-	2	14	17	39
Male	323	8	2	1	3	-	2	-	2	10	13	25
Female	285	2	-	-	-	1	-	-	-	4	4	14
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	1	1	4
Male	8	-	-	-	-	-	-	-	-	1	-	2
Female	10	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	1	-	6	-	6
Male	64	-	-	-	-	-	-	1	-	5	-	4
Female	37	-	-	-	-	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
North Carolina												
All races, both sexes	2,706	34	5	-	2	4	6	9	19	34	84	149
Male	1,341	21	-	-	1	2	3	7	9	23	50	103
Female	1,365	13	5	-	1	2	3	2	10	11	34	46
White, both sexes	2,122	18	2	-	1	3	4	9	12	22	55	116
Male	1,023	9	-	-	-	1	1	7	6	15	32	78
Female	1,099	9	2	-	1	2	3	2	6	7	23	38
Black, both sexes	537	16	3	-	1	1	1	-	6	11	26	33
Male	294	12	-	-	1	1	1	-	3	7	16	25
Female	243	4	3	-	-	-	-	-	3	4	10	8
American Indian 1/, both sexes	35	-	-	-	-	-	1	-	1	1	3	-
Male	20	-	-	-	-	-	1	-	-	1	2	-
Female	15	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	236	1	2	-	-	1	1	1	1	4	10	12
Male	120	1	2	-	-	1	1	1	1	2	6	6
Female	116	-	-	-	-	-	-	-	-	2	4	6
White, both sexes	210	1	2	-	-	1	1	1	-	2	6	11
Male	105	1	2	-	-	1	1	1	-	1	4	5
Female	105	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	2	4	1
Male	15	-	-	-	-	-	-	-	1	1	2	1
Female	11	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Ohio												
All races, both sexes	3,816	18	3	2	4	3	5	13	16	57	123	217
Male	1,867	13	2	1	2	1	3	6	7	32	77	143
Female	1,949	5	1	1	2	2	2	7	9	25	46	74
White, both sexes	3,402	13	2	2	4	1	5	10	14	47	101	173
Male	1,641	9	1	1	2	-	3	5	7	26	62	111
Female	1,761	4	1	1	2	1	2	5	7	21	39	62
Black, both sexes	395	5	1	-	-	2	-	3	2	6	19	44
Male	217	4	1	-	-	1	-	1	-	4	12	32
Female	178	1	-	-	-	1	-	2	2	2	7	12
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	3	2	-
Male	5	-	-	-	-	-	-	-	-	2	2	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	1,350	7	3	1	1	3	4	7	4	17	71	81
Male	649	4	1	-	1	-	2	1	2	14	38	48
Female	701	3	2	1	-	3	2	6	2	3	33	33
White, both sexes	1,148	6	2	-	1	1	4	4	4	12	48	59
Male	547	3	-	-	1	-	2	-	2	11	29	38
Female	601	3	2	-	-	1	2	4	2	1	19	21
Black, both sexes	67	1	1	1	-	-	-	2	-	1	5	8
Male	33	1	1	-	-	-	-	1	-	1	2	4
Female	34	-	-	1	-	-	-	1	-	-	3	4
American Indian 1/, both sexes	130	-	-	-	-	2	-	1	-	4	18	14
Male	67	-	-	-	-	-	-	-	-	2	7	6
Female	63	-	-	-	-	2	-	1	-	2	11	8
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Oregon												
All races, both sexes	1,186	8	1	-	1	2	1	3	8	20	33	85
Male	591	4	1	-	1	-	1	2	4	12	20	49
Female	595	4	-	-	-	2	-	1	4	8	13	36
White, both sexes	1,140	7	1	-	1	2	1	2	8	20	30	78
Male	567	4	1	-	1	-	1	1	4	12	18	45
Female	573	3	-	-	-	2	-	1	4	8	12	33
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	-	-	1	6
Male	12	-	-	-	-	-	-	1	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	3
Asian or Pacific Islander 2/, both sexes	18	1	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	1	-	-	-	-	-	-	-	-	1	-
Pennsylvania												
All races, both sexes	4,439	14	4	1	1	5	7	8	18	48	123	235
Male	2,077	12	4	1	1	2	4	3	12	32	76	156
Female	2,362	2	-	-	-	3	3	5	6	16	47	79
White, both sexes	4,027	9	3	1	-	5	1	6	14	42	112	187
Male	1,882	8	3	1	-	2	1	2	10	29	73	127
Female	2,145	1	-	-	-	3	-	4	4	13	39	60
Black, both sexes	394	5	1	-	1	-	6	2	4	6	11	47
Male	188	4	1	-	1	-	3	1	2	3	3	29
Female	206	1	-	-	-	-	3	1	2	3	8	18
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Rhode Island												
All races, both sexes	377	2	1	1	-	-	-	2	-	5	9	13
Male	167	2	1	1	-	-	-	1	-	5	3	7
Female	210	-	-	-	-	-	-	1	-	-	6	6
White, both sexes	362	1	1	1	-	-	-	2	-	4	9	13
Male	158	1	1	1	-	-	-	1	-	4	3	7
Female	204	-	-	-	-	-	-	1	-	-	6	6
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,495	13	3	3	2	-	2	4	8	20	58	102
Male	760	8	1	1	-	-	1	3	5	8	33	65
Female	735	5	2	2	2	-	1	1	3	12	25	37
White, both sexes	1,107	5	-	2	1	-	-	3	4	12	38	72
Male	565	3	-	-	-	-	-	2	3	5	25	46
Female	542	2	-	2	1	-	-	1	1	7	13	26
Black, both sexes	379	8	3	1	1	-	2	1	4	8	20	29
Male	189	5	1	1	-	-	1	1	2	3	8	18
Female	190	3	2	-	1	-	1	-	2	5	12	11
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
South Dakota												
All races, both sexes	279	2	-	-	-	-	1	1	3	9	10	20
Male	142	2	-	-	-	-	-	1	3	4	7	15
Female	137	-	-	-	-	-	1	-	-	5	3	5
White, both sexes	228	1	-	-	-	-	1	-	1	3	3	11
Male	112	1	-	-	-	-	-	-	1	1	2	8
Female	116	-	-	-	-	-	1	-	-	2	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	50	1	-	-	-	-	-	1	2	6	7	8
Male	29	1	-	-	-	-	-	1	2	3	5	6
Female	21	-	-	-	-	-	-	-	-	3	2	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,939	11	2	-	1	-	1	7	12	24	74	112
Male	982	5	1	-	-	-	-	6	8	15	50	73
Female	957	6	1	-	1	-	1	1	4	9	24	39
White, both sexes	1,699	3	2	-	1	-	1	6	9	20	63	92
Male	852	2	1	-	-	-	-	5	5	13	44	59
Female	847	1	1	-	1	-	1	1	4	7	19	33
Black, both sexes	231	6	-	-	-	-	-	1	2	4	11	20
Male	125	3	-	-	-	-	-	1	2	2	6	14
Female	106	3	-	-	-	-	-	-	-	2	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	2	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	3	2	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Texas												
All races, both sexes	6,187	69	8	8	4	7	14	28	41	106	285	477
Male	3,315	31	3	4	2	4	9	15	30	79	193	338
Female	2,872	38	5	4	2	3	5	13	11	27	92	139
White, both sexes	5,541	56	7	6	2	4	10	26	32	92	260	418
Male	2,966	24	3	2	1	2	6	15	22	71	178	302
Female	2,575	32	4	4	1	2	4	11	10	21	82	116
Black, both sexes	575	12	1	2	2	3	4	2	7	12	19	54
Male	308	6	-	2	1	2	3	-	6	6	11	32
Female	267	6	1	-	1	1	1	2	1	6	8	22
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	3	1
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	58	1	-	-	-	-	-	-	2	2	3	4
Male	35	1	-	-	-	-	-	-	2	2	2	4
Female	23	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	473	2	-	-	-	-	-	3	2	9	7	29
Male	244	1	-	-	-	-	-	1	2	9	5	19
Female	229	1	-	-	-	-	-	2	-	-	2	10
White, both sexes	457	2	-	-	-	-	-	3	2	7	7	26
Male	232	1	-	-	-	-	-	1	2	7	5	16
Female	225	1	-	-	-	-	-	2	-	-	2	10
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	3
Male	5	-	-	-	-	-	-	-	-	1	-	3
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Vermont												
All races, both sexes	193	-	-	-	-	1	-	1	-	2	5	8
Male	84	-	-	-	-	1	-	-	-	1	4	5
Female	109	-	-	-	-	-	-	1	-	1	1	3
White, both sexes	191	-	-	-	-	1	-	1	-	2	5	8
Male	83	-	-	-	-	1	-	-	-	1	4	5
Female	108	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,003	12	2	-	-	2	-	6	13	34	69	119
Male	1,019	6	1	-	-	1	-	3	4	19	45	85
Female	984	6	1	-	-	1	-	3	9	15	24	34
White, both sexes	1,588	6	1	-	-	2	-	3	11	25	55	90
Male	808	2	-	-	-	1	-	2	4	13	39	66
Female	780	4	1	-	-	1	-	1	7	12	16	24
Black, both sexes	395	6	1	-	-	-	-	3	1	8	14	29
Male	206	4	1	-	-	-	-	1	-	6	6	19
Female	189	2	-	-	-	-	-	2	1	2	8	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Washington												
All races, both sexes	1,660	16	3	1	-	1	2	5	8	22	54	98
Male	811	11	2	1	-	-	1	3	4	15	31	66
Female	849	5	1	-	-	1	1	2	4	7	23	32
White, both sexes	1,525	10	3	1	-	1	2	5	6	16	48	83
Male	733	7	2	1	-	-	1	3	2	9	28	55
Female	792	3	1	-	-	1	1	2	4	7	20	28
Black, both sexes	44	3	-	-	-	-	-	-	-	1	2	7
Male	27	2	-	-	-	-	-	-	-	1	1	6
Female	17	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	50	2	-	-	-	-	-	-	2	3	4	7
Male	28	1	-	-	-	-	-	-	2	3	2	4
Female	22	1	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, both sexes	41	1	-	-	-	-	-	-	-	2	-	1
Male	23	1	-	-	-	-	-	-	-	2	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	706	4	1	1	-	-	1	3	-	7	26	36
Male	343	2	-	1	-	-	1	2	-	5	17	25
Female	363	2	1	-	-	-	-	1	-	2	9	11
White, both sexes	682	3	1	1	-	-	1	3	-	7	22	34
Male	334	1	-	1	-	-	1	2	-	5	16	25
Female	348	2	1	-	-	-	-	1	-	2	6	9
Black, both sexes	24	1	-	-	-	-	-	-	-	-	4	2
Male	9	1	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
Wisconsin												
All races, both sexes	1,700	9	2	-	-	-	4	2	5	21	57	84
Male	797	4	2	-	-	-	2	-	3	12	40	59
Female	903	5	-	-	-	-	2	2	2	9	17	25
White, both sexes	1,584	7	-	-	-	-	3	2	3	19	49	67
Male	743	4	-	-	-	-	1	-	2	11	34	47
Female	841	3	-	-	-	-	2	2	1	8	15	20
Black, both sexes	73	2	1	-	-	-	-	-	-	-	4	14
Male	31	-	1	-	-	-	-	-	-	-	3	11
Female	42	2	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	33	-	1	-	-	-	1	-	1	1	4	3
Male	17	-	1	-	-	-	1	-	-	-	3	1
Female	16	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	186	1	-	-	-	-	-	1	2	2	6	17
Male	91	1	-	-	-	-	-	1	2	1	2	5
Female	95	-	-	-	-	-	-	-	-	1	4	12
White, both sexes	174	1	-	-	-	-	-	1	2	-	5	15
Male	85	1	-	-	-	-	-	1	2	-	2	4
Female	89	-	-	-	-	-	-	-	-	-	3	11
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	2	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
United States												
All races, both sexes	6,995	7,228	6,592	6,641	7,838	9,658	11,226	9,844	6,517	2,351	407	8
Male	4,897	4,862	4,112	3,701	4,045	4,547	4,618	3,382	1,853	512	78	8
Female	2,098	2,366	2,480	2,940	3,793	5,111	6,608	6,462	4,664	1,839	329	-
White, both sexes	5,709	6,077	5,562	5,671	6,872	8,590	10,176	9,079	6,022	2,122	357	7
Male	4,055	4,121	3,487	3,157	3,563	4,049	4,202	3,121	1,721	454	61	7
Female	1,654	1,956	2,075	2,514	3,309	4,541	5,974	5,958	4,301	1,668	296	-
Black, both sexes	1,036	961	818	765	764	819	816	585	390	182	42	1
Male	676	613	495	441	391	382	307	183	93	39	12	1
Female	360	348	323	324	373	437	509	402	297	143	30	-
American Indian 1/, both sexes	148	112	106	80	65	57	56	32	18	9	2	-
Male	91	74	70	35	25	24	21	11	6	3	1	-
Female	57	38	36	45	40	33	35	21	12	6	1	-
Asian or Pacific Islander 2/, both sexes	102	78	106	125	137	192	178	148	87	38	6	-
Male	75	54	60	68	66	92	88	67	33	16	4	-
Female	27	24	46	57	71	100	90	81	54	22	2	-
Alabama												
All races, both sexes	132	140	128	147	147	181	206	162	92	33	7	-
Male	89	95	80	79	69	82	86	45	17	9	3	-
Female	43	45	48	68	78	99	120	117	75	24	4	-
White, both sexes	93	106	103	116	120	153	173	137	77	24	3	-
Male	61	72	66	62	57	69	74	42	14	7	-	-
Female	32	34	37	54	63	84	99	95	63	17	3	-
Black, both sexes	39	33	25	30	27	28	33	25	15	9	4	-
Male	28	22	14	17	12	13	12	3	3	2	3	-
Female	11	11	11	13	15	15	21	22	12	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Alaska												
All races, both sexes	16	11	14	9	14	11	16	10	5	1	1	-
Male	10	8	11	3	9	5	4	3	-	1	-	-
Female	6	3	3	6	5	6	12	7	5	-	1	-
White, both sexes	12	8	10	4	12	7	12	6	4	1	1	-
Male	7	6	8	2	8	4	4	1	-	1	-	-
Female	5	2	2	2	4	3	8	5	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	4	5	1	4	3	4	1	-	-	-
Male	2	2	3	1	-	1	-	2	-	-	-	-
Female	1	1	1	4	1	3	3	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	182	159	164	147	175	164	196	156	118	33	8	1
Male	137	105	94	93	100	77	96	52	40	9	1	1
Female	45	54	70	54	75	87	100	104	78	24	7	-
White, both sexes	150	142	143	134	159	150	181	154	114	31	8	1
Male	115	95	84	87	94	72	87	51	39	7	1	1
Female	35	47	59	47	65	78	94	103	75	24	7	-
Black, both sexes	3	4	7	4	4	5	4	-	2	-	-	-
Male	1	2	4	1	2	2	4	-	1	-	-	-
Female	2	2	3	3	2	3	-	-	1	-	-	-
American Indian 1/, both sexes	29	13	13	7	11	7	11	2	2	2	-	-
Male	21	8	6	4	3	2	5	1	-	2	-	-
Female	8	5	7	3	8	5	6	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	2	-	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Arkansas												
All races, both sexes	77	84	70	74	104	99	127	111	73	23	6	-
Male	53	52	37	45	54	41	49	38	19	3	1	-
Female	24	32	33	29	50	58	78	73	54	20	5	-
White, both sexes	66	69	61	63	92	88	117	104	66	18	3	-
Male	45	47	34	38	51	32	45	37	17	2	-	-
Female	21	22	27	25	41	56	72	67	49	16	3	-
Black, both sexes	10	14	9	9	12	11	10	7	7	5	3	-
Male	8	5	3	5	3	9	4	1	2	1	1	-
Female	2	9	6	4	9	2	6	6	5	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	908	902	761	743	754	892	1,074	898	548	206	33	3
Male	635	598	514	418	416	456	481	343	187	50	9	3
Female	273	304	247	325	338	436	593	555	361	156	24	-
White, both sexes	775	794	654	630	645	753	934	803	487	179	30	3
Male	550	537	449	351	355	384	414	310	161	44	6	3
Female	225	257	205	279	290	369	520	493	326	135	24	-
Black, both sexes	80	70	58	57	50	49	54	28	19	8	1	-
Male	49	37	34	36	31	25	22	8	7	1	1	-
Female	31	33	24	21	19	24	32	20	12	7	-	-
American Indian 1/, both sexes	13	15	12	10	9	5	5	3	2	1	-	-
Male	8	9	10	3	5	3	2	-	-	-	-	-
Female	5	6	2	7	4	2	3	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	40	23	37	46	50	85	81	64	40	18	2	-
Male	28	15	21	28	25	44	43	25	19	5	2	-
Female	12	8	16	18	25	41	38	39	21	13	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Colorado												
All races, both sexes	120	127	90	92	95	150	118	132	74	38	4	-
Male	80	83	55	53	47	71	52	46	22	11	-	-
Female	40	44	35	39	48	79	66	86	52	27	4	-
White, both sexes	111	124	79	81	94	144	115	132	72	37	4	-
Male	72	81	50	49	46	70	52	46	21	11	-	-
Female	39	43	29	32	48	74	63	86	51	26	4	-
Black, both sexes	8	-	7	4	-	3	3	-	1	1	-	-
Male	7	-	3	2	-	-	-	-	-	-	-	-
Female	1	-	4	2	-	3	3	-	1	1	-	-
American Indian 1/, both sexes	1	3	3	3	1	2	-	-	1	-	-	-
Male	1	2	2	-	1	1	-	-	1	-	-	-
Female	-	1	1	3	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	4	-	1	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	68	80	79	66	78	145	155	173	123	63	7	-
Male	36	57	46	38	37	68	59	68	30	8	2	-
Female	32	23	33	28	41	77	96	105	93	55	5	-
White, both sexes	57	73	67	62	75	136	147	170	120	63	6	-
Male	32	53	38	36	35	64	56	68	30	8	2	-
Female	25	20	29	26	40	72	91	102	90	55	4	-
Black, both sexes	11	6	10	4	3	8	7	3	1	-	1	-
Male	4	3	6	2	2	4	3	-	-	-	-	-
Female	7	3	4	2	1	4	4	3	1	-	1	-
American Indian 1/, both sexes	-	1	1	-	-	1	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Delaware												
All races, both sexes	16	28	16	14	21	27	31	30	14	3	1	-
Male	10	20	12	7	8	13	16	11	4	-	-	-
Female	6	8	4	7	13	14	15	19	10	3	1	-
White, both sexes	13	25	15	11	19	25	28	26	13	3	1	-
Male	8	18	12	6	8	13	16	9	3	-	-	-
Female	5	7	3	5	11	12	12	17	10	3	1	-
Black, both sexes	3	3	1	3	2	2	3	4	-	-	-	-
Male	2	2	-	1	-	-	-	2	-	-	-	-
Female	1	1	1	2	2	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	31	15	12	16	13	24	19	10	7	1	-
Male	16	20	10	9	12	8	9	11	2	2	-	-
Female	4	11	5	3	4	5	15	8	8	5	1	-
White, both sexes	3	3	1	3	2	3	5	4	5	3	1	-
Male	2	2	1	3	2	2	3	2	2	-	-	-
Female	1	1	-	-	-	1	2	2	3	3	1	-
Black, both sexes	16	28	13	9	14	10	17	13	5	4	-	-
Male	13	18	8	6	10	6	4	8	-	2	-	-
Female	3	10	5	3	4	4	13	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	2	1	-	-	-	-
Male	1	-	1	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Florida												
All races, both sexes	450	553	459	431	494	672	699	612	394	138	25	-
Male	322	384	294	250	261	323	310	224	109	37	8	-
Female	128	169	165	181	233	349	389	388	285	101	17	-
White, both sexes	394	494	409	389	453	627	656	575	375	127	22	-
Male	284	344	269	225	244	302	294	210	104	37	7	-
Female	110	150	140	164	209	325	362	365	271	90	15	-
Black, both sexes	49	55	48	39	38	42	38	34	16	10	3	-
Male	32	36	24	23	16	21	15	12	3	-	1	-
Female	17	19	24	16	22	21	23	22	13	10	2	-
American Indian 1/, both sexes	3	1	-	-	1	-	1	-	-	-	-	-
Male	3	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	2	3	2	3	4	3	3	1	-	-
Male	3	3	1	2	1	-	-	2	2	-	-	-
Female	1	-	1	1	1	3	4	1	1	1	-	-
Georgia												
All races, both sexes	236	246	198	201	223	249	267	219	122	41	8	-
Male	152	164	116	119	120	111	101	64	29	7	1	-
Female	84	82	82	82	103	138	166	155	93	34	7	-
White, both sexes	160	175	148	149	174	197	209	181	101	32	6	-
Male	109	120	85	91	93	88	85	57	26	4	1	-
Female	51	55	63	58	81	109	124	124	75	28	5	-
Black, both sexes	74	71	48	51	46	49	57	38	20	9	2	-
Male	42	44	30	28	25	21	15	7	3	3	-	-
Female	32	27	18	23	21	28	42	31	17	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	1	3	3	1	-	1	-	-	-
Male	1	-	1	-	2	2	1	-	-	-	-	-
Female	1	-	1	1	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Hawaii												
All races, both sexes	26	23	29	20	31	33	43	40	24	13	1	-
Male	18	15	19	15	16	17	20	14	9	7	1	-
Female	8	8	10	5	15	16	23	26	15	6	-	-
White, both sexes	11	13	16	10	8	11	16	6	4	6	-	-
Male	7	9	10	9	3	7	6	2	2	1	-	-
Female	4	4	6	1	5	4	10	4	2	5	-	-
Black, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	9	13	9	22	22	27	34	19	7	1	-
Male	11	6	9	5	13	10	14	12	7	6	1	-
Female	4	3	4	4	9	12	13	22	12	1	-	-
Idaho												
All races, both sexes	31	31	31	28	34	38	63	60	29	11	-	-
Male	20	23	18	18	17	17	25	26	10	3	-	-
Female	11	8	13	10	17	21	38	34	19	8	-	-
White, both sexes	30	30	30	28	32	37	63	60	29	11	-	-
Male	19	23	18	18	15	16	25	26	10	3	-	-
Female	11	7	12	10	17	21	38	34	19	8	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Illinois												
All races, both sexes	236	273	250	240	304	404	496	461	299	101	21	-
Male	156	181	156	143	150	170	188	146	84	13	6	-
Female	80	92	94	97	154	234	308	315	215	88	15	-
White, both sexes	185	211	203	203	273	359	457	432	271	94	17	-
Male	122	135	127	117	138	153	175	135	77	12	3	-
Female	63	76	76	86	135	206	282	297	194	82	14	-
Black, both sexes	47	58	46	33	25	38	33	26	27	7	3	-
Male	33	42	28	23	10	13	12	8	7	1	3	-
Female	14	16	18	10	15	25	21	18	20	6	-	-
American Indian 1/, both sexes	4	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	2	6	7	6	3	1	-	1	-
Male	-	4	1	1	2	4	1	3	-	-	-	-
Female	-	-	-	1	4	3	5	-	1	-	1	-
Indiana												
All races, both sexes	149	133	142	148	163	213	237	218	140	41	7	-
Male	101	92	78	72	77	95	98	70	35	13	2	-
Female	48	41	64	76	86	118	139	148	105	28	5	-
White, both sexes	120	122	129	139	154	196	225	210	133	41	6	-
Male	79	86	71	68	74	89	93	67	34	13	2	-
Female	41	36	58	71	80	107	132	143	99	28	4	-
Black, both sexes	29	11	12	8	9	14	12	8	7	-	1	-
Male	22	6	6	4	3	5	5	3	1	-	-	-
Female	7	5	6	4	6	9	7	5	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Iowa												
All races, both sexes	56	36	52	60	68	113	134	121	113	33	7	-
Male	40	24	30	31	31	57	52	36	34	6	-	-
Female	16	12	22	29	37	56	82	85	79	27	7	-
White, both sexes	50	33	50	59	66	109	133	120	113	32	7	-
Male	35	23	30	30	31	56	52	35	34	6	-	-
Female	15	10	20	29	35	53	81	85	79	26	7	-
Black, both sexes	3	1	2	-	1	3	1	1	-	1	-	-
Male	3	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	2	-	1	2	1	-	-	1	-	-
American Indian 1/, both sexes	3	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	56	63	45	61	54	98	119	118	99	30	7	-
Male	39	36	31	40	24	49	46	46	23	7	3	-
Female	17	27	14	21	30	49	73	72	76	23	4	-
White, both sexes	55	59	40	52	50	93	114	115	98	29	7	-
Male	38	35	28	34	22	47	44	45	23	6	3	-
Female	17	24	12	18	28	46	70	70	75	23	4	-
Black, both sexes	-	4	3	4	3	2	3	1	1	1	-	-
Male	-	1	2	3	1	2	1	-	-	1	-	-
Female	-	3	1	1	2	-	2	1	1	-	-	-
American Indian 1/, both sexes	1	-	1	3	1	3	-	1	-	-	-	-
Male	1	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	3	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	2	1	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Kentucky												
All races, both sexes	104	101	114	115	115	143	175	142	93	23	4	-
Male	78	67	61	62	60	70	62	52	29	5	2	-
Female	26	34	53	53	55	73	113	90	64	18	2	-
White, both sexes	93	95	108	108	107	134	169	133	87	23	4	-
Male	67	62	57	59	56	64	59	49	28	5	2	-
Female	26	33	51	49	51	70	110	84	59	18	2	-
Black, both sexes	10	4	5	7	8	9	5	8	6	-	-	-
Male	10	3	3	3	4	6	2	3	1	-	-	-
Female	-	1	2	4	4	3	3	5	5	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	121	113	123	124	134	158	155	135	72	30	3	-
Male	85	81	77	60	67	75	58	38	14	7	1	-
Female	36	32	46	64	67	83	97	97	58	23	2	-
White, both sexes	80	75	96	84	96	119	120	107	58	21	3	-
Male	60	54	59	41	47	57	45	28	11	3	1	-
Female	20	21	37	43	49	62	75	79	47	18	2	-
Black, both sexes	41	35	27	40	37	38	32	26	14	9	-	-
Male	25	24	18	19	20	17	13	10	3	4	-	-
Female	16	11	9	21	17	21	19	16	11	5	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	3	1	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Maine												
All races, both sexes	36	31	36	30	33	64	74	75	47	17	4	-
Male	32	22	22	14	15	37	28	37	12	6	3	-
Female	4	9	14	16	18	27	46	38	35	11	1	-
White, both sexes	35	31	35	30	33	64	74	74	47	17	4	-
Male	31	22	22	14	15	37	28	36	12	6	3	-
Female	4	9	13	16	18	27	46	38	35	11	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	112	131	128	119	125	170	203	181	104	38	6	1
Male	87	92	83	70	53	84	86	59	30	5	2	1
Female	25	39	45	49	72	86	117	122	74	33	4	-
White, both sexes	69	88	76	84	100	133	168	158	86	31	4	-
Male	52	61	46	47	43	68	75	53	25	4	1	-
Female	17	27	30	37	57	65	93	105	61	27	3	-
Black, both sexes	40	42	49	32	24	34	35	21	17	7	2	1
Male	33	31	35	22	10	16	11	5	5	1	1	1
Female	7	11	14	10	14	18	24	16	12	6	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	3	1	3	-	2	-	-	-	-
Male	2	-	2	1	-	-	-	1	-	-	-	-
Female	-	1	1	2	1	3	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Massachusetts												
All races, both sexes	156	146	125	127	189	254	309	273	204	89	15	-
Male	111	88	72	73	104	127	124	98	69	21	2	-
Female	45	58	53	54	85	127	185	175	135	68	13	-
White, both sexes	150	134	117	122	173	235	301	269	200	84	15	-
Male	106	81	68	70	99	115	120	97	68	18	2	-
Female	44	53	49	52	74	120	181	172	132	66	13	-
Black, both sexes	4	12	8	5	12	12	7	4	2	2	-	-
Male	4	7	4	3	5	7	3	1	-	1	-	-
Female	-	5	4	2	7	5	4	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	4	7	1	-	2	3	-	-
Male	1	-	-	-	-	5	1	-	1	2	-	-
Female	1	-	-	-	4	2	-	-	1	1	-	-
Michigan												
All races, both sexes	272	266	230	222	297	321	388	333	228	74	13	-
Male	187	190	142	122	153	146	145	121	60	12	2	-
Female	85	76	88	100	144	175	243	212	168	62	11	-
White, both sexes	210	203	193	179	254	281	348	301	203	68	13	-
Male	144	147	116	93	130	128	126	111	52	10	2	-
Female	66	56	77	86	124	153	222	190	151	58	11	-
Black, both sexes	56	58	31	39	40	40	34	28	25	6	-	-
Male	37	39	22	28	21	18	19	8	8	2	-	-
Female	19	19	9	11	19	22	15	20	17	4	-	-
American Indian 1/, both sexes	5	3	5	1	1	-	1	-	-	-	-	-
Male	5	2	4	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	3	2	-	5	4	-	-	-	-
Male	1	2	-	1	1	-	-	2	-	-	-	-
Female	-	-	1	2	1	-	5	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Minnesota												
All races, both sexes	83	82	83	103	101	110	181	175	163	51	10	-
Male	62	55	53	49	57	56	91	62	51	13	2	-
Female	21	27	30	54	44	54	90	113	112	38	8	-
White, both sexes	75	75	75	100	97	103	178	173	162	51	10	-
Male	56	51	45	49	55	53	88	61	50	13	2	-
Female	19	24	30	51	42	50	90	112	112	38	8	-
Black, both sexes	4	4	3	1	-	1	2	-	-	-	-	-
Male	3	1	3	-	-	-	2	-	-	-	-	-
Female	1	3	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	5	1	2	4	-	1	-	-	-	-
Male	2	1	5	-	1	2	-	-	-	-	-	-
Female	1	-	-	1	1	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	1	2	2	1	1	1	-	-	-
Male	1	2	-	-	1	1	1	1	1	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	82	93	67	75	97	109	103	87	37	29	2	-
Male	64	68	39	41	52	50	38	25	13	8	-	-
Female	18	25	28	34	45	59	65	62	24	21	2	-
White, both sexes	49	69	51	48	76	80	83	63	29	16	-	-
Male	38	51	30	27	39	38	31	17	10	4	-	-
Female	11	18	21	21	37	42	52	46	19	12	-	-
Black, both sexes	31	24	16	26	20	28	20	24	8	13	2	-
Male	25	17	9	14	13	12	7	8	3	4	-	-
Female	6	7	7	12	7	16	13	16	5	9	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Missouri												
All races, both sexes	123	105	115	124	184	213	262	233	165	52	13	-
Male	90	67	59	59	103	110	113	74	40	13	2	-
Female	33	38	56	65	81	103	149	159	125	39	11	-
White, both sexes	104	88	95	111	169	197	248	227	152	51	12	-
Male	79	55	53	54	94	99	107	71	39	13	1	-
Female	25	33	42	57	75	98	141	156	113	38	11	-
Black, both sexes	18	17	18	12	14	14	13	6	13	1	1	-
Male	11	12	6	5	8	10	5	3	1	-	1	-
Female	7	5	12	7	6	4	8	3	12	1	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	37	25	15	28	25	35	40	42	36	9	-	-
Male	24	14	12	15	11	21	14	20	13	1	-	-
Female	13	11	3	13	14	14	26	22	23	8	-	-
White, both sexes	29	22	14	24	25	32	37	41	35	9	-	-
Male	22	13	11	13	11	20	14	20	13	1	-	-
Female	7	9	3	11	14	12	23	21	22	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	3	1	4	-	3	3	1	1	-	-	-
Male	2	1	1	2	-	1	-	-	-	-	-	-
Female	6	2	-	2	-	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Nebraska												
All races, both sexes	32	36	33	38	43	66	86	73	53	39	2	-
Male	26	25	19	22	24	34	40	23	12	9	-	-
Female	6	11	14	16	19	32	46	50	41	30	2	-
White, both sexes	28	29	31	35	40	63	84	73	52	38	2	-
Male	23	20	19	21	21	31	40	23	11	9	-	-
Female	5	9	12	14	19	32	44	50	41	29	2	-
Black, both sexes	1	3	2	2	3	2	1	-	1	1	-	-
Male	1	3	-	-	3	2	-	-	1	-	-	-
Female	-	-	2	2	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	3	4	-	1	-	1	-	-	-	-	-	-
Male	2	2	-	1	-	1	-	-	-	-	-	-
Female	1	2	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Nevada												
All races, both sexes	67	79	84	73	74	79	80	53	26	5	2	-
Male	42	53	59	46	40	44	42	25	5	2	2	-
Female	25	26	25	27	34	35	38	28	21	3	-	-
White, both sexes	62	72	77	65	69	72	74	51	25	5	2	-
Male	40	49	55	40	38	42	38	25	5	2	2	-
Female	22	23	22	25	31	30	36	26	20	3	-	-
Black, both sexes	4	5	4	3	3	4	3	1	-	-	-	-
Male	1	2	3	2	-	2	1	-	-	-	-	-
Female	3	3	1	1	3	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	1	2	3	1	1	-	-	-
Male	1	2	1	3	1	-	3	-	-	-	-	-
Female	-	-	1	1	-	2	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
New Hampshire												
All races, both sexes	27	24	25	25	34	58	69	57	31	16	2	-
Male	21	13	13	15	14	25	30	27	11	3	-	-
Female	6	11	12	10	20	33	39	30	20	13	2	-
White, both sexes	27	23	25	25	34	58	69	57	31	16	2	-
Male	21	13	13	15	14	25	30	27	11	3	-	-
Female	6	10	12	10	20	33	39	30	20	13	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	182	174	163	165	236	261	338	283	191	69	9	-
Male	127	115	104	94	127	122	153	92	60	16	2	-
Female	55	59	59	71	109	139	185	191	131	53	7	-
White, both sexes	142	149	134	136	202	231	302	260	175	63	6	-
Male	102	101	87	78	111	110	135	87	57	14	2	-
Female	40	48	47	58	91	121	167	173	118	49	4	-
Black, both sexes	36	24	25	20	27	24	31	20	13	4	3	-
Male	22	13	14	8	13	9	16	4	3	1	-	-
Female	14	11	11	12	14	15	15	16	10	3	3	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	4	9	6	5	4	3	3	2	-	-
Male	3	1	3	8	2	2	2	1	-	1	-	-
Female	1	-	1	1	4	3	2	2	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
New Mexico												
All races, both sexes	75	67	50	60	63	64	82	52	33	16	-	-
Male	56	40	36	37	36	32	40	18	7	5	-	-
Female	19	27	14	23	27	32	42	34	26	11	-	-
White, both sexes	55	56	40	51	51	58	76	48	31	13	-	-
Male	45	33	31	32	33	29	35	16	5	4	-	-
Female	10	23	9	19	18	29	41	32	26	9	-	-
Black, both sexes	1	-	1	1	-	-	2	2	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	1	-	1	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	19	11	9	8	11	6	4	2	2	2	-	-
Male	11	7	5	4	3	3	3	2	2	1	-	-
Female	8	4	4	4	8	3	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
New York												
All races, both sexes	315	323	348	365	455	577	593	597	390	146	25	-
Male	222	218	223	201	208	273	243	196	120	23	1	-
Female	93	105	125	164	247	304	350	401	270	123	24	-
White, both sexes	233	266	270	297	382	491	520	542	359	134	23	-
Male	168	180	168	162	170	238	213	175	115	22	1	-
Female	65	86	102	135	212	253	307	367	244	112	22	-
Black, both sexes	74	52	64	57	65	65	60	41	27	10	2	-
Male	47	35	45	32	36	22	22	13	4	1	-	-
Female	27	17	19	25	29	43	38	28	23	9	2	-
American Indian 1/, both sexes	1	2	2	2	1	-	3	1	-	-	-	-
Male	1	2	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	2	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	12	9	7	21	10	13	4	2	-	-
Male	6	1	9	7	2	13	7	8	1	-	-	-
Female	1	2	3	2	5	8	3	5	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
North Carolina												
All races, both sexes	218	225	226	235	278	311	335	278	189	58	7	-
Male	149	144	145	148	128	151	120	74	54	9	-	-
Female	69	81	81	87	150	160	215	204	135	49	7	-
White, both sexes	154	167	172	180	223	264	274	235	160	45	6	-
Male	115	109	106	113	101	122	97	61	45	5	-	-
Female	39	58	66	67	122	142	177	174	115	40	6	-
Black, both sexes	54	54	46	50	53	44	57	39	28	13	1	-
Male	29	33	33	33	27	28	22	11	8	4	-	-
Female	25	21	13	17	26	16	35	28	20	9	1	-
American Indian 1/, both sexes	9	3	5	5	1	-	3	2	1	-	-	-
Male	5	1	4	2	-	-	1	2	1	-	-	-
Female	4	2	1	3	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	-	1	3	1	2	-	-	-	-
Male	-	1	2	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	1	2	1	2	-	-	-	-
North Dakota												
All races, both sexes	12	17	13	14	15	21	25	33	34	17	2	-
Male	9	13	9	10	7	10	16	11	12	2	-	-
Female	3	4	4	4	8	11	9	22	22	15	2	-
White, both sexes	9	15	11	13	13	20	21	31	34	16	2	-
Male	6	12	8	9	7	10	12	11	12	2	-	-
Female	3	3	3	4	6	10	9	20	22	14	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	2	2	1	2	1	4	2	-	1	-	-
Male	3	1	1	1	-	-	4	-	-	-	-	-
Female	-	1	1	-	2	1	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Ohio												
All races, both sexes	302	282	274	291	358	444	535	470	286	105	8	-
Male	207	186	164	157	187	200	212	166	76	24	1	-
Female	95	96	110	134	171	244	323	304	210	81	7	-
White, both sexes	257	247	240	258	324	402	496	446	264	89	7	-
Male	174	165	141	139	168	181	198	154	71	22	1	-
Female	83	82	99	119	156	221	298	292	193	67	6	-
Black, both sexes	44	35	32	31	33	39	37	24	21	16	1	-
Male	32	21	22	17	19	18	14	12	5	2	-	-
Female	12	14	10	14	14	21	23	12	16	14	1	-
American Indian 1/, both sexes	1	-	1	-	1	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	2	2	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	2	2	-	1	-	-	-
Oklahoma												
All races, both sexes	97	139	115	104	118	132	157	135	109	39	6	-
Male	66	92	67	56	52	63	58	49	29	6	-	-
Female	31	47	48	48	66	69	99	86	80	33	6	-
White, both sexes	82	107	97	84	106	127	142	123	99	35	5	-
Male	59	69	55	45	46	61	53	43	25	5	-	-
Female	23	38	42	39	60	66	89	80	74	30	5	-
Black, both sexes	4	8	5	8	2	2	7	4	5	2	1	-
Male	3	4	2	5	1	1	1	2	3	1	-	-
Female	1	4	3	3	1	1	6	2	2	1	1	-
American Indian 1/, both sexes	11	24	12	12	10	3	6	8	3	2	-	-
Male	4	19	9	6	5	1	3	4	1	-	-	-
Female	7	5	3	6	5	2	3	4	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	2	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Oregon												
All races, both sexes	91	100	95	83	89	116	152	160	92	41	5	-
Male	65	71	59	41	42	61	58	59	31	10	-	-
Female	26	29	36	42	47	55	94	101	61	31	5	-
White, both sexes	89	95	89	78	87	111	148	156	91	41	5	-
Male	65	68	54	39	42	59	57	56	30	10	-	-
Female	24	27	35	39	45	52	91	100	61	31	5	-
Black, both sexes	-	-	-	-	-	3	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	2	2	2	3	-	1	2	-	-	-	-	-
Male	-	2	2	2	-	1	-	-	-	-	-	-
Female	2	-	-	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	4	2	2	1	1	2	-	-	-	-
Male	-	1	3	-	-	-	1	2	-	-	-	-
Female	-	2	1	2	2	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	264	292	237	281	414	579	709	636	398	130	33	2
Male	186	193	144	155	227	256	284	192	110	18	7	2
Female	78	99	93	126	187	323	425	444	288	112	26	-
White, both sexes	218	252	198	251	389	532	668	608	377	121	31	2
Male	154	169	128	140	217	238	269	184	102	16	7	2
Female	64	83	70	111	172	294	399	424	275	105	24	-
Black, both sexes	43	40	36	27	25	46	38	27	19	8	2	-
Male	29	24	16	14	10	18	14	7	7	2	-	-
Female	14	16	20	13	15	28	24	20	12	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	3	3	-	1	3	1	2	1	-	-
Male	3	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	3	2	-	1	2	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Rhode Island												
All races, both sexes	28	24	20	21	36	49	68	42	40	13	3	-
Male	16	17	11	11	15	22	30	15	8	1	1	-
Female	12	7	9	10	21	27	38	27	32	12	2	-
White, both sexes	25	22	18	19	35	46	68	42	40	13	3	-
Male	14	15	11	10	15	20	30	15	8	1	1	-
Female	11	7	7	9	20	26	38	27	32	12	2	-
Black, both sexes	-	2	-	1	1	2	-	-	-	-	-	-
Male	-	2	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	140	134	132	125	164	158	176	139	81	28	3	-
Male	104	89	80	68	88	71	61	45	23	6	-	-
Female	36	45	52	57	76	87	115	94	58	22	3	-
White, both sexes	94	94	97	89	128	126	143	111	64	22	2	-
Male	72	63	62	47	64	58	52	39	19	5	-	-
Female	22	31	35	42	64	68	91	72	45	17	2	-
Black, both sexes	45	39	33	35	36	31	32	28	17	5	1	-
Male	31	25	18	20	24	13	8	6	4	-	-	-
Female	14	14	15	15	12	18	24	22	13	5	1	-
American Indian 1/, both sexes	1	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
South Dakota												
All races, both sexes	17	13	16	13	13	35	44	42	20	17	3	-
Male	12	8	12	7	5	17	23	16	6	4	-	-
Female	5	5	4	6	8	18	21	26	14	13	3	-
White, both sexes	13	9	10	12	10	32	41	41	20	17	3	-
Male	9	6	8	6	5	16	23	16	6	4	-	-
Female	4	3	2	6	5	16	18	25	14	13	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	4	6	1	3	3	3	1	-	-	-	-
Male	3	2	4	1	-	1	-	-	-	-	-	-
Female	1	2	2	-	3	2	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	152	175	168	152	173	222	252	216	137	40	8	-
Male	114	128	115	83	82	99	94	70	33	5	1	-
Female	38	47	53	69	91	123	158	146	104	35	7	-
White, both sexes	127	155	144	129	149	203	232	198	127	32	6	-
Male	96	112	96	71	70	89	86	66	32	4	1	-
Female	31	43	48	58	79	114	146	132	95	28	5	-
Black, both sexes	24	17	24	23	23	19	20	18	9	8	2	-
Male	17	13	19	12	12	10	8	4	1	1	-	-
Female	7	4	5	11	11	9	12	14	8	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	-	1	-	-	-	-	-	-	-
Male	1	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Texas												
All races, both sexes	597	582	519	503	586	644	649	523	377	124	36	-
Male	434	381	336	266	328	271	284	172	102	30	3	-
Female	163	201	183	237	258	373	365	351	275	94	33	-
White, both sexes	524	508	463	451	526	581	593	482	358	111	31	-
Male	392	336	297	238	294	240	261	156	98	26	2	-
Female	132	172	166	213	232	341	332	326	260	85	29	-
Black, both sexes	67	69	49	44	50	60	49	39	17	11	2	-
Male	38	43	35	26	27	28	20	15	4	3	-	-
Female	29	26	14	18	23	32	29	24	13	8	2	-
American Indian 1/, both sexes	3	1	2	2	-	-	-	-	-	-	1	-
Male	2	-	1	1	-	-	-	-	-	-	-	-
Female	1	1	1	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	4	5	6	10	3	7	2	2	2	2	-
Male	2	2	3	1	7	3	3	1	-	1	1	-
Female	1	2	2	5	3	-	4	1	2	1	1	-
Utah												
All races, both sexes	26	30	41	43	50	49	78	57	39	6	1	1
Male	21	21	24	25	25	23	36	18	9	3	1	1
Female	5	9	17	18	25	26	42	39	30	3	-	-
White, both sexes	26	29	40	41	49	46	76	56	39	6	1	1
Male	21	20	23	25	24	22	34	17	9	3	1	1
Female	5	9	17	16	25	24	42	39	30	3	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	1	1	-	-	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Vermont												
All races, both sexes	10	10	12	11	14	24	27	25	23	17	3	-
Male	5	8	7	4	9	10	10	10	6	3	1	-
Female	5	2	5	7	5	14	17	15	17	14	2	-
White, both sexes	10	10	12	10	14	23	27	25	23	17	3	-
Male	5	8	7	4	9	9	10	10	6	3	1	-
Female	5	2	5	6	5	14	17	15	17	14	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	152	153	153	160	190	221	285	233	145	49	5	-
Male	101	108	98	88	97	104	109	96	41	11	2	-
Female	51	45	55	72	93	117	176	137	104	38	3	-
White, both sexes	105	111	115	122	142	187	236	207	127	40	3	-
Male	76	78	74	66	75	89	91	84	37	10	1	-
Female	29	33	41	56	67	98	145	123	90	30	2	-
Black, both sexes	45	41	36	35	46	30	46	25	18	9	2	-
Male	24	29	23	22	21	15	17	12	4	1	1	-
Female	21	12	13	13	25	15	29	13	14	8	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	3	2	4	3	1	-	-	-	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	3	1	4	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Washington												
All races, both sexes	112	146	117	121	145	168	228	186	158	57	12	-
Male	70	100	73	55	71	79	93	68	47	19	2	-
Female	42	46	44	66	74	89	135	118	111	38	10	-
White, both sexes	97	129	104	115	131	158	220	176	152	57	11	-
Male	59	87	68	53	64	74	91	62	47	19	1	-
Female	38	42	36	62	67	84	129	114	105	38	10	-
Black, both sexes	6	6	2	4	2	3	3	2	3	-	-	-
Male	4	6	-	-	1	3	1	2	-	-	-	-
Female	2	-	2	4	1	-	2	-	3	-	-	-
American Indian 1/, both sexes	4	6	10	-	4	2	2	1	2	-	1	-
Male	3	4	5	-	2	1	-	-	-	-	1	-
Female	1	2	5	-	2	1	2	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	1	2	8	5	3	7	1	-	-	-
Male	4	3	-	2	4	1	1	4	-	-	-	-
Female	1	2	1	-	4	4	2	3	1	-	-	-
West Virginia												
All races, both sexes	44	63	52	65	65	95	108	72	44	14	5	-
Male	23	51	27	39	32	50	37	19	9	3	-	-
Female	21	12	25	26	33	45	71	53	35	11	5	-
White, both sexes	43	60	50	63	64	93	108	68	44	13	4	-
Male	22	49	27	38	32	49	37	17	9	3	-	-
Female	21	11	23	25	32	44	71	51	35	10	4	-
Black, both sexes	1	3	2	2	1	2	-	4	-	1	1	-
Male	1	2	-	1	-	1	-	2	-	-	-	-
Female	-	1	2	1	1	1	-	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Wisconsin												
All races, both sexes	113	110	105	118	138	185	238	243	179	74	13	-
Male	73	77	54	67	71	88	87	84	51	22	1	-
Female	40	33	51	51	67	97	151	159	128	52	12	-
White, both sexes	94	97	92	110	134	180	230	239	174	72	12	-
Male	64	67	50	65	71	85	85	83	51	22	1	-
Female	30	30	42	45	63	95	145	156	123	50	11	-
Black, both sexes	10	7	10	3	4	2	6	4	4	1	1	-
Male	4	5	2	1	-	1	2	1	-	-	-	-
Female	6	2	8	2	4	1	4	3	4	1	1	-
American Indian 1/, both sexes	7	3	2	4	-	3	2	-	-	1	-	-
Male	3	3	2	1	-	2	-	-	-	-	-	-
Female	4	-	-	3	-	1	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	1	-	-	-	-	1	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	17	17	15	15	12	21	17	23	16	4	-	-
Male	14	10	12	8	7	6	7	8	6	1	-	-
Female	3	7	3	7	5	15	10	15	10	3	-	-
White, both sexes	16	14	15	14	12	20	17	23	15	4	-	-
Male	13	9	12	8	7	5	7	8	5	1	-	-
Female	3	5	3	6	5	15	10	15	10	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	-	1	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
United States												
All races, both sexes	1,974	10	7	1	2	4	4	4	6	16	28	47
Male	928	5	5	-	1	-	4	1	4	10	18	33
Female	1,046	5	2	1	1	4	-	3	2	6	10	14
White, both sexes	1,795	6	6	1	2	4	3	4	4	14	25	41
Male	837	4	4	-	1	-	3	1	3	10	15	29
Female	958	2	2	1	1	4	-	3	1	4	10	12
Black, both sexes	140	2	1	-	-	-	1	-	2	2	2	6
Male	67	1	1	-	-	-	1	-	1	-	2	4
Female	73	1	-	-	-	-	-	-	1	2	-	2
American Indian 1/, both sexes	14	1	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	1	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	36	-	-	-	-	-	-	-	-	1	2	-
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	30	-	-	-	-	-	-	-	-	1	2	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Arkansas												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	184	1	1	1	-	-	-	-	1	4	4	4
Male	99	1	1	-	-	-	-	-	-	3	2	4
Female	85	-	-	1	-	-	-	-	1	1	2	-
White, both sexes	164	1	1	1	-	-	-	-	1	4	4	4
Male	88	1	1	-	-	-	-	-	-	3	2	4
Female	76	-	-	1	-	-	-	-	1	1	2	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Colorado												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	2	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	2	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Florida												
All races, both sexes	96	-	-	-	-	-	-	-	-	-	1	1
Male	54	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	90	-	-	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	49	-	-	-	-	-	1	1	3	1	-	3
Male	22	-	-	-	-	-	1	-	3	1	-	2
Female	27	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	39	-	-	-	-	-	1	1	2	1	-	2
Male	16	-	-	-	-	-	1	-	2	1	-	1
Female	23	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Illinois												
All races, both sexes	72	-	-	-	-	-	-	-	-	1	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	69	-	-	-	-	-	-	-	-	1	-	1
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	2	3
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	43	-	-	-	-	-	-	-	-	-	2	3
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Iowa												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	3	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	3	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	21	1	-	-	-	-	-	-	-	-	2	3
Male	13	1	-	-	-	-	-	-	-	-	2	3
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	1	-	-	-	-	-	-	-	-	1	3
Male	12	1	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Kentucky												
All races, both sexes	33	-	1	-	-	-	-	-	-	-	1	2
Male	14	-	1	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	29	-	1	-	-	-	-	-	-	-	1	2
Male	12	-	1	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Massachusetts												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	47	-	1	-	-	1	-	-	-	1	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	44	-	1	-	-	1	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Minnesota												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	1	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Missouri												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Nebraska												
All races, both sexes	27	1	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	1	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	1	-	-	-	1	-
Male	7	-	-	-	-	-	1	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	1	-	-	-	1	-
Male	7	-	-	-	-	-	1	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
New Mexico												
All races, both sexes	18	1	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	85	-	-	-	-	-	-	-	-	1	1	1
Male	40	-	-	-	-	-	-	-	-	1	1	1
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	1	1	-
Male	34	-	-	-	-	-	-	-	-	1	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
North Carolina												
All races, both sexes	68	-	-	-	1	-	-	1	1	-	-	-
Male	34	-	-	-	-	-	-	1	1	-	-	-
Female	34	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	62	-	-	-	1	-	-	1	1	-	-	-
Male	30	-	-	-	-	-	-	1	1	-	-	-
Female	32	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	9	-	1	-	-	-	-	-	-	-	-	2
Male	8	-	1	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	-	-	-	-	-	-	2
Male	8	-	1	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Ohio												
All races, both sexes	103	-	-	-	-	1	-	-	-	-	3	3
Male	39	-	-	-	-	-	-	-	-	-	2	2
Female	64	-	-	-	-	1	-	-	-	-	1	1
White, both sexes	90	-	-	-	-	1	-	-	-	-	2	3
Male	33	-	-	-	-	-	-	-	-	-	1	2
Female	57	-	-	-	-	1	-	-	-	-	1	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	39	1	1	-	-	-	-	-	-	-	1	3
Male	17	1	-	-	-	-	-	-	-	-	1	3
Female	22	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	34	1	1	-	-	-	-	-	-	-	1	3
Male	16	1	-	-	-	-	-	-	-	-	1	3
Female	18	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Oregon												
All races, both sexes	47	-	-	-	-	1	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	1	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	84	-	-	-	-	-	-	-	-	-	-	3
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	79	-	-	-	-	-	-	-	-	-	-	2
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Rhode Island												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	29	1	-	-	-	-	-	-	-	1	-	-
Male	15	1	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	23	1	-	-	-	-	-	-	-	-	-	-
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
South Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	55	1	-	-	-	-	-	-	-	1	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	32	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	1	2
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Texas												
All races, both sexes	143	2	1	-	1	-	2	1	-	2	1	2
Male	59	1	1	-	1	-	2	-	-	2	-	1
Female	84	1	-	-	-	-	-	1	-	-	1	1
White, both sexes	129	1	1	-	1	-	1	1	-	2	1	-
Male	52	-	1	-	1	-	1	-	-	2	-	-
Female	77	1	-	-	-	-	-	1	-	-	1	-
Black, both sexes	14	1	-	-	-	-	1	-	-	-	-	2
Male	7	1	-	-	-	-	1	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	54	1	-	-	-	-	-	-	-	2	1	-
Male	26	-	-	-	-	-	-	-	-	1	1	-
Female	28	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	2	1	-
Male	23	-	-	-	-	-	-	-	-	1	1	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Washington												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	1
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of esophagus (K20-K22)												
Wisconsin												
All races, both sexes	48	-	-	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	47	-	-	-	-	-	-	-	-	1	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
United States												
All races, both sexes	53	59	77	73	109	212	384	401	320	134	23	-
Male	37	36	46	45	49	121	184	164	124	38	3	-
Female	16	23	31	28	60	91	200	237	196	96	20	-
White, both sexes	42	49	68	56	92	194	353	383	301	126	21	-
Male	29	29	40	33	42	111	174	155	113	38	3	-
Female	13	20	28	23	50	83	179	228	188	88	18	-
Black, both sexes	11	9	6	12	15	14	25	12	11	7	2	-
Male	8	6	4	8	6	6	7	6	6	-	-	-
Female	3	3	2	4	9	8	18	6	5	7	2	-
American Indian 1/, both sexes	-	1	-	1	1	1	3	2	2	1	-	-
Male	-	1	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	1	-	-	3	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	4	1	3	3	4	6	-	-	-
Male	-	-	2	4	-	3	3	2	4	-	-	-
Female	-	-	1	-	1	-	-	2	2	-	-	-
Alabama												
All races, both sexes	2	2	2	2	2	3	11	5	3	1	-	-
Male	1	1	1	2	1	2	3	2	-	-	-	-
Female	1	1	1	-	1	1	8	3	3	1	-	-
White, both sexes	2	-	2	-	2	3	10	4	3	1	-	-
Male	1	-	1	-	1	2	3	1	-	-	-	-
Female	1	-	1	-	1	1	7	3	3	1	-	-
Black, both sexes	-	2	-	2	-	-	1	1	-	-	-	-
Male	-	1	-	2	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Alaska												
All races, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	1	1	-	-	2	6	2	3	2	1	-
Male	-	1	-	-	-	2	2	1	2	-	1	-
Female	1	-	1	-	-	-	4	1	1	2	-	-
White, both sexes	1	1	1	-	-	2	6	2	3	2	1	-
Male	-	1	-	-	-	2	2	1	2	-	1	-
Female	1	-	1	-	-	-	4	1	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Arkansas												
All races, both sexes	2	-	1	-	2	2	6	1	2	1	-	-
Male	2	-	-	-	-	1	1	-	1	1	-	-
Female	-	-	1	-	2	1	5	1	1	-	-	-
White, both sexes	2	-	1	-	-	1	5	1	2	1	-	-
Male	2	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	1	-	-	1	5	1	1	-	-	-
Black, both sexes	-	-	-	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	7	3	4	6	18	33	34	37	18	3	-
Male	4	3	2	2	3	10	18	16	21	8	1	-
Female	1	4	1	2	3	8	15	18	16	10	2	-
White, both sexes	4	6	1	3	5	16	29	33	32	16	3	-
Male	3	3	1	1	3	10	15	15	17	8	1	-
Female	1	3	-	2	2	6	14	18	15	8	2	-
Black, both sexes	1	1	-	-	1	2	1	-	1	2	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	2	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	3	1	4	-	-	-
Male	-	-	1	1	-	-	3	1	3	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Colorado												
All races, both sexes	-	-	-	1	2	4	6	10	8	2	-	-
Male	-	-	-	-	1	3	3	3	3	1	-	-
Female	-	-	-	1	1	1	3	7	5	1	-	-
White, both sexes	-	-	-	1	1	4	6	10	7	2	-	-
Male	-	-	-	-	-	3	3	3	2	1	-	-
Female	-	-	-	1	1	1	3	7	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	2	1	-	4	1	5	9	1	-	-
Male	1	1	-	1	-	1	-	2	3	-	-	-
Female	-	-	2	-	-	3	1	3	6	1	-	-
White, both sexes	1	1	2	1	-	4	1	5	9	1	-	-
Male	1	1	-	1	-	1	-	2	3	-	-	-
Female	-	-	2	-	-	3	1	3	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Delaware												
All races, both sexes	2	-	-	-	-	1	1	2	1	1	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	2	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	1	-	-	-	-	1	1	2	1	1	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	-	1	-	-	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Florida												
All races, both sexes	4	2	3	3	5	10	19	23	15	10	-	-
Male	3	2	2	3	3	7	10	10	7	5	-	-
Female	1	-	1	-	2	3	9	13	8	5	-	-
White, both sexes	3	2	3	2	5	8	19	23	14	10	-	-
Male	2	2	2	2	3	5	10	10	7	5	-	-
Female	1	-	1	-	2	3	9	13	7	5	-	-
Black, both sexes	1	-	-	1	-	2	-	-	-	-	-	-
Male	1	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Georgia												
All races, both sexes	3	-	-	3	1	6	8	8	6	4	1	-
Male	3	-	-	2	1	3	6	-	-	-	-	-
Female	-	-	-	1	-	3	2	8	6	4	1	-
White, both sexes	2	-	-	1	1	4	6	7	6	4	1	-
Male	2	-	-	1	1	2	5	-	-	-	-	-
Female	-	-	-	-	-	2	1	7	6	4	1	-
Black, both sexes	1	-	-	2	-	2	2	1	-	-	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Hawaii												
All races, both sexes	-	-	1	1	-	2	-	1	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	-	-	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	1	-	1	2	2	5	2	1	-	-
Male	-	-	1	-	1	1	-	2	1	-	-	-
Female	-	-	-	-	-	1	2	3	1	1	-	-
White, both sexes	-	-	1	-	1	1	2	5	2	1	-	-
Male	-	-	1	-	1	-	-	2	1	-	-	-
Female	-	-	-	-	-	1	2	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Illinois												
All races, both sexes	1	2	3	6	5	12	12	12	9	7	1	-
Male	-	1	2	4	4	6	5	4	4	2	-	-
Female	1	1	1	2	1	6	7	8	5	5	1	-
White, both sexes	1	1	3	6	5	11	12	11	9	7	1	-
Male	-	-	2	4	4	6	5	4	4	2	-	-
Female	1	1	1	2	1	5	7	7	5	5	1	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	4	4	2	4	5	9	7	2	-	-
Male	-	-	3	2	1	3	2	2	4	-	-	-
Female	1	1	1	2	1	1	3	7	3	2	-	-
White, both sexes	1	1	4	4	2	4	5	9	6	2	-	-
Male	-	-	3	2	1	3	2	2	4	-	-	-
Female	1	1	1	2	1	1	3	7	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Iowa												
All races, both sexes	1	-	2	-	2	6	13	7	8	3	1	-
Male	1	-	2	-	-	3	5	2	5	1	-	-
Female	-	-	-	-	2	3	8	5	3	2	1	-
White, both sexes	1	-	2	-	2	6	12	7	8	3	1	-
Male	1	-	2	-	-	3	5	2	5	1	-	-
Female	-	-	-	-	2	3	7	5	3	2	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	1	-	-	-	4	5	3	-	1	-
Male	1	-	-	-	-	-	2	2	1	-	1	-
Female	-	-	1	-	-	-	2	3	2	-	-	-
White, both sexes	1	-	1	-	-	-	4	5	3	-	1	-
Male	1	-	-	-	-	-	2	2	1	-	1	-
Female	-	-	1	-	-	-	2	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Kentucky												
All races, both sexes	1	1	1	2	1	4	6	5	5	2	1	-
Male	-	1	-	1	-	4	4	-	2	-	-	-
Female	1	-	1	1	1	-	2	5	3	2	1	-
White, both sexes	1	1	1	2	-	4	5	4	4	2	1	-
Male	-	1	-	1	-	4	3	-	1	-	-	-
Female	1	-	1	1	-	-	2	4	3	2	1	-
Black, both sexes	-	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	1	3	1	2	5	5	3	-	-	-
Male	-	1	1	1	-	2	1	3	1	-	-	-
Female	1	-	-	2	1	-	4	2	2	-	-	-
White, both sexes	-	1	1	2	-	2	5	4	3	-	-	-
Male	-	1	1	1	-	2	1	3	1	-	-	-
Female	-	-	-	1	-	-	4	1	2	-	-	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Maine												
All races, both sexes	1	-	1	-	2	2	7	6	1	1	-	-
Male	1	-	-	-	-	-	3	4	-	-	-	-
Female	-	-	1	-	2	2	4	2	1	1	-	-
White, both sexes	1	-	1	-	2	2	7	6	1	1	-	-
Male	1	-	-	-	-	-	3	4	-	-	-	-
Female	-	-	1	-	2	2	4	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	-	1	-	2	6	4	5	-	-	-
Male	-	-	-	1	-	-	3	3	2	-	-	-
Female	-	1	-	-	-	2	3	1	3	-	-	-
White, both sexes	-	1	-	-	-	2	3	4	5	-	-	-
Male	-	-	-	-	-	-	3	3	2	-	-	-
Female	-	1	-	-	-	2	-	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Massachusetts												
All races, both sexes	-	4	-	-	3	6	8	9	5	3	2	-
Male	-	3	-	-	1	3	4	3	1	1	-	-
Female	-	1	-	-	2	3	4	6	4	2	2	-
White, both sexes	-	4	-	-	2	6	7	9	5	3	2	-
Male	-	3	-	-	1	3	4	3	1	1	-	-
Female	-	1	-	-	1	3	3	6	4	2	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	1	2	3	10	9	8	6	1	1	-
Male	2	-	1	-	1	3	5	3	2	-	-	-
Female	-	-	-	2	2	7	4	5	4	1	1	-
White, both sexes	1	-	1	2	3	10	8	7	6	1	1	-
Male	1	-	1	-	1	3	4	2	2	-	-	-
Female	-	-	-	2	2	7	4	5	4	1	1	-
Black, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Minnesota												
All races, both sexes	1	-	3	1	2	3	5	6	9	5	1	-
Male	1	-	2	-	1	1	3	2	5	1	-	-
Female	-	-	1	1	1	2	2	4	4	4	1	-
White, both sexes	1	-	3	1	2	3	5	6	9	5	1	-
Male	1	-	2	-	1	1	3	2	5	1	-	-
Female	-	-	1	1	1	2	2	4	4	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	1	1	3	-	2	2	4	2	1	-
Male	-	1	1	1	1	-	-	1	2	-	-	-
Female	-	-	-	-	2	-	2	1	2	2	1	-
White, both sexes	-	1	1	1	2	-	1	2	3	1	-	-
Male	-	1	1	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	2	-	1	1	2	1	-	-
Black, both sexes	-	-	-	-	1	-	1	-	1	1	1	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Missouri												
All races, both sexes	1	-	4	1	-	3	12	12	8	2	-	-
Male	1	-	2	1	-	2	6	3	3	2	-	-
Female	-	-	2	-	-	1	6	9	5	-	-	-
White, both sexes	1	-	3	1	-	3	12	12	6	2	-	-
Male	1	-	2	1	-	2	6	3	3	2	-	-
Female	-	-	1	-	-	1	6	9	3	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	1	1	3	2	1	1	-	-
Male	-	-	1	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	1	-	-
White, both sexes	-	-	1	-	1	1	3	2	1	1	-	-
Male	-	-	1	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Nebraska												
All races, both sexes	-	-	-	2	2	3	5	8	4	2	-	-
Male	-	-	-	1	2	2	4	6	1	-	-	-
Female	-	-	-	1	-	1	1	2	3	2	-	-
White, both sexes	-	-	-	2	2	3	5	8	4	2	-	-
Male	-	-	-	1	2	2	4	6	1	-	-	-
Female	-	-	-	1	-	1	1	2	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	-	2	1	2	1	-	-	-
Male	-	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	-	2	1	2	1	-	-	-
Male	-	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
New Hampshire												
All races, both sexes	1	1	-	-	1	3	4	1	4	2	-	-
Male	1	1	-	-	-	-	4	1	2	-	-	-
Female	-	-	-	-	1	3	-	-	2	2	-	-
White, both sexes	1	1	-	-	1	3	4	1	4	2	-	-
Male	1	1	-	-	-	-	4	1	2	-	-	-
Female	-	-	-	-	1	3	-	-	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	1	1	2	-	6	10	5	8	2	-	-
Male	3	-	-	-	-	4	5	3	3	-	-	-
Female	-	1	1	2	-	2	5	2	5	2	-	-
White, both sexes	3	1	1	1	-	6	8	5	8	2	-	-
Male	3	-	-	-	-	4	5	3	3	-	-	-
Female	-	1	1	1	-	2	3	2	5	2	-	-
Black, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
New Mexico												
All races, both sexes	-	-	-	1	1	3	4	3	3	1	-	-
Male	-	-	-	-	1	2	1	2	-	1	-	-
Female	-	-	-	1	-	1	3	1	3	-	-	-
White, both sexes	-	-	-	1	1	3	4	3	3	1	-	-
Male	-	-	-	-	1	2	1	2	-	1	-	-
Female	-	-	-	1	-	1	3	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	4	5	1	4	13	16	24	9	4	1	-
Male	1	3	4	1	1	8	8	8	2	1	-	-
Female	-	1	1	-	3	5	8	16	7	3	1	-
White, both sexes	-	3	3	1	4	11	14	22	9	3	1	-
Male	-	3	2	1	1	8	7	7	2	1	-	-
Female	-	-	1	-	3	3	7	15	7	2	1	-
Black, both sexes	1	1	2	-	-	2	2	-	-	1	-	-
Male	1	-	2	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	2	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
North Carolina												
All races, both sexes	-	1	3	5	9	7	14	16	8	2	-	-
Male	-	-	3	5	3	4	7	9	1	-	-	-
Female	-	1	-	-	6	3	7	7	7	2	-	-
White, both sexes	-	1	3	4	8	6	14	14	8	1	-	-
Male	-	-	3	4	3	3	7	7	1	-	-	-
Female	-	1	-	-	5	3	7	7	7	1	-	-
Black, both sexes	-	-	-	1	1	1	-	1	-	1	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	2	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	2	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Ohio												
All races, both sexes	3	3	6	4	9	9	13	23	14	11	1	-
Male	3	1	2	3	2	4	4	10	4	2	-	-
Female	-	2	4	1	7	5	9	13	10	9	1	-
White, both sexes	1	3	6	1	8	8	12	22	13	9	1	-
Male	1	1	2	1	2	4	4	9	4	2	-	-
Female	-	2	4	-	6	4	8	13	9	7	1	-
Black, both sexes	2	-	-	2	1	1	1	1	1	2	-	-
Male	2	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	1	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	2	-	1	1	2	6	5	8	4	1	-
Male	1	2	-	-	-	2	3	1	3	-	-	-
Female	2	-	-	1	1	-	3	4	5	4	1	-
White, both sexes	3	1	-	-	1	2	6	4	7	3	1	-
Male	1	1	-	-	-	2	3	1	3	-	-	-
Female	2	-	-	-	1	-	3	3	4	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	1	1	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Oregon												
All races, both sexes	-	2	6	-	2	2	7	13	9	4	1	-
Male	-	1	4	-	-	2	4	4	3	2	-	-
Female	-	1	2	-	2	-	3	9	6	2	1	-
White, both sexes	-	2	6	-	2	2	6	13	8	4	1	-
Male	-	1	4	-	-	2	4	4	2	2	-	-
Female	-	1	2	-	2	-	2	9	6	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	5	1	4	3	13	20	18	13	3	1	-
Male	-	2	-	4	2	7	11	9	6	1	-	-
Female	-	3	1	-	1	6	9	9	7	2	1	-
White, both sexes	-	5	1	4	2	12	20	17	12	3	1	-
Male	-	2	-	4	1	7	11	8	5	1	-	-
Female	-	3	1	-	1	5	9	9	7	2	1	-
Black, both sexes	-	-	-	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Rhode Island												
All races, both sexes	-	1	-	1	1	1	3	-	3	1	-	-
Male	-	1	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	3	1	-	-
White, both sexes	-	1	-	1	1	-	3	-	3	1	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	1	5	2	5	5	5	1	-	-
Male	1	1	1	-	3	1	3	-	4	-	-	-
Female	-	-	-	1	2	1	2	5	1	1	-	-
White, both sexes	1	1	1	1	3	2	3	5	4	1	-	-
Male	1	1	1	-	1	1	2	-	3	-	-	-
Female	-	-	-	1	2	1	1	5	1	1	-	-
Black, both sexes	-	-	-	-	2	-	2	-	1	-	-	-
Male	-	-	-	-	2	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	5	3	1	1	-	-
Male	-	-	-	-	-	-	4	1	1	-	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
White, both sexes	-	-	-	-	-	-	4	3	1	1	-	-
Male	-	-	-	-	-	-	4	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	2	2	1	2	2	9	12	13	3	1	-
Male	3	-	-	-	1	2	5	5	4	1	-	-
Female	-	2	2	1	1	-	4	7	9	2	1	-
White, both sexes	2	2	2	1	1	2	9	11	13	3	-	-
Male	2	-	-	-	-	2	5	4	4	1	-	-
Female	-	2	2	1	1	-	4	7	9	2	-	-
Black, both sexes	1	-	-	-	1	-	-	1	-	-	1	-
Male	1	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Texas												
All races, both sexes	3	6	8	4	12	11	28	26	21	10	2	-
Male	2	4	4	3	6	3	13	9	5	2	-	-
Female	1	2	4	1	6	8	15	17	16	8	2	-
White, both sexes	2	5	7	3	10	11	26	24	21	10	2	-
Male	2	3	4	2	5	3	12	9	5	2	-	-
Female	-	2	3	1	5	8	14	15	16	8	2	-
Black, both sexes	1	1	1	1	2	-	2	2	-	-	-	-
Male	-	1	-	1	1	-	1	-	-	-	-	-
Female	1	-	1	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	1	-	-	1	1	2	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	1	-	-	1	-	-	1	1	2	1	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Vermont												
All races, both sexes	1	-	-	-	1	2	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	1	-	-	-	1	2	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	1	1	4	8	12	12	7	4	-	-
Male	-	1	-	1	2	6	6	6	1	1	-	-
Female	-	-	1	-	2	2	6	6	6	3	-	-
White, both sexes	-	-	1	-	3	7	10	12	6	4	-	-
Male	-	-	-	-	2	5	6	6	1	1	-	-
Female	-	-	1	-	1	2	4	6	5	3	-	-
Black, both sexes	-	1	-	1	1	1	2	-	1	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Washington												
All races, both sexes	1	3	3	3	5	3	12	17	10	4	-	-
Male	-	2	3	1	5	3	5	5	6	1	-	-
Female	1	1	-	2	-	-	7	12	4	3	-	-
White, both sexes	1	2	3	3	5	3	12	17	10	4	-	-
Male	-	1	3	1	5	3	5	5	6	1	-	-
Female	1	1	-	2	-	-	7	12	4	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	-	2	1	2	3	4	2	-	1	-
Male	-	1	-	1	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	2	3	2	-	1	-
White, both sexes	-	1	-	2	1	2	3	3	2	-	1	-
Male	-	1	-	1	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	2	2	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Wisconsin												
All races, both sexes	-	-	2	1	1	7	9	11	13	1	-	-
Male	-	-	2	1	-	3	1	7	5	1	-	-
Female	-	-	-	-	1	4	8	4	8	-	-	-
White, both sexes	-	-	1	1	1	7	9	11	13	1	-	-
Male	-	-	1	1	-	3	1	7	5	1	-	-
Female	-	-	-	-	1	4	8	4	8	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	2	1	2	1	-	-
Male	1	-	-	-	-	-	1	1	2	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	2	1	1	1	-	-
Male	1	-	-	-	-	-	1	1	1	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
United States												
All races, both sexes	3,478	2	1	-	1	1	7	11	14	31	77	115
Male	1,722	1	-	-	1	-	6	7	10	19	54	85
Female	1,756	1	1	-	-	1	1	4	4	12	23	30
White, both sexes	2,995	1	-	-	1	1	3	8	9	23	54	81
Male	1,453	1	-	-	1	-	3	6	6	13	37	58
Female	1,542	-	-	-	-	1	-	2	3	10	17	23
Black, both sexes	363	1	1	-	-	-	4	3	5	7	21	30
Male	211	-	-	-	-	-	3	1	4	5	15	24
Female	152	1	1	-	-	-	1	2	1	2	6	6
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	108	-	-	-	-	-	-	-	-	-	2	4
Male	52	-	-	-	-	-	-	-	-	-	2	3
Female	56	-	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	46	-	-	-	-	-	-	1	-	-	1	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	19	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	1	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	1	3
Male	33	-	-	-	-	-	-	-	-	-	1	3
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	1	3
Male	32	-	-	-	-	-	-	-	-	-	1	3
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Arkansas												
All races, both sexes	41	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	597	-	-	-	-	-	-	2	-	7	11	17
Male	319	-	-	-	-	-	-	2	-	4	8	12
Female	278	-	-	-	-	-	-	-	-	3	3	5
White, both sexes	478	-	-	-	-	-	-	2	-	7	7	11
Male	259	-	-	-	-	-	-	2	-	4	5	8
Female	219	-	-	-	-	-	-	-	-	3	2	3
Black, both sexes	40	-	-	-	-	-	-	-	-	-	2	5
Male	20	-	-	-	-	-	-	-	-	-	1	3
Female	20	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	76	-	-	-	-	-	-	-	-	-	2	1
Male	38	-	-	-	-	-	-	-	-	-	2	1
Female	38	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Colorado												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	3	1
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	25	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	3	1
Male	16	-	-	-	-	-	-	-	-	-	2	1
Female	23	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Florida												
All races, both sexes	222	-	-	-	1	-	-	-	3	1	6	8
Male	116	-	-	-	1	-	-	-	2	-	4	5
Female	106	-	-	-	-	-	-	-	1	1	2	3
White, both sexes	201	-	-	-	1	-	-	-	2	1	5	6
Male	107	-	-	-	1	-	-	-	1	-	4	4
Female	94	-	-	-	-	-	-	-	1	1	1	2
Black, both sexes	20	-	-	-	-	-	-	-	1	-	1	2
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	95	-	-	-	-	-	-	1	1	1	1	4
Male	53	-	-	-	-	-	-	1	1	1	1	4
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	1	-	3
Male	38	-	-	-	-	-	-	-	-	1	-	3
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	1	1	-	1	1
Male	15	-	-	-	-	-	-	1	1	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Hawaii												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Illinois												
All races, both sexes	147	-	-	-	-	-	-	-	-	-	3	7
Male	65	-	-	-	-	-	-	-	-	-	2	6
Female	82	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	126	-	-	-	-	-	-	-	-	-	1	6
Male	52	-	-	-	-	-	-	-	-	-	-	5
Female	74	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	21	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	2	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	66	-	-	-	-	-	-	-	1	-	1	2
Male	31	-	-	-	-	-	-	-	1	-	1	1
Female	35	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	61	-	-	-	-	-	-	-	1	-	1	1
Male	28	-	-	-	-	-	-	-	1	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Iowa												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	28	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Kentucky												
All races, both sexes	47	1	-	-	-	-	-	-	-	1	1	-
Male	21	1	-	-	-	-	-	-	-	1	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	1	-	-	-	-	-	-	-	1	1	-
Male	20	1	-	-	-	-	-	-	-	1	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	48	1	-	-	-	-	2	-	3	-	4	-
Male	18	-	-	-	-	-	2	-	2	-	3	-
Female	30	1	-	-	-	-	-	-	1	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	1	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	21	1	-	-	-	-	2	-	2	-	2	-
Male	7	-	-	-	-	-	2	-	2	-	1	-
Female	14	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Maine												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	44	-	-	-	-	-	-	-	-	2	2	3
Male	26	-	-	-	-	-	-	-	-	1	2	3
Female	18	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	2	1
Male	19	-	-	-	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	2	-	2
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Massachusetts												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	115	-	-	-	-	-	-	-	1	-	-	7
Male	62	-	-	-	-	-	-	-	1	-	-	3
Female	53	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	98	-	-	-	-	-	-	-	1	-	-	5
Male	51	-	-	-	-	-	-	-	1	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Minnesota												
All races, both sexes	49	-	-	-	-	-	-	-	1	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Missouri												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Nebraska												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	24	-	-	-	-	-	-	-	-	2	-	2
Male	12	-	-	-	-	-	-	-	-	2	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
New Hampshire												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	86	-	-	-	-	-	-	-	-	2	3	2
Male	45	-	-	-	-	-	-	-	-	1	1	1
Female	41	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	73	-	-	-	-	-	-	-	-	2	1	1
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	38	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	2	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
New Mexico												
All races, both sexes	27	-	-	-	-	-	2	-	-	-	1	2
Male	16	-	-	-	-	-	2	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	2	-	-	-	1	2
Male	16	-	-	-	-	-	2	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	220	-	-	-	-	-	1	1	1	2	4	3
Male	120	-	-	-	-	-	1	1	1	1	3	3
Female	100	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	175	-	-	-	-	-	-	1	1	-	2	2
Male	91	-	-	-	-	-	-	1	1	-	1	2
Female	84	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	42	-	-	-	-	-	1	-	-	2	2	1
Male	27	-	-	-	-	-	1	-	-	1	2	1
Female	15	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
North Carolina												
All races, both sexes	96	-	1	-	-	-	-	-	2	1	5	10
Male	48	-	-	-	-	-	-	-	1	1	3	6
Female	48	-	1	-	-	-	-	-	1	-	2	4
White, both sexes	77	-	-	-	-	-	-	-	2	1	2	8
Male	36	-	-	-	-	-	-	-	1	1	1	5
Female	41	-	-	-	-	-	-	-	1	-	1	3
Black, both sexes	19	-	1	-	-	-	-	-	-	-	3	2
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	7	-	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Ohio												
All races, both sexes	151	-	-	-	-	-	-	-	-	1	2	6
Male	70	-	-	-	-	-	-	-	-	1	2	5
Female	81	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	140	-	-	-	-	-	-	-	-	1	2	5
Male	65	-	-	-	-	-	-	-	-	1	2	4
Female	75	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	53	-	-	-	-	-	-	1	-	-	2	-
Male	28	-	-	-	-	-	-	-	-	-	2	-
Female	25	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	2	-
Male	26	-	-	-	-	-	-	-	-	-	2	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Oregon												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	2	2
Male	32	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	56	-	-	-	-	-	-	-	-	-	2	2
Male	32	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	148	-	-	-	-	1	1	1	1	1	5	2
Male	61	-	-	-	-	-	-	1	1	-	3	2
Female	87	-	-	-	-	1	1	-	-	1	2	-
White, both sexes	135	-	-	-	-	1	-	1	1	1	5	1
Male	57	-	-	-	-	-	-	1	1	-	3	1
Female	78	-	-	-	-	1	-	-	-	1	2	-
Black, both sexes	12	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	65	-	-	-	-	-	-	1	-	-	2	2
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	40	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	61	-	-	-	-	-	-	1	-	-	2	2
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Texas												
All races, both sexes	175	-	-	-	-	-	1	1	-	3	5	6
Male	83	-	-	-	-	-	1	-	-	2	3	4
Female	92	-	-	-	-	-	-	1	-	1	2	2
White, both sexes	150	-	-	-	-	-	1	1	-	2	3	4
Male	66	-	-	-	-	-	1	-	-	1	2	3
Female	84	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	24	-	-	-	-	-	-	-	-	1	2	2
Male	16	-	-	-	-	-	-	-	-	1	1	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	43	-	-	-	-	-	-	-	-	1	3	2
Male	28	-	-	-	-	-	-	-	-	1	2	1
Female	15	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	32	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Washington												
All races, both sexes	59	-	-	-	-	-	-	1	-	-	1	3
Male	35	-	-	-	-	-	-	1	-	-	-	3
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	57	-	-	-	-	-	-	1	-	-	1	3
Male	33	-	-	-	-	-	-	1	-	-	-	3
Female	24	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Wisconsin												
All races, both sexes	72	-	-	-	-	-	-	-	-	2	1	2
Male	34	-	-	-	-	-	-	-	-	1	1	2
Female	38	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	2	1	-
Male	32	-	-	-	-	-	-	-	-	1	1	-
Female	38	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	1
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
United States												
All races, both sexes	166	164	212	222	322	429	611	579	376	119	18	-
Male	118	108	136	124	181	229	283	203	124	30	3	-
Female	48	56	76	98	141	200	328	376	252	89	15	-
White, both sexes	128	139	175	188	271	370	553	526	344	103	17	-
Male	88	90	114	102	151	195	263	187	112	24	2	-
Female	40	49	61	86	120	175	290	339	232	79	15	-
Black, both sexes	32	22	28	27	42	35	40	33	24	8	-	-
Male	26	18	18	17	25	21	12	11	8	3	-	-
Female	6	4	10	10	17	14	28	22	16	5	-	-
American Indian 1/, both sexes	3	1	-	-	1	3	1	2	-	-	-	-
Male	2	-	-	-	1	2	-	-	-	-	-	-
Female	1	1	-	-	-	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	9	7	8	21	17	18	8	8	1	-
Male	2	-	4	5	4	11	8	5	4	3	1	-
Female	1	2	5	2	4	10	9	13	4	5	-	-
Alabama												
All races, both sexes	2	3	3	3	7	7	5	5	5	1	-	-
Male	2	2	2	2	5	4	4	3	-	1	-	-
Female	-	1	1	1	2	3	1	2	5	-	-	-
White, both sexes	1	3	2	3	4	7	5	5	4	1	-	-
Male	1	2	1	2	3	4	4	3	-	1	-	-
Female	-	1	1	1	1	3	1	2	4	-	-	-
Black, both sexes	1	-	1	-	3	-	-	-	1	-	-	-
Male	1	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	4	2	3	9	5	12	11	8	5	-	-
Male	4	3	-	1	5	2	4	5	2	3	-	-
Female	1	1	2	2	4	3	8	6	6	2	-	-
White, both sexes	4	4	1	2	9	4	11	11	8	5	-	-
Male	3	3	-	1	5	2	4	5	2	3	-	-
Female	1	1	1	1	4	2	7	6	6	2	-	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Arkansas												
All races, both sexes	3	1	2	1	6	7	7	8	2	3	-	-
Male	2	1	1	1	3	3	1	2	-	-	-	-
Female	1	-	1	-	3	4	6	6	2	3	-	-
White, both sexes	1	1	2	1	5	7	7	7	2	2	-	-
Male	-	1	1	1	3	3	1	2	-	-	-	-
Female	1	-	1	-	2	4	6	5	2	2	-	-
Black, both sexes	2	-	-	-	1	-	-	1	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	30	21	37	40	51	79	114	90	74	21	3	-
Male	25	13	22	24	29	48	61	33	32	5	1	-
Female	5	8	15	16	22	31	53	57	42	16	2	-
White, both sexes	24	18	29	29	41	59	95	74	64	16	2	-
Male	20	12	22	14	24	37	52	29	26	4	-	-
Female	4	6	7	15	17	22	43	45	38	12	2	-
Black, both sexes	3	2	4	6	3	3	4	3	5	-	-	-
Male	3	1	-	5	2	2	1	-	2	-	-	-
Female	-	1	4	1	1	1	3	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	4	5	6	15	15	13	5	5	1	-
Male	2	-	-	5	2	8	8	4	4	1	1	-
Female	1	1	4	-	4	7	7	9	1	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Colorado												
All races, both sexes	3	1	3	-	4	9	7	4	5	2	-	-
Male	3	-	2	-	1	2	3	1	2	-	-	-
Female	-	1	1	-	3	7	4	3	3	2	-	-
White, both sexes	2	1	3	-	4	9	6	4	5	1	-	-
Male	2	-	2	-	1	2	3	1	2	-	-	-
Female	-	1	1	-	3	7	3	3	3	1	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	2	3	2	5	5	10	6	4	1	-	-
Male	-	-	3	-	3	3	6	1	1	1	-	-
Female	-	2	-	2	2	2	4	5	3	-	-	-
White, both sexes	-	2	2	2	4	5	10	6	4	1	-	-
Male	-	-	2	-	2	3	6	1	1	1	-	-
Female	-	2	-	2	2	2	4	5	3	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Delaware												
All races, both sexes	-	1	-	-	2	1	-	1	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	1	-	-	-
White, both sexes	-	1	-	-	2	1	-	1	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	2	3	-	-	-	1	-	-	-	-
Male	-	2	2	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	2	2	2	-	-	-	-	-	-	-	-
Male	-	2	2	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Florida												
All races, both sexes	9	9	17	17	20	29	42	32	19	9	-	-
Male	6	8	13	11	9	18	16	16	4	3	-	-
Female	3	1	4	6	11	11	26	16	15	6	-	-
White, both sexes	8	8	16	15	18	27	38	30	18	8	-	-
Male	6	7	12	10	8	17	15	15	4	3	-	-
Female	2	1	4	5	10	10	23	15	14	5	-	-
Black, both sexes	1	1	1	2	2	2	3	2	1	1	-	-
Male	-	1	1	1	1	1	1	1	-	-	-	-
Female	1	-	-	1	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	6	4	5	5	11	11	22	15	6	1	1	-
Male	4	4	3	2	8	7	10	6	1	-	-	-
Female	2	-	2	3	3	4	12	9	5	1	1	-
White, both sexes	4	3	4	3	7	8	19	13	6	1	1	-
Male	2	3	3	1	6	5	9	4	1	-	-	-
Female	2	-	1	2	1	3	10	9	5	1	1	-
Black, both sexes	2	1	1	2	4	3	3	2	-	-	-	-
Male	2	1	-	1	2	2	1	2	-	-	-	-
Female	-	-	1	1	2	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Hawaii												
All races, both sexes	-	2	1	1	-	-	-	4	1	2	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	2	1	-	-	-	-	4	1	1	-	-
White, both sexes	-	1	1	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	4	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	4	1	-	-	-
Idaho												
All races, both sexes	2	1	1	3	-	3	5	5	1	1	-	-
Male	1	1	-	3	-	2	2	4	-	1	-	-
Female	1	-	1	-	-	1	3	1	1	-	-	-
White, both sexes	2	1	1	3	-	3	5	5	1	1	-	-
Male	1	1	-	3	-	2	2	4	-	1	-	-
Female	1	-	1	-	-	1	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Illinois												
All races, both sexes	5	9	15	8	11	16	23	26	17	6	1	-
Male	4	7	8	6	7	5	12	2	5	1	-	-
Female	1	2	7	2	4	11	11	24	12	5	1	-
White, both sexes	3	6	12	6	10	15	22	23	15	6	1	-
Male	2	5	6	4	6	5	11	2	5	1	-	-
Female	1	1	6	2	4	10	11	21	10	5	1	-
Black, both sexes	2	3	3	2	1	1	1	3	2	-	-	-
Male	2	2	2	2	1	-	1	-	-	-	-	-
Female	-	1	1	-	-	1	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	3	-	3	7	10	14	14	7	1	-	-
Male	1	3	-	1	5	5	5	5	3	-	-	-
Female	2	-	-	2	2	5	9	9	4	1	-	-
White, both sexes	3	3	-	3	7	8	13	13	7	1	-	-
Male	1	3	-	1	5	4	5	4	3	-	-	-
Female	2	-	-	2	2	4	8	9	4	1	-	-
Black, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Iowa												
All races, both sexes	2	1	5	3	4	11	13	9	9	-	1	-
Male	1	-	3	1	1	4	7	6	3	-	-	-
Female	1	1	2	2	3	7	6	3	6	-	1	-
White, both sexes	1	1	5	3	4	11	13	9	9	-	1	-
Male	1	-	3	1	1	4	7	6	3	-	-	-
Female	-	1	2	2	3	7	6	3	6	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	4	2	4	2	3	13	8	3	1	1	-
Male	-	1	-	1	1	2	5	4	-	-	-	-
Female	-	3	2	3	1	1	8	4	3	1	1	-
White, both sexes	-	3	1	4	2	3	13	8	3	1	1	-
Male	-	1	-	1	1	2	5	4	-	-	-	-
Female	-	2	1	3	1	1	8	4	3	1	1	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Kentucky												
All races, both sexes	2	1	1	4	1	7	8	13	6	-	1	-
Male	2	-	1	2	-	5	3	2	2	-	1	-
Female	-	1	-	2	1	2	5	11	4	-	-	-
White, both sexes	2	1	1	4	1	6	8	11	5	-	1	-
Male	2	-	1	2	-	4	3	2	2	-	1	-
Female	-	1	-	2	1	2	5	9	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	5	3	2	7	8	6	4	1	-	-
Male	1	1	3	-	1	3	-	1	-	1	-	-
Female	-	-	2	3	1	4	8	5	4	-	-	-
White, both sexes	1	1	4	2	2	5	2	4	2	1	-	-
Male	1	1	2	-	1	2	-	1	-	1	-	-
Female	-	-	2	2	1	3	2	3	2	-	-	-
Black, both sexes	-	-	1	1	-	2	6	2	2	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	6	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Maine												
All races, both sexes	-	2	2	-	1	2	4	4	2	-	-	-
Male	-	1	2	-	-	2	2	2	2	-	-	-
Female	-	1	-	-	1	-	2	2	-	-	-	-
White, both sexes	-	2	2	-	1	2	4	3	2	-	-	-
Male	-	1	2	-	-	2	2	1	2	-	-	-
Female	-	1	-	-	1	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	3	4	1	1	9	7	8	4	-	-	-
Male	-	1	4	1	-	6	3	4	1	-	-	-
Female	-	2	-	-	1	3	4	4	3	-	-	-
White, both sexes	-	3	3	1	1	8	7	7	3	-	-	-
Male	-	1	3	1	-	5	3	3	-	-	-	-
Female	-	2	-	-	1	3	4	4	3	-	-	-
Black, both sexes	-	-	1	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Massachusetts												
All races, both sexes	-	3	6	1	4	5	11	10	13	1	-	-
Male	-	2	4	-	2	3	2	4	7	-	-	-
Female	-	1	2	1	2	2	9	6	6	1	-	-
White, both sexes	-	3	6	1	3	5	11	10	13	1	-	-
Male	-	2	4	-	2	3	2	4	7	-	-	-
Female	-	1	2	1	1	2	9	6	6	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	6	7	6	9	18	14	19	17	9	2	-	-
Male	6	5	5	6	10	8	11	4	3	-	-	-
Female	-	2	1	3	8	6	8	13	6	2	-	-
White, both sexes	4	6	5	7	13	12	19	15	9	2	-	-
Male	4	5	5	5	6	6	11	4	3	-	-	-
Female	-	1	-	2	7	6	8	11	6	2	-	-
Black, both sexes	1	1	-	1	4	2	-	2	-	-	-	-
Male	1	-	-	1	3	2	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Minnesota												
All races, both sexes	1	1	2	5	5	7	7	9	8	1	-	-
Male	1	1	1	1	3	4	3	5	1	-	-	-
Female	-	-	1	4	2	3	4	4	7	1	-	-
White, both sexes	1	1	2	5	5	5	7	9	8	1	-	-
Male	1	1	1	1	3	3	3	5	1	-	-	-
Female	-	-	1	4	2	2	4	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	3	2	2	5	6	4	6	4	1	-	-
Male	-	3	1	-	3	3	3	1	2	1	-	-
Female	1	-	1	2	2	3	1	5	2	-	-	-
White, both sexes	1	2	1	1	5	4	4	4	4	-	-	-
Male	-	2	-	-	3	3	3	-	2	-	-	-
Female	1	-	1	1	2	1	1	4	2	-	-	-
Black, both sexes	-	1	1	1	-	1	-	2	-	1	-	-
Male	-	1	1	-	-	-	-	1	-	1	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Missouri												
All races, both sexes	5	3	3	4	7	4	17	10	8	-	1	-
Male	4	2	2	2	1	2	8	1	-	-	-	-
Female	1	1	1	2	6	2	9	9	8	-	1	-
White, both sexes	1	2	3	4	5	3	16	10	8	-	1	-
Male	1	1	2	2	-	1	8	1	-	-	-	-
Female	-	1	1	2	5	2	8	9	8	-	1	-
Black, both sexes	4	1	-	-	2	1	1	-	-	-	-	-
Male	3	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	1	-	1	1	2	3	3	-	-	-
Male	-	-	1	-	-	-	-	2	1	-	-	-
Female	-	1	-	-	1	1	2	1	2	-	-	-
White, both sexes	-	1	1	-	1	1	2	3	3	-	-	-
Male	-	-	1	-	-	-	-	2	1	-	-	-
Female	-	1	-	-	1	1	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Nebraska												
All races, both sexes	-	3	1	1	-	4	1	4	3	5	-	-
Male	-	1	-	1	-	1	-	2	1	1	-	-
Female	-	2	1	-	-	3	1	2	2	4	-	-
White, both sexes	-	3	1	1	-	4	-	4	3	5	-	-
Male	-	1	-	1	-	1	-	2	1	1	-	-
Female	-	2	1	-	-	3	-	2	2	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Nevada												
All races, both sexes	3	1	-	-	2	4	5	2	2	-	1	-
Male	2	-	-	-	1	3	1	-	-	-	1	-
Female	1	1	-	-	1	1	4	2	2	-	-	-
White, both sexes	3	1	-	-	2	4	4	2	2	-	1	-
Male	2	-	-	-	1	3	1	-	-	-	1	-
Female	1	1	-	-	1	1	3	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
New Hampshire												
All races, both sexes	1	-	-	2	2	2	5	4	1	-	-	-
Male	1	-	-	1	1	1	1	2	1	-	-	-
Female	-	-	-	1	1	1	4	2	-	-	-	-
White, both sexes	1	-	-	2	2	2	5	4	1	-	-	-
Male	1	-	-	1	1	1	1	2	1	-	-	-
Female	-	-	-	1	1	1	4	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8	6	5	3	8	9	15	14	5	6	-	-
Male	6	4	2	1	6	4	9	7	1	2	-	-
Female	2	2	3	2	2	5	6	7	4	4	-	-
White, both sexes	5	4	4	3	7	9	14	14	5	4	-	-
Male	4	2	1	1	5	4	8	7	1	-	-	-
Female	1	2	3	2	2	5	6	7	4	4	-	-
Black, both sexes	3	2	-	-	1	-	1	-	-	1	-	-
Male	2	2	-	-	1	-	1	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
New Mexico												
All races, both sexes	4	4	1	1	2	-	3	3	4	-	-	-
Male	3	3	-	1	2	-	1	-	1	-	-	-
Female	1	1	1	-	-	-	2	3	3	-	-	-
White, both sexes	4	4	1	1	2	-	3	3	4	-	-	-
Male	3	3	-	1	2	-	1	-	1	-	-	-
Female	1	1	1	-	-	-	2	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	11	10	18	22	28	41	34	26	5	-	-
Male	9	9	7	10	14	16	23	12	8	2	-	-
Female	4	2	3	8	8	12	18	22	18	3	-	-
White, both sexes	10	10	7	15	15	22	36	30	20	4	-	-
Male	6	8	5	8	8	13	20	10	6	2	-	-
Female	4	2	2	7	7	9	16	20	14	2	-	-
Black, both sexes	3	1	3	3	7	4	5	4	5	1	-	-
Male	3	1	2	2	6	1	3	2	2	-	-	-
Female	-	-	1	1	1	3	2	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
North Carolina												
All races, both sexes	5	4	5	6	7	10	14	13	10	2	1	-
Male	2	4	3	3	6	5	7	2	4	1	-	-
Female	3	-	2	3	1	5	7	11	6	1	1	-
White, both sexes	2	3	3	6	6	9	14	12	7	1	1	-
Male	1	3	1	3	5	4	7	2	2	-	-	-
Female	1	-	2	3	1	5	7	10	5	1	1	-
Black, both sexes	3	1	2	-	1	1	-	1	3	1	-	-
Male	1	1	2	-	1	1	-	-	2	1	-	-
Female	2	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	2	-	-	2	4	5	2	-	-
Male	1	-	-	2	-	-	2	-	3	-	-	-
Female	-	-	-	-	-	-	-	4	2	2	-	-
White, both sexes	1	-	-	2	-	-	2	2	5	2	-	-
Male	1	-	-	2	-	-	2	-	3	-	-	-
Female	-	-	-	-	-	-	-	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Ohio												
All races, both sexes	8	6	11	8	19	15	30	27	11	7	-	-
Male	6	4	7	5	11	7	12	8	1	1	-	-
Female	2	2	4	3	8	8	18	19	10	6	-	-
White, both sexes	8	6	9	8	17	14	26	26	11	7	-	-
Male	6	4	5	5	10	7	11	8	1	1	-	-
Female	2	2	4	3	7	7	15	18	10	6	-	-
Black, both sexes	-	-	2	-	2	1	4	1	-	-	-	-
Male	-	-	2	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	6	3	3	6	6	11	8	5	-	-
Male	-	1	5	2	2	2	4	3	4	3	-	-
Female	-	1	1	1	1	4	2	8	4	2	-	-
White, both sexes	-	1	5	3	2	6	6	11	7	5	-	-
Male	-	1	4	2	1	2	4	3	4	3	-	-
Female	-	-	1	1	1	4	2	8	3	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Oregon												
All races, both sexes	2	4	4	5	2	2	12	13	8	1	-	-
Male	1	3	1	5	1	2	7	8	2	-	-	-
Female	1	1	3	-	1	-	5	5	6	1	-	-
White, both sexes	2	4	4	5	2	2	11	13	8	1	-	-
Male	1	3	1	5	1	2	7	8	2	-	-	-
Female	1	1	3	-	1	-	4	5	6	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	2	5	7	19	18	23	37	15	7	-	-
Male	1	2	4	2	10	10	11	10	4	-	-	-
Female	2	-	1	5	9	8	12	27	11	7	-	-
White, both sexes	2	1	5	7	15	17	23	35	14	6	-	-
Male	-	1	4	2	10	9	11	10	4	-	-	-
Female	2	-	1	5	5	8	12	25	10	6	-	-
Black, both sexes	1	1	-	-	4	1	-	2	1	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	4	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Rhode Island												
All races, both sexes	1	-	-	2	2	-	4	-	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	2	2	-	2	-	-	1	-	-
White, both sexes	1	-	-	2	2	-	4	-	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	2	2	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	5	3	6	5	6	4	5	4	-	-	-
Male	3	3	2	3	-	2	-	3	3	-	-	-
Female	-	2	1	3	5	4	4	2	1	-	-	-
White, both sexes	1	4	3	3	4	6	1	4	4	-	-	-
Male	1	2	2	2	-	2	-	2	3	-	-	-
Female	-	2	1	1	4	4	1	2	1	-	-	-
Black, both sexes	2	1	-	3	1	-	3	1	-	-	-	-
Male	2	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	2	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
South Dakota												
All races, both sexes	-	-	-	-	-	2	2	3	1	-	1	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	1	-
White, both sexes	-	-	-	-	-	2	2	3	1	-	1	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	6	4	5	6	12	11	5	5	2	1	-
Male	3	2	4	3	3	7	1	-	1	-	-	-
Female	-	4	-	2	3	5	10	5	4	2	1	-
White, both sexes	3	6	4	4	6	10	11	5	4	2	1	-
Male	3	2	4	2	3	5	1	-	1	-	-	-
Female	-	4	-	2	3	5	10	5	3	2	1	-
Black, both sexes	-	-	-	1	-	2	-	-	1	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Texas												
All races, both sexes	9	8	10	14	22	22	23	31	16	3	1	-
Male	5	6	7	7	11	7	13	11	6	-	-	-
Female	4	2	3	7	11	15	10	20	10	3	1	-
White, both sexes	8	5	6	13	22	18	19	29	15	3	1	-
Male	5	3	3	6	11	5	11	10	5	-	-	-
Female	3	2	3	7	11	13	8	19	10	3	1	-
Black, both sexes	1	3	3	1	-	4	4	2	1	-	-	-
Male	-	3	3	1	-	2	2	1	1	-	-	-
Female	1	-	-	-	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	3	-	1	3	3	6	2	-	-	-
Male	-	-	1	-	1	1	-	3	1	-	-	-
Female	1	-	2	-	-	2	3	3	1	-	-	-
White, both sexes	1	-	3	-	1	2	3	6	2	-	-	-
Male	-	-	1	-	1	1	-	3	1	-	-	-
Female	1	-	2	-	-	1	3	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Vermont												
All races, both sexes	-	-	1	1	1	3	-	3	-	-	-	-
Male	-	-	-	-	1	2	-	2	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
White, both sexes	-	-	1	-	1	2	-	3	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	3	2	3	5	4	3	10	5	2	-	-
Male	-	2	1	2	4	3	2	7	2	1	-	-
Female	-	1	1	1	1	1	1	3	3	1	-	-
White, both sexes	-	3	1	2	2	3	2	9	5	1	-	-
Male	-	2	1	1	2	2	1	6	2	1	-	-
Female	-	1	-	1	-	1	1	3	3	-	-	-
Black, both sexes	-	-	1	1	3	1	1	1	-	1	-	-
Male	-	-	-	1	2	1	1	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Washington												
All races, both sexes	4	-	5	4	4	2	11	8	10	5	1	-
Male	4	-	5	3	3	1	6	5	3	1	-	-
Female	-	-	-	1	1	1	5	3	7	4	1	-
White, both sexes	3	-	5	4	3	2	11	8	10	5	1	-
Male	3	-	5	3	2	1	6	5	3	1	-	-
Female	-	-	-	1	1	1	5	3	7	4	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	3	3	4	1	5	7	5	1	1	-	-
Male	1	1	2	3	1	4	3	-	1	-	-	-
Female	2	2	1	1	-	1	4	5	-	1	-	-
White, both sexes	2	3	3	4	1	5	7	5	1	1	-	-
Male	-	1	2	3	1	4	3	-	1	-	-	-
Female	2	2	1	1	-	1	4	5	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Wisconsin												
All races, both sexes	7	3	3	1	5	10	11	10	10	4	3	-
Male	2	2	2	1	4	7	6	2	4	-	-	-
Female	5	1	1	-	1	3	5	8	6	4	3	-
White, both sexes	7	3	3	1	5	10	11	10	10	4	3	-
Male	2	2	2	1	4	7	6	2	4	-	-	-
Female	5	1	1	-	1	3	5	8	6	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	1	-	4	-	2	-	1	-	-
Male	1	-	-	1	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	1	-	-
White, both sexes	1	-	-	1	-	3	-	2	-	1	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
United States												
All races, both sexes	1,041	-	-	-	-	1	1	6	6	12	34	41
Male	475	-	-	-	-	-	1	3	4	6	26	32
Female	566	-	-	-	-	1	-	3	2	6	8	9
White, both sexes	877	-	-	-	-	1	1	5	4	9	20	28
Male	389	-	-	-	-	-	1	3	2	4	14	20
Female	488	-	-	-	-	1	-	2	2	5	6	8
Black, both sexes	134	-	-	-	-	-	-	1	2	2	13	11
Male	73	-	-	-	-	-	-	-	2	1	11	11
Female	61	-	-	-	-	-	-	1	-	1	2	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	15	-	-	-	-	-	-	1	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	1	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Arkansas												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	181	-	-	-	-	-	-	1	-	3	5	3
Male	93	-	-	-	-	-	-	1	-	1	4	3
Female	88	-	-	-	-	-	-	-	-	2	1	-
White, both sexes	146	-	-	-	-	-	-	1	-	3	3	2
Male	78	-	-	-	-	-	-	1	-	1	2	2
Female	68	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Colorado												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Florida												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	3	4
Male	22	-	-	-	-	-	-	-	-	-	1	2
Female	35	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	51	-	-	-	-	-	-	-	-	-	2	4
Male	21	-	-	-	-	-	-	-	-	-	1	2
Female	30	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	34	-	-	-	-	-	-	-	1	-	1	3
Male	18	-	-	-	-	-	-	-	1	-	1	3
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	1	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Illinois												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	2	3
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	34	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	44	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Iowa												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Kentucky												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	17	-	-	-	-	-	-	-	2	-	2	-
Male	6	-	-	-	-	-	-	-	1	-	1	-
Female	11	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	1	-	2	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Maine												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Massachusetts												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	38	-	-	-	-	-	-	-	1	-	-	3
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	32	-	-	-	-	-	-	-	1	-	-	3
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Minnesota												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Missouri												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Nebraska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
New Hampshire												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	23	-	-	-	-	-	-	-	-	2	1	1
Male	9	-	-	-	-	-	-	-	-	1	1	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	2	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
New Mexico												
All races, both sexes	10	-	-	-	-	-	1	-	-	-	1	2
Male	8	-	-	-	-	-	1	-	-	-	1	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	1	-	-	-	1	2
Male	8	-	-	-	-	-	1	-	-	-	1	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	1	2
Male	36	-	-	-	-	-	-	-	-	-	1	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
North Carolina												
All races, both sexes	32	-	-	-	-	-	-	-	2	-	4	2
Male	18	-	-	-	-	-	-	-	1	-	3	2
Female	14	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	28	-	-	-	-	-	-	-	2	-	2	2
Male	14	-	-	-	-	-	-	-	1	-	1	2
Female	14	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Ohio												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Oregon												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	51	-	-	-	-	1	-	1	-	1	1	-
Male	18	-	-	-	-	-	-	1	-	-	1	-
Female	33	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	46	-	-	-	-	1	-	1	-	1	1	-
Male	16	-	-	-	-	-	-	1	-	-	1	-
Female	30	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Texas												
All races, both sexes	56	-	-	-	-	-	-	1	-	2	2	3
Male	23	-	-	-	-	-	-	-	-	2	2	2
Female	33	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	44	-	-	-	-	-	-	1	-	1	1	2
Male	15	-	-	-	-	-	-	-	-	1	1	1
Female	29	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Washington												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
Wisconsin												
All races, both sexes	25	-	-	-	-	-	-	-	-	2	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	2	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
United States												
All races, both sexes	65	53	69	68	79	119	174	173	100	33	7	-
Male	45	34	35	29	38	59	76	47	33	5	2	-
Female	20	19	34	39	41	60	98	126	67	28	5	-
White, both sexes	46	44	57	55	70	102	153	157	90	28	7	-
Male	31	28	29	23	34	48	69	46	30	5	2	-
Female	15	16	28	32	36	54	84	111	60	23	5	-
Black, both sexes	16	8	10	12	8	11	16	12	9	3	-	-
Male	12	6	6	5	4	7	4	1	3	-	-	-
Female	4	2	4	7	4	4	12	11	6	3	-	-
American Indian 1/, both sexes	2	-	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	1	4	5	4	1	2	-	-
Male	1	-	-	1	-	2	3	-	-	-	-	-
Female	-	1	2	-	1	2	2	4	1	2	-	-
Alabama												
All races, both sexes	-	-	1	2	2	2	1	1	2	1	-	-
Male	-	-	1	2	2	1	1	1	-	1	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
White, both sexes	-	-	-	2	1	2	1	1	1	1	-	-
Male	-	-	-	2	1	1	1	1	-	1	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	1	-	2	-	3	2	2	2	-	-
Male	2	-	-	-	2	-	2	-	1	1	-	-
Female	-	-	1	-	-	-	1	2	1	1	-	-
White, both sexes	2	-	-	-	2	-	3	2	2	2	-	-
Male	2	-	-	-	2	-	2	-	1	1	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Arkansas												
All races, both sexes	1	-	-	1	2	1	-	5	-	1	-	-
Male	-	-	-	1	1	1	-	2	-	-	-	-
Female	1	-	-	-	1	-	-	3	-	1	-	-
White, both sexes	1	-	-	1	2	1	-	5	-	-	-	-
Male	-	-	-	1	1	1	-	2	-	-	-	-
Female	1	-	-	-	1	-	-	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	10	9	11	14	25	36	30	15	5	2	-
Male	12	6	4	6	10	14	16	9	7	-	-	-
Female	-	4	5	5	4	11	20	21	8	5	2	-
White, both sexes	10	8	7	8	12	20	30	25	11	4	2	-
Male	10	5	4	4	9	12	13	9	6	-	-	-
Female	-	3	3	4	3	8	17	16	5	4	2	-
Black, both sexes	1	1	1	2	1	1	1	1	3	-	-	-
Male	1	1	-	1	1	-	-	-	1	-	-	-
Female	-	-	1	1	-	1	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	1	3	5	4	1	1	-	-
Male	1	-	-	1	-	1	3	-	-	-	-	-
Female	-	1	1	-	1	2	2	4	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Colorado												
All races, both sexes	-	-	3	-	2	2	-	2	2	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	1	2	-	1	2	-	-	-
White, both sexes	-	-	3	-	2	2	-	2	2	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	1	2	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	1	1	-	2	2	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	2	2	-	-	-	-
White, both sexes	-	1	-	1	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Florida												
All races, both sexes	2	2	4	4	3	9	11	6	6	3	-	-
Male	-	1	2	-	1	6	5	1	2	1	-	-
Female	2	1	2	4	2	3	6	5	4	2	-	-
White, both sexes	1	2	4	3	3	9	9	6	6	2	-	-
Male	-	1	2	-	1	6	4	1	2	1	-	-
Female	1	1	2	3	2	3	5	5	4	1	-	-
Black, both sexes	1	-	-	1	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	3	1	3	3	2	4	4	3	-	1	-
Male	3	3	-	1	2	1	1	2	-	-	-	-
Female	2	-	1	2	1	1	3	2	3	-	1	-
White, both sexes	3	2	-	2	2	-	4	3	3	-	1	-
Male	1	2	-	1	2	-	1	1	-	-	-	-
Female	2	-	-	1	-	-	3	2	3	-	1	-
Black, both sexes	2	1	1	1	1	2	-	1	-	-	-	-
Male	2	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Illinois												
All races, both sexes	1	3	5	1	4	4	8	12	5	2	-	-
Male	-	2	2	1	1	1	3	2	1	-	-	-
Female	1	1	3	-	3	3	5	10	4	2	-	-
White, both sexes	1	1	5	-	4	4	8	11	5	2	-	-
Male	-	1	2	-	1	1	3	2	1	-	-	-
Female	1	-	3	-	3	3	5	9	4	2	-	-
Black, both sexes	-	2	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	-	-	2	1	6	4	2	-	-	-
Male	-	1	-	-	1	-	2	1	-	-	-	-
Female	2	-	-	-	1	1	4	3	2	-	-	-
White, both sexes	2	1	-	-	2	1	5	4	2	-	-	-
Male	-	1	-	-	1	-	2	1	-	-	-	-
Female	2	-	-	-	1	1	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Iowa												
All races, both sexes	1	1	3	1	-	5	7	5	3	-	-	-
Male	-	-	1	1	-	3	5	4	2	-	-	-
Female	1	1	2	-	-	2	2	1	1	-	-	-
White, both sexes	-	1	3	1	-	5	7	5	3	-	-	-
Male	-	-	1	1	-	3	5	4	2	-	-	-
Female	-	1	2	-	-	2	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	2	2	-	-	4	2	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	1	2	2	-	-	3	-	-	-	-	-
White, both sexes	-	-	1	2	-	-	4	2	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	1	2	-	-	3	-	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Kentucky												
All races, both sexes	1	-	-	2	1	-	1	7	2	-	1	-
Male	1	-	-	1	-	-	1	-	-	-	1	-
Female	-	-	-	1	1	-	-	7	2	-	-	-
White, both sexes	1	-	-	2	1	-	1	5	1	-	1	-
Male	1	-	-	1	-	-	1	-	-	-	1	-
Female	-	-	-	1	1	-	-	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	2	2	-	1	2	4	-	1	-	-
Male	-	1	-	-	-	1	-	1	-	1	-	-
Female	-	-	2	2	-	-	2	3	-	-	-	-
White, both sexes	-	1	2	1	-	1	-	2	-	1	-	-
Male	-	1	-	-	-	1	-	1	-	1	-	-
Female	-	-	2	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Maine												
All races, both sexes	-	1	-	-	-	-	2	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	2	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	-	-	-	3	1	3	2	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	3	2	-	-	-
White, both sexes	-	1	-	-	-	3	1	3	2	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Massachusetts												
All races, both sexes	-	-	1	-	-	-	5	3	5	-	-	-
Male	-	-	1	-	-	-	1	1	3	-	-	-
Female	-	-	-	-	-	-	4	2	2	-	-	-
White, both sexes	-	-	1	-	-	-	5	3	5	-	-	-
Male	-	-	1	-	-	-	1	1	3	-	-	-
Female	-	-	-	-	-	-	4	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	2	2	4	3	3	8	4	4	-	-	-
Male	4	2	1	2	1	2	4	-	1	-	-	-
Female	-	-	1	2	2	1	4	4	3	-	-	-
White, both sexes	2	2	1	3	3	2	8	3	4	-	-	-
Male	2	2	1	1	1	1	4	-	1	-	-	-
Female	-	-	-	2	2	1	4	3	3	-	-	-
Black, both sexes	1	-	-	1	-	1	-	1	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Minnesota												
All races, both sexes	1	-	2	-	3	2	3	2	2	-	-	-
Male	1	-	1	-	1	2	1	1	-	-	-	-
Female	-	-	1	-	2	-	2	1	2	-	-	-
White, both sexes	1	-	2	-	3	1	3	2	2	-	-	-
Male	1	-	1	-	1	1	1	1	-	-	-	-
Female	-	-	1	-	2	-	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	1	2	-	1	1	2	1	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	2	-	-	-	2	-	-	-	-
White, both sexes	-	1	1	1	-	1	1	2	1	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Missouri												
All races, both sexes	4	1	1	1	4	1	2	2	-	-	-	-
Male	3	-	1	-	-	-	2	-	-	-	-	-
Female	1	1	-	1	4	1	-	2	-	-	-	-
White, both sexes	-	1	1	1	4	1	2	2	-	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	1	-	1	4	1	-	2	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Nebraska												
All races, both sexes	-	1	-	-	-	2	-	2	1	2	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	1	-	2	-	2	-	-
White, both sexes	-	1	-	-	-	2	-	2	1	2	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	1	-	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	1	1	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
New Hampshire												
All races, both sexes	1	-	-	-	-	2	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	2	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	-	1	1	2	4	2	4	-	2	-	-
Male	2	-	-	-	1	1	1	1	-	-	-	-
Female	1	-	1	1	1	3	1	3	-	2	-	-
White, both sexes	1	-	1	1	2	4	2	4	-	2	-	-
Male	1	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	1	1	1	3	1	3	-	2	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
New Mexico												
All races, both sexes	1	2	1	-	1	-	-	-	1	-	-	-
Male	1	2	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	1	2	1	-	1	-	-	-	1	-	-	-
Male	1	2	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	4	4	7	5	5	11	5	8	1	-	-
Male	6	3	4	3	3	4	7	-	3	-	-	-
Female	-	1	-	4	2	1	4	5	5	1	-	-
White, both sexes	4	4	3	5	3	4	9	4	4	-	-	-
Male	4	3	3	2	2	3	5	-	1	-	-	-
Female	-	1	-	3	1	1	4	4	3	-	-	-
Black, both sexes	2	-	1	2	2	1	2	1	4	1	-	-
Male	2	-	1	1	1	1	2	-	2	-	-	-
Female	-	-	-	1	1	-	-	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
North Carolina												
All races, both sexes	1	2	4	1	1	2	5	4	3	1	-	-
Male	-	2	2	1	1	1	4	-	1	-	-	-
Female	1	-	2	-	-	1	1	4	2	1	-	-
White, both sexes	1	2	2	1	1	2	5	4	3	1	-	-
Male	-	2	-	1	1	1	4	-	1	-	-	-
Female	1	-	2	-	-	1	1	4	2	1	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	1	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Ohio												
All races, both sexes	3	3	4	1	5	7	11	3	2	1	-	-
Male	3	3	2	-	2	2	4	1	-	-	-	-
Female	-	-	2	1	3	5	7	2	2	1	-	-
White, both sexes	3	3	4	1	5	6	9	2	2	1	-	-
Male	3	3	2	-	2	2	4	1	-	-	-	-
Female	-	-	2	1	3	4	5	1	2	1	-	-
Black, both sexes	-	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	1	1	-	1	1	5	3	-	-	-
Male	-	-	1	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	5	2	-	-	-
White, both sexes	-	-	1	1	-	1	1	5	3	-	-	-
Male	-	-	1	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	5	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Oregon												
All races, both sexes	1	1	2	2	1	-	3	3	3	-	-	-
Male	-	-	-	2	-	-	1	2	1	-	-	-
Female	1	1	2	-	1	-	2	1	2	-	-	-
White, both sexes	1	1	2	2	1	-	2	3	3	-	-	-
Male	-	-	-	2	-	-	1	2	1	-	-	-
Female	1	1	2	-	1	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	2	2	5	11	9	11	4	2	-	-
Male	1	-	1	-	2	4	2	4	2	-	-	-
Female	-	-	1	2	3	7	7	7	2	2	-	-
White, both sexes	-	-	2	2	4	10	9	10	4	1	-	-
Male	-	-	1	-	2	3	2	4	2	-	-	-
Female	-	-	1	2	2	7	7	6	2	1	-	-
Black, both sexes	1	-	-	-	1	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Rhode Island												
All races, both sexes	-	-	-	1	1	-	3	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	1	-	-
White, both sexes	-	-	-	1	1	-	3	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	4	1	1	3	2	2	1	1	-	-	-
Male	1	2	-	1	-	1	-	-	1	-	-	-
Female	-	2	1	-	3	1	2	1	-	-	-	-
White, both sexes	1	3	1	1	2	2	-	1	1	-	-	-
Male	1	1	-	1	-	1	-	-	1	-	-	-
Female	-	2	1	-	2	1	-	1	-	-	-	-
Black, both sexes	-	1	-	-	1	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	2	3	-	3	2	1	1	1	-	-
Male	-	-	2	1	-	2	-	-	-	-	-	-
Female	-	1	-	2	-	1	2	1	1	1	-	-
White, both sexes	-	1	2	3	-	2	2	1	1	1	-	-
Male	-	-	2	1	-	1	-	-	-	-	-	-
Female	-	1	-	2	-	1	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Texas												
All races, both sexes	5	2	4	3	6	7	5	12	4	-	-	-
Male	2	2	4	-	2	2	1	3	1	-	-	-
Female	3	-	-	3	4	5	4	9	3	-	-	-
White, both sexes	4	-	2	3	6	5	4	11	4	-	-	-
Male	2	-	2	-	2	1	1	3	1	-	-	-
Female	2	-	-	3	4	4	3	8	3	-	-	-
Black, both sexes	1	2	2	-	-	2	1	1	-	-	-	-
Male	-	2	2	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	2	-	2	-	1	1	3	1	-	-	-
Male	-	1	-	1	-	-	1	2	-	-	-	-
Female	-	1	-	1	-	1	-	1	1	-	-	-
White, both sexes	-	2	-	1	-	1	-	3	1	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	1	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Washington												
All races, both sexes	-	-	1	2	1	-	1	5	3	3	-	-
Male	-	-	1	2	1	-	-	3	-	1	-	-
Female	-	-	-	-	-	-	1	2	3	2	-	-
White, both sexes	-	-	1	2	1	-	1	5	3	3	-	-
Male	-	-	1	2	1	-	-	3	-	1	-	-
Female	-	-	-	-	-	-	1	2	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	-	2	2	-	2	1	2	-	1	-	-
Male	1	-	1	1	-	1	1	-	-	-	-	-
Female	2	-	1	1	-	1	-	2	-	1	-	-
White, both sexes	2	-	2	2	-	2	1	2	-	1	-	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	2	-	1	1	-	1	-	2	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Wisconsin												
All races, both sexes	2	2	1	-	2	4	3	4	2	-	2	-
Male	-	1	1	-	1	2	2	1	1	-	-	-
Female	2	1	-	-	1	2	1	3	1	-	2	-
White, both sexes	2	2	1	-	2	4	3	4	2	-	2	-
Male	-	1	1	-	1	2	2	1	1	-	-	-
Female	2	1	-	-	1	2	1	3	1	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
United States												
All races, both sexes	1,246	2	1	-	1	-	4	3	6	13	31	49
Male	695	1	-	-	1	-	4	2	4	8	22	39
Female	551	1	1	-	-	-	-	1	2	5	9	10
White, both sexes	1,079	1	-	-	1	-	2	1	4	8	23	35
Male	589	1	-	-	1	-	2	1	3	4	18	28
Female	490	-	-	-	-	-	-	-	1	4	5	7
Black, both sexes	122	1	1	-	-	-	2	2	2	5	7	13
Male	81	-	-	-	-	-	2	1	1	4	3	10
Female	41	1	1	-	-	-	-	1	1	1	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Arkansas												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	196	-	-	-	-	-	-	-	-	2	5	8
Male	119	-	-	-	-	-	-	-	-	1	3	6
Female	77	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	156	-	-	-	-	-	-	-	-	2	3	5
Male	93	-	-	-	-	-	-	-	-	1	2	4
Female	63	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	3
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Colorado												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Florida												
All races, both sexes	93	-	-	-	1	-	-	-	2	1	3	2
Male	53	-	-	-	1	-	-	-	1	-	3	2
Female	40	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	89	-	-	-	1	-	-	-	2	1	3	2
Male	51	-	-	-	1	-	-	-	1	-	3	2
Female	38	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	33	-	-	-	-	-	-	1	-	1	-	1
Male	18	-	-	-	-	-	-	1	-	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Illinois												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	1	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	3
Male	22	-	-	-	-	-	-	-	-	-	-	3
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Iowa												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Kentucky												
All races, both sexes	15	1	-	-	-	-	-	-	-	1	-	-
Male	8	1	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	1	-	-	-	-	-	-	-	1	-	-
Male	8	1	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	13	1	-	-	-	-	2	-	1	-	1	-
Male	7	-	-	-	-	-	2	-	1	-	1	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	-	2	-	1	-	-	-
Male	4	-	-	-	-	-	2	-	1	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	20	-	-	-	-	-	-	-	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	1	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Massachusetts												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	4
Male	27	-	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Minnesota												
All races, both sexes	23	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Missouri												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Nebraska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	-	-	-	-	-	-	-	-	2	-	2
Male	7	-	-	-	-	-	-	-	-	2	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
New Hampshire												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
New Mexico												
All races, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	86	-	-	-	-	-	-	-	1	2	1	1
Male	46	-	-	-	-	-	-	-	1	1	1	1
Female	40	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	75	-	-	-	-	-	-	-	1	-	-	1
Male	38	-	-	-	-	-	-	-	1	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	2	1	-
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
North Carolina												
All races, both sexes	29	-	1	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	1	-	-	-	-	-	-	-	1	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	1	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Ohio												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	2	4
Male	33	-	-	-	-	-	-	-	-	-	2	4
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	2	3
Male	29	-	-	-	-	-	-	-	-	-	2	3
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Oregon												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	51	-	-	-	-	-	-	-	1	-	3	2
Male	25	-	-	-	-	-	-	-	1	-	2	2
Female	26	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	45	-	-	-	-	-	-	-	1	-	3	1
Male	23	-	-	-	-	-	-	-	1	-	2	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Texas												
All races, both sexes	54	-	-	-	-	-	1	-	-	-	3	1
Male	30	-	-	-	-	-	1	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	46	-	-	-	-	-	1	-	-	-	2	-
Male	25	-	-	-	-	-	1	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Washington												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	-	1
Male	15	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	1	-	-	-	1
Male	14	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Wisconsin												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
United States												
All races, both sexes	54	66	89	81	120	161	211	202	114	34	4	-
Male	44	48	67	52	81	90	108	81	34	9	-	-
Female	10	18	22	29	39	71	103	121	80	25	4	-
White, both sexes	45	57	74	72	100	137	197	184	105	29	4	-
Male	36	40	55	45	68	77	103	73	28	6	-	-
Female	9	17	19	27	32	60	94	111	77	23	4	-
Black, both sexes	7	9	10	7	15	14	9	9	5	4	-	-
Male	6	8	9	6	10	8	3	4	3	3	-	-
Female	1	1	1	1	5	6	6	5	2	1	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	5	2	4	10	5	9	4	1	-	-
Male	1	-	3	1	2	5	2	4	3	-	-	-
Female	-	-	2	1	2	5	3	5	1	1	-	-
Alabama												
All races, both sexes	1	1	2	-	-	3	1	2	2	-	-	-
Male	1	1	1	-	-	2	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	2	-	-	-
White, both sexes	-	1	2	-	-	3	1	2	2	-	-	-
Male	-	1	1	-	-	2	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	4	1	1	2	4	4	8	1	1	-	-
Male	2	3	-	-	1	2	1	5	-	-	-	-
Female	-	1	1	1	1	2	3	3	1	1	-	-
White, both sexes	1	4	1	1	2	3	4	8	1	1	-	-
Male	1	3	-	-	1	2	1	5	-	-	-	-
Female	-	1	1	1	1	1	3	3	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Arkansas												
All races, both sexes	-	-	1	-	3	5	1	-	1	1	-	-
Male	-	-	1	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	1	4	1	-	1	1	-	-
White, both sexes	-	-	1	-	3	5	1	-	1	1	-	-
Male	-	-	1	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	1	4	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	11	6	16	15	18	26	36	29	21	3	-	-
Male	7	4	12	10	12	18	24	12	9	1	-	-
Female	4	2	4	5	6	8	12	17	12	2	-	-
White, both sexes	9	6	13	11	14	19	31	24	17	2	-	-
Male	5	4	12	6	10	13	21	9	5	1	-	-
Female	4	2	1	5	4	6	10	15	12	1	-	-
Black, both sexes	1	-	1	3	-	-	1	1	1	-	-	-
Male	1	-	-	3	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	3	7	4	4	3	1	-	-
Male	1	-	-	1	1	5	2	3	3	-	-	-
Female	-	-	2	-	2	2	2	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Colorado												
All races, both sexes	2	-	-	-	-	4	4	-	3	2	-	-
Male	2	-	-	-	-	2	2	-	2	-	-	-
Female	-	-	-	-	-	2	2	-	1	2	-	-
White, both sexes	2	-	-	-	-	4	3	-	3	1	-	-
Male	2	-	-	-	-	2	2	-	2	-	-	-
Female	-	-	-	-	-	2	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	2	-	3	2	5	2	2	-	-	-
Male	-	-	2	-	2	1	4	1	-	-	-	-
Female	-	-	-	-	1	1	1	1	2	-	-	-
White, both sexes	-	-	1	-	3	2	5	2	2	-	-	-
Male	-	-	1	-	2	1	4	1	-	-	-	-
Female	-	-	-	-	1	1	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Delaware												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	2	2	-	-	-	1	-	-	-	-
Male	-	2	2	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	2	2	1	-	-	-	-	-	-	-	-
Male	-	2	2	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Florida												
All races, both sexes	3	4	10	7	7	12	13	17	7	4	-	-
Male	3	4	9	6	3	7	4	7	1	2	-	-
Female	-	-	1	1	4	5	9	10	6	2	-	-
White, both sexes	3	4	9	7	7	11	12	16	7	4	-	-
Male	3	4	8	6	3	6	4	7	1	2	-	-
Female	-	-	1	1	4	5	8	9	6	2	-	-
Black, both sexes	-	-	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	2	2	3	3	7	9	2	-	-	-
Male	1	1	1	1	2	2	3	3	1	-	-	-
Female	-	-	1	1	1	1	4	6	1	-	-	-
White, both sexes	1	1	2	1	2	3	6	8	2	-	-	-
Male	1	1	1	-	2	2	3	2	1	-	-	-
Female	-	-	1	1	-	1	3	6	1	-	-	-
Black, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Hawaii												
All races, both sexes	-	1	1	-	-	-	-	4	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	4	1	-	-	-
White, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	4	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	4	1	-	-	-
Idaho												
All races, both sexes	-	-	-	2	-	1	2	3	-	-	-	-
Male	-	-	-	2	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	2	-	1	2	3	-	-	-	-
Male	-	-	-	2	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Illinois												
All races, both sexes	2	3	6	5	5	6	8	4	2	1	-	-
Male	2	2	5	4	4	2	5	-	1	-	-	-
Female	-	1	1	1	1	4	3	4	1	1	-	-
White, both sexes	1	2	4	5	4	5	7	2	2	1	-	-
Male	1	1	3	4	3	2	4	-	1	-	-	-
Female	-	1	1	1	1	3	3	2	1	1	-	-
Black, both sexes	1	1	2	-	1	1	1	2	-	-	-	-
Male	1	1	2	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	-	1	4	6	4	1	3	-	-	-
Male	1	1	-	-	4	5	1	1	1	-	-	-
Female	-	-	-	1	-	1	3	-	2	-	-	-
White, both sexes	1	1	-	1	4	4	4	1	3	-	-	-
Male	1	1	-	-	4	4	1	1	1	-	-	-
Female	-	-	-	1	-	-	3	-	2	-	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Iowa												
All races, both sexes	-	-	2	1	2	3	5	2	3	-	-	-
Male	-	-	2	-	-	-	2	1	-	-	-	-
Female	-	-	-	1	2	3	3	1	3	-	-	-
White, both sexes	-	-	2	1	2	3	5	2	3	-	-	-
Male	-	-	2	-	-	-	2	1	-	-	-	-
Female	-	-	-	1	2	3	3	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	3	-	1	1	-	6	5	2	1	1	-
Male	-	1	-	1	1	-	1	1	-	-	-	-
Female	-	2	-	-	-	-	5	4	2	1	1	-
White, both sexes	-	3	-	1	1	-	6	5	2	1	1	-
Male	-	1	-	1	1	-	1	1	-	-	-	-
Female	-	2	-	-	-	-	5	4	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Kentucky												
All races, both sexes	-	-	1	2	-	4	4	1	1	-	-	-
Male	-	-	1	1	-	2	1	1	-	-	-	-
Female	-	-	-	1	-	2	3	-	1	-	-	-
White, both sexes	-	-	1	2	-	4	4	1	1	-	-	-
Male	-	-	1	1	-	2	1	1	-	-	-	-
Female	-	-	-	1	-	2	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	5	1	1	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	3	1	1	-	-	-	-
White, both sexes	1	-	-	-	-	3	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Maine												
All races, both sexes	-	-	1	-	-	2	1	3	-	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	1	-	-	2	1	2	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	2	3	-	-	5	4	1	1	-	-	-
Male	-	1	3	-	-	4	2	1	1	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	2	3	-	-	4	4	1	-	-	-	-
Male	-	1	3	-	-	3	2	1	-	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Massachusetts												
All races, both sexes	-	3	4	1	2	1	6	5	3	1	-	-
Male	-	2	2	-	2	1	1	2	1	-	-	-
Female	-	1	2	1	-	-	5	3	2	1	-	-
White, both sexes	-	3	4	1	2	1	6	5	3	1	-	-
Male	-	2	2	-	2	1	1	2	1	-	-	-
Female	-	1	2	1	-	-	5	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	4	2	4	6	6	4	6	3	2	-	-
Male	1	2	2	4	4	4	3	3	1	-	-	-
Female	-	2	-	-	2	2	1	3	2	2	-	-
White, both sexes	1	3	2	4	4	5	4	6	3	2	-	-
Male	1	2	2	4	2	3	3	3	1	-	-	-
Female	-	1	-	-	2	2	1	3	2	2	-	-
Black, both sexes	-	1	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Minnesota												
All races, both sexes	-	1	-	4	1	5	1	5	3	1	-	-
Male	-	1	-	1	1	2	-	3	-	-	-	-
Female	-	-	-	3	-	3	1	2	3	1	-	-
White, both sexes	-	1	-	4	1	4	1	5	3	1	-	-
Male	-	1	-	1	1	2	-	3	-	-	-	-
Female	-	-	-	3	-	2	1	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	2	-	-	2	1	3	2	2	1	-	-
Male	-	2	-	-	2	-	2	1	1	1	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	1	-	-	2	-	3	1	2	-	-	-
Male	-	1	-	-	2	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Missouri												
All races, both sexes	1	1	-	1	2	2	7	4	4	-	1	-
Male	1	1	-	-	1	1	5	-	-	-	-	-
Female	-	-	-	1	1	1	2	4	4	-	1	-
White, both sexes	1	1	-	1	-	1	7	4	4	-	1	-
Male	1	1	-	-	-	-	5	-	-	-	-	-
Female	-	-	-	1	-	1	2	4	4	-	1	-
Black, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	1	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
White, both sexes	-	-	1	-	1	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Nebraska												
All races, both sexes	-	1	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	1	-	-	-
White, both sexes	-	1	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Nevada												
All races, both sexes	1	1	-	-	1	1	4	-	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	1	-	3	-	-	-	-	-
White, both sexes	1	1	-	-	1	1	4	-	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
New Hampshire												
All races, both sexes	-	-	-	1	1	-	4	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	1	1	-	3	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	4	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	1	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	3	4	2	2	2	4	5	3	2	-	-
Male	2	2	2	1	2	1	3	3	-	1	-	-
Female	-	1	2	1	-	1	1	2	3	1	-	-
White, both sexes	2	1	3	2	2	2	4	5	3	1	-	-
Male	2	-	1	1	2	1	3	3	-	-	-	-
Female	-	1	2	1	-	1	1	2	3	1	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	1	-	-
Male	-	2	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
New Mexico												
All races, both sexes	3	2	-	-	-	-	2	2	-	-	-	-
Male	2	1	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	1	2	-	-	-	-
White, both sexes	3	2	-	-	-	-	2	2	-	-	-	-
Male	2	1	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	4	4	6	14	9	15	16	9	2	-	-
Male	1	4	3	3	10	4	9	5	2	1	-	-
Female	1	-	1	3	4	5	6	11	7	1	-	-
White, both sexes	2	4	3	6	10	7	15	15	9	2	-	-
Male	1	4	2	3	6	4	9	4	2	1	-	-
Female	1	-	1	3	4	3	6	11	7	1	-	-
Black, both sexes	-	-	1	-	4	2	-	1	-	-	-	-
Male	-	-	1	-	4	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
North Carolina												
All races, both sexes	3	2	1	2	1	3	5	3	5	1	-	-
Male	2	2	1	1	1	2	2	2	2	1	-	-
Female	1	-	-	1	-	1	3	1	3	-	-	-
White, both sexes	1	1	1	2	1	3	5	3	3	-	-	-
Male	1	1	1	1	1	2	2	2	1	-	-	-
Female	-	-	-	1	-	1	3	1	2	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	2	1	-	-
Male	1	1	-	-	-	-	-	-	1	1	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Ohio												
All races, both sexes	2	1	5	3	8	5	8	11	4	2	-	-
Male	2	1	4	3	4	4	4	4	-	1	-	-
Female	-	-	1	-	4	1	4	7	4	1	-	-
White, both sexes	2	1	3	3	7	5	6	11	4	2	-	-
Male	2	1	2	3	4	4	3	4	-	1	-	-
Female	-	-	1	-	3	1	3	7	4	1	-	-
Black, both sexes	-	-	2	-	1	-	2	-	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	4	1	-	5	2	2	2	1	-	-
Male	-	-	3	-	-	2	2	1	2	1	-	-
Female	-	-	1	1	-	3	-	1	-	-	-	-
White, both sexes	-	-	4	1	-	5	2	2	2	1	-	-
Male	-	-	3	-	-	2	2	1	2	1	-	-
Female	-	-	1	1	-	3	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Oregon												
All races, both sexes	1	1	2	1	1	-	4	3	2	-	-	-
Male	1	1	1	1	1	-	4	3	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
White, both sexes	1	1	2	1	1	-	4	3	2	-	-	-
Male	1	1	1	1	1	-	4	3	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	2	-	1	8	6	7	14	3	3	-	-
Male	-	2	-	-	5	5	6	2	-	-	-	-
Female	1	-	-	1	3	1	1	12	3	3	-	-
White, both sexes	1	1	-	1	6	6	7	13	2	3	-	-
Male	-	1	-	-	5	5	6	2	-	-	-	-
Female	1	-	-	1	1	1	1	11	2	3	-	-
Black, both sexes	-	1	-	-	2	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Rhode Island												
All races, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	1	-	2	1	2	1	3	3	-	-	-
Male	2	1	-	1	-	-	-	2	2	-	-	-
Female	-	-	-	1	1	2	1	1	1	-	-	-
White, both sexes	-	1	-	1	1	2	1	3	3	-	-	-
Male	-	1	-	1	-	-	-	2	2	-	-	-
Female	-	-	-	-	1	2	1	1	1	-	-	-
Black, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	2	-	-	3	3	3	3	1	-	-	-
Male	2	2	-	-	2	1	-	-	1	-	-	-
Female	-	-	-	-	1	2	3	3	-	-	-	-
White, both sexes	2	2	-	-	3	3	3	3	1	-	-	-
Male	2	2	-	-	2	1	-	-	1	-	-	-
Female	-	-	-	-	1	2	3	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Texas												
All races, both sexes	3	3	5	5	7	8	5	7	4	2	-	-
Male	2	2	3	3	6	4	3	5	-	-	-	-
Female	1	1	2	2	1	4	2	2	4	2	-	-
White, both sexes	3	3	3	4	7	7	4	6	4	2	-	-
Male	2	2	1	2	6	3	3	4	-	-	-	-
Female	1	1	2	2	1	4	1	2	4	2	-	-
Black, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	1	1	1	4	-	-	-	-
Male	-	-	-	-	1	-	-	3	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	1	-	1	-	1	4	-	-	-	-
Male	-	-	-	-	1	-	-	3	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Vermont												
All races, both sexes	-	-	1	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	1	1	3	2	1	3	2	-	-	-
Male	-	-	1	1	3	2	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	1	1	1	1	1	3	2	-	-	-
Male	-	-	1	1	1	1	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Washington												
All races, both sexes	3	-	3	1	3	-	5	2	1	1	1	-
Male	3	-	3	1	2	-	2	1	1	-	-	-
Female	-	-	-	-	1	-	3	1	-	1	1	-
White, both sexes	3	-	3	1	2	-	5	2	1	1	1	-
Male	3	-	3	1	1	-	2	1	1	-	-	-
Female	-	-	-	-	1	-	3	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	3	1	1	1	1	4	3	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	4	3	-	-	-	-
White, both sexes	-	3	1	1	1	1	4	3	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	4	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Wisconsin												
All races, both sexes	2	1	-	1	2	1	4	2	4	2	-	-
Male	2	1	-	1	2	1	3	1	2	-	-	-
Female	-	-	-	-	-	-	1	1	2	2	-	-
White, both sexes	2	1	-	1	2	1	4	2	4	2	-	-
Male	2	1	-	1	2	1	3	1	2	-	-	-
Female	-	-	-	-	-	-	1	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
United States												
All races, both sexes	1,164	-	-	-	-	-	2	1	2	5	12	22
Male	537	-	-	-	-	-	1	1	2	4	6	13
Female	627	-	-	-	-	-	1	-	-	1	6	9
White, both sexes	1,014	-	-	-	-	-	-	1	1	5	11	16
Male	460	-	-	-	-	-	-	1	1	4	5	9
Female	554	-	-	-	-	-	-	-	-	1	6	7
Black, both sexes	106	-	-	-	-	-	2	-	1	-	1	5
Male	57	-	-	-	-	-	1	-	1	-	1	3
Female	49	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Arkansas												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	218	-	-	-	-	-	-	1	-	2	1	6
Male	105	-	-	-	-	-	-	1	-	2	1	3
Female	113	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	174	-	-	-	-	-	-	1	-	2	1	4
Male	86	-	-	-	-	-	-	1	-	2	1	2
Female	88	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Colorado												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Florida												
All races, both sexes	71	-	-	-	-	-	-	-	1	-	-	2
Male	40	-	-	-	-	-	-	-	1	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	1	-	-	2
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Illinois												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	24	-	-	-	-	-	-	-	1	-	-	1
Male	11	-	-	-	-	-	-	-	1	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Iowa												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Kentucky												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Massachusetts												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Minnesota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Missouri												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Nebraska												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
New Mexico												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	74	-	-	-	-	-	1	-	-	-	2	-
Male	37	-	-	-	-	-	1	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	1	-	-	-	1	-
Male	8	-	-	-	-	-	1	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
North Carolina												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	-	5
Male	12	-	-	-	-	-	-	-	-	1	-	3
Female	20	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	26	-	-	-	-	-	-	-	-	1	-	4
Male	8	-	-	-	-	-	-	-	-	1	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Ohio												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	53	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Oregon												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	45	-	-	-	-	-	1	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Texas												
All races, both sexes	65	-	-	-	-	-	-	-	-	1	-	2
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	1	-	2
Male	26	-	-	-	-	-	-	-	-	-	-	2
Female	34	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Washington												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
Wisconsin												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
United States												
All races, both sexes	47	44	52	70	119	147	222	201	159	52	7	-
Male	29	26	32	41	59	80	96	74	56	16	1	-
Female	18	18	20	29	60	67	126	127	103	36	6	-
White, both sexes	37	37	42	58	97	129	199	183	146	46	6	-
Male	21	22	28	32	46	70	88	67	53	13	-	-
Female	16	15	14	26	51	59	111	116	93	33	6	-
Black, both sexes	9	5	8	8	19	10	15	12	10	1	-	-
Male	8	4	3	6	11	6	5	6	2	-	-	-
Female	1	1	5	2	8	4	10	6	8	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	4	3	7	7	5	3	5	1	-
Male	-	-	1	3	2	4	3	1	1	3	1	-
Female	1	1	1	1	1	3	4	4	2	2	-	-
Alabama												
All races, both sexes	1	2	-	1	5	2	3	2	1	-	-	-
Male	1	1	-	-	3	1	2	1	-	-	-	-
Female	-	1	-	1	2	1	1	1	1	-	-	-
White, both sexes	1	2	-	1	3	2	3	2	1	-	-	-
Male	1	1	-	-	2	1	2	1	-	-	-	-
Female	-	1	-	1	1	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	2	5	-	5	1	5	2	-	-
Male	-	-	-	1	2	-	1	-	1	2	-	-
Female	1	-	-	1	3	-	4	1	4	-	-	-
White, both sexes	1	-	-	1	5	-	4	1	5	2	-	-
Male	-	-	-	1	2	-	1	-	1	2	-	-
Female	1	-	-	-	3	-	3	1	4	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Arkansas												
All races, both sexes	2	1	1	-	1	1	6	2	1	1	-	-
Male	2	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	1	-	5	2	1	1	-	-
White, both sexes	-	1	1	-	-	1	6	1	1	1	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	5	1	1	1	-	-
Black, both sexes	2	-	-	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	5	12	14	19	28	40	31	38	13	1	-
Male	6	3	6	8	7	16	19	12	16	4	1	-
Female	1	2	6	6	12	12	21	19	22	9	-	-
White, both sexes	5	4	9	10	15	20	32	25	36	10	-	-
Male	5	3	6	4	5	12	16	11	15	3	-	-
Female	-	1	3	6	10	8	16	14	21	7	-	-
Black, both sexes	1	1	2	1	2	2	2	1	1	-	-	-
Male	1	-	-	1	1	2	-	-	-	-	-	-
Female	-	1	2	-	1	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	2	5	6	5	1	3	1	-
Male	-	-	-	3	1	2	3	1	1	1	1	-
Female	1	-	1	-	1	3	3	4	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Colorado												
All races, both sexes	1	1	-	-	2	3	3	2	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	2	3	2	2	-	-	-	-
White, both sexes	-	1	-	-	2	3	3	2	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	2	3	2	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	1	1	1	3	3	2	1	1	-	-
Male	-	-	1	-	-	2	2	-	-	1	-	-
Female	-	1	-	1	1	1	1	2	1	-	-	-
White, both sexes	-	1	1	1	1	3	3	2	1	1	-	-
Male	-	-	1	-	-	2	2	-	-	1	-	-
Female	-	1	-	1	1	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Delaware												
All races, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Florida												
All races, both sexes	4	3	2	6	10	8	18	9	6	2	-	-
Male	3	3	1	5	5	5	7	8	1	-	-	-
Female	1	-	1	1	5	3	11	1	5	2	-	-
White, both sexes	4	2	2	5	8	7	17	8	5	2	-	-
Male	3	2	1	4	4	5	7	7	1	-	-	-
Female	1	-	1	1	4	2	10	1	4	2	-	-
Black, both sexes	-	1	-	1	2	1	-	1	1	-	-	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	-	-	2	-	5	6	11	2	1	1	-	-
Male	-	-	2	-	4	4	6	1	-	-	-	-
Female	-	-	-	-	1	2	5	1	1	1	-	-
White, both sexes	-	-	2	-	3	5	9	2	1	1	-	-
Male	-	-	2	-	2	3	5	1	-	-	-	-
Female	-	-	-	-	1	2	4	1	1	1	-	-
Black, both sexes	-	-	-	-	2	1	2	-	-	-	-	-
Male	-	-	-	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Hawaii												
All races, both sexes	-	1	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	1	1	1	-	2	1	2	-	1	-	-
Male	-	1	-	1	-	1	-	2	-	1	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	1	1	1	1	-	2	1	2	-	1	-	-
Male	-	1	-	1	-	1	-	2	-	1	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Illinois												
All races, both sexes	2	3	4	2	2	6	7	10	10	3	1	-
Male	2	3	1	1	2	2	4	-	3	1	-	-
Female	-	-	3	1	-	4	3	10	7	2	1	-
White, both sexes	1	3	3	1	2	6	7	10	8	3	1	-
Male	1	3	1	-	2	2	4	-	3	1	-	-
Female	-	-	2	1	-	4	3	10	5	2	1	-
Black, both sexes	1	-	1	1	-	-	-	-	2	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	1	-	1	1	3	4	9	2	1	-	-
Male	-	1	-	1	-	-	2	3	2	-	-	-
Female	-	-	-	-	1	3	2	6	-	1	-	-
White, both sexes	-	1	-	1	1	3	4	8	2	1	-	-
Male	-	1	-	1	-	-	2	2	2	-	-	-
Female	-	-	-	-	1	3	2	6	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Iowa												
All races, both sexes	1	-	-	1	2	3	1	2	3	-	1	-
Male	1	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	1	1	2	1	1	2	-	1	-
White, both sexes	1	-	-	1	2	3	1	2	3	-	1	-
Male	1	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	1	1	2	1	1	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	1	3	2	1	1	-	-	-
Male	-	-	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	1	1	-	-	1	-	-	-
White, both sexes	-	-	-	1	1	3	2	1	1	-	-	-
Male	-	-	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Kentucky												
All races, both sexes	1	-	-	-	-	3	3	5	3	-	-	-
Male	1	-	-	-	-	3	1	1	2	-	-	-
Female	-	-	-	-	-	-	2	4	1	-	-	-
White, both sexes	1	-	-	-	-	2	3	5	3	-	-	-
Male	1	-	-	-	-	2	1	1	2	-	-	-
Female	-	-	-	-	-	-	2	4	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	3	1	2	1	5	1	4	-	-	-
Male	-	-	3	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	5	1	4	-	-	-
White, both sexes	-	-	2	1	2	1	2	1	2	-	-	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	2	1	2	-	-	-
Black, both sexes	-	-	1	-	-	-	3	-	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Maine												
All races, both sexes	-	1	1	-	1	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	1	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	1	1	1	2	4	-	-	-	-
Male	-	-	1	1	-	-	-	3	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
White, both sexes	-	-	-	1	1	1	2	3	-	-	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Massachusetts												
All races, both sexes	-	-	1	-	2	4	-	2	5	-	-	-
Male	-	-	1	-	-	2	-	1	3	-	-	-
Female	-	-	-	-	2	2	-	1	2	-	-	-
White, both sexes	-	-	1	-	1	4	-	2	5	-	-	-
Male	-	-	1	-	-	2	-	1	3	-	-	-
Female	-	-	-	-	1	2	-	1	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	2	1	8	5	7	7	2	-	-	-
Male	1	1	2	-	4	2	4	1	1	-	-	-
Female	-	-	-	1	4	3	3	6	1	-	-	-
White, both sexes	1	1	2	-	5	5	7	6	2	-	-	-
Male	1	1	2	-	2	2	4	1	1	-	-	-
Female	-	-	-	-	3	3	3	5	1	-	-	-
Black, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Minnesota												
All races, both sexes	-	-	-	1	-	-	3	2	2	-	-	-
Male	-	-	-	-	-	-	2	1	1	-	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	1	-	-	3	2	2	-	-	-
Male	-	-	-	-	-	-	2	1	1	-	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	1	-	3	4	-	2	1	-	-	-
Male	-	-	1	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	2	2	-	2	1	-	-	-
White, both sexes	1	-	-	-	3	3	-	1	1	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	2	1	-	1	1	-	-	-
Black, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Missouri												
All races, both sexes	-	1	2	1	1	1	8	4	4	-	-	-
Male	-	1	1	1	-	1	1	1	-	-	-	-
Female	-	-	1	-	1	-	7	3	4	-	-	-
White, both sexes	-	-	2	1	1	1	7	4	4	-	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	-	1	-	1	-	6	3	4	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	-	-	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Nebraska												
All races, both sexes	-	1	1	-	-	1	-	2	1	3	-	-
Male	-	1	-	-	-	-	-	2	-	1	-	-
Female	-	-	1	-	-	1	-	-	1	2	-	-
White, both sexes	-	1	1	-	-	1	-	2	1	3	-	-
Male	-	1	-	-	-	-	-	2	-	1	-	-
Female	-	-	1	-	-	1	-	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	1	2	-	1	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	2	-	-	-	1	2	-	1	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
New Hampshire												
All races, both sexes	-	-	-	1	1	-	1	2	-	-	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	2	-	-	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	3	-	-	4	3	8	5	2	2	-	-
Male	2	2	-	-	3	2	5	3	1	1	-	-
Female	1	1	-	-	1	1	3	2	1	1	-	-
White, both sexes	2	3	-	-	3	3	7	5	2	1	-	-
Male	1	2	-	-	2	2	4	3	1	-	-	-
Female	1	1	-	-	1	1	3	2	1	1	-	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
New Mexico												
All races, both sexes	-	-	-	1	1	-	1	1	3	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	3	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	3	2	5	3	14	15	13	9	2	-	-
Male	2	2	-	4	1	8	7	7	3	1	-	-
Female	3	1	2	1	2	6	8	6	6	1	-	-
White, both sexes	4	2	1	4	2	11	12	11	7	2	-	-
Male	1	1	-	3	-	6	6	6	3	1	-	-
Female	3	1	1	1	2	5	6	5	4	1	-	-
Black, both sexes	1	1	1	1	1	1	3	2	1	-	-	-
Male	1	1	-	1	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
North Carolina												
All races, both sexes	1	-	-	3	4	5	4	6	2	-	1	-
Male	-	-	-	1	3	2	1	-	1	-	-	-
Female	1	-	-	2	1	3	3	6	1	-	1	-
White, both sexes	-	-	-	3	3	4	4	5	1	-	1	-
Male	-	-	-	1	2	1	1	-	-	-	-	-
Female	-	-	-	2	1	3	3	5	1	-	1	-
Black, both sexes	1	-	-	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	1	5	1	-	-
Male	1	-	-	-	-	-	-	-	3	-	-	-
Female	-	-	-	-	-	-	-	1	2	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	5	1	-	-
Male	1	-	-	-	-	-	-	-	3	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Ohio												
All races, both sexes	3	2	2	3	6	3	11	13	5	4	-	-
Male	1	-	1	1	5	1	4	3	1	-	-	-
Female	2	2	1	2	1	2	7	10	4	4	-	-
White, both sexes	3	2	2	3	5	3	11	13	5	4	-	-
Male	1	-	1	1	4	1	4	3	1	-	-	-
Female	2	2	1	2	1	2	7	10	4	4	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	1	1	3	-	3	4	3	4	-	-
Male	-	1	1	1	2	-	1	2	1	2	-	-
Female	-	1	-	-	1	-	2	2	2	2	-	-
White, both sexes	-	1	-	1	2	-	3	4	2	4	-	-
Male	-	1	-	1	1	-	1	2	1	2	-	-
Female	-	-	-	-	1	-	2	2	1	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Oregon												
All races, both sexes	-	2	-	2	-	2	5	7	3	1	-	-
Male	-	2	-	2	-	2	2	3	1	-	-	-
Female	-	-	-	-	-	-	3	4	2	1	-	-
White, both sexes	-	2	-	2	-	2	5	7	3	1	-	-
Male	-	2	-	2	-	2	2	3	1	-	-	-
Female	-	-	-	-	-	-	3	4	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	3	4	5	1	7	12	8	2	-	-
Male	-	-	3	2	3	1	3	4	2	-	-	-
Female	1	-	-	2	2	-	4	8	6	2	-	-
White, both sexes	1	-	3	4	4	1	7	12	8	2	-	-
Male	-	-	3	2	3	1	3	4	2	-	-	-
Female	1	-	-	2	1	-	4	8	6	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Rhode Island												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	3	1	2	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	2	1	1	1	-	-	-	-	-
White, both sexes	-	-	1	1	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
South Dakota												
All races, both sexes	-	-	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	3	2	2	3	6	6	1	3	1	1	-
Male	1	-	2	2	1	4	1	-	-	-	-	-
Female	-	3	-	-	2	2	5	1	3	1	1	-
White, both sexes	1	3	2	1	3	5	6	1	2	1	1	-
Male	1	-	2	1	1	3	1	-	-	-	-	-
Female	-	3	-	-	2	2	5	1	2	1	1	-
Black, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Texas												
All races, both sexes	1	3	1	6	9	7	13	12	8	1	1	-
Male	1	2	-	4	3	1	9	3	5	-	-	-
Female	-	1	1	2	6	6	4	9	3	1	1	-
White, both sexes	1	2	1	6	9	6	11	12	7	1	1	-
Male	1	1	-	4	3	1	7	3	4	-	-	-
Female	-	1	1	2	6	5	4	9	3	1	1	-
Black, both sexes	-	1	-	-	-	1	2	-	1	-	-	-
Male	-	1	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	1	-	-	1	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	1	-	1	-	-	1	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Vermont												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	1	-	2	1	1	4	2	2	-	-
Male	-	1	-	-	1	1	-	3	1	1	-	-
Female	-	-	1	-	1	-	1	1	1	1	-	-
White, both sexes	-	1	-	-	1	1	1	3	2	1	-	-
Male	-	1	-	-	1	1	-	2	1	1	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	1	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Washington												
All races, both sexes	1	-	1	1	-	1	5	1	6	1	-	-
Male	1	-	1	-	-	1	4	1	2	-	-	-
Female	-	-	-	1	-	-	1	-	4	1	-	-
White, both sexes	-	-	1	1	-	1	5	1	6	1	-	-
Male	-	-	1	-	-	1	4	1	2	-	-	-
Female	-	-	-	1	-	-	1	-	4	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	-	2	2	-	1	-	-	-
Male	-	-	-	1	-	2	2	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	2	2	-	1	-	-	-
Male	-	-	-	1	-	2	2	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Wisconsin												
All races, both sexes	3	-	2	-	1	5	4	4	3	2	1	-
Male	-	-	1	-	1	4	1	-	-	-	-	-
Female	3	-	1	-	-	1	3	4	3	2	1	-
White, both sexes	3	-	2	-	1	5	4	4	3	2	1	-
Male	-	-	1	-	1	4	1	-	-	-	-	-
Female	3	-	1	-	-	1	3	4	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	2	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	2	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
United States												
All races, both sexes	27	-	-	-	-	-	-	1	-	1	-	3
Male	15	-	-	-	-	-	-	1	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	25	-	-	-	-	-	-	1	-	1	-	2
Male	15	-	-	-	-	-	-	1	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Florida												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
North Carolina												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastrojejunal ulcer (K28)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
United States												
All races, both sexes	-	1	2	3	4	2	4	3	3	-	-	-
Male	-	-	2	2	3	-	3	1	1	-	-	-
Female	-	1	-	1	1	2	1	2	2	-	-	-
White, both sexes	-	1	2	3	4	2	4	2	3	-	-	-
Male	-	-	2	2	3	-	3	1	1	-	-	-
Female	-	1	-	1	1	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Florida												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Kentucky												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Minnesota												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Missouri												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
North Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Ohio												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Washington												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
United States												
All races, both sexes	333	-	1	1	-	3	1	2	2	5	10	20
Male	169	-	1	1	-	3	1	1	2	5	6	15
Female	164	-	-	-	-	-	-	1	-	-	4	5
White, both sexes	290	-	1	-	-	3	1	1	1	3	6	17
Male	145	-	1	-	-	3	1	1	1	3	2	13
Female	145	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	36	-	-	1	-	-	-	1	-	2	4	3
Male	19	-	-	1	-	-	-	-	-	2	4	2
Female	17	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Arkansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	52	-	1	-	-	1	-	-	-	-	3	2
Male	29	-	1	-	-	1	-	-	-	-	2	2
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	44	-	1	-	-	1	-	-	-	-	1	2
Male	22	-	1	-	-	1	-	-	-	-	-	2
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Colorado												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Florida												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Illinois												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Iowa												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Kentucky												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Maine												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Massachusetts												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	18	-	-	-	-	-	1	-	-	1	-	4
Male	13	-	-	-	-	-	1	-	-	1	-	3
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	1	-	-	1	-	3
Male	13	-	-	-	-	-	1	-	-	1	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Minnesota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	-	-	1	-	-	-	-	1	-	-	-
Male	3	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Missouri												
All races, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Nebraska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
New Mexico												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
North Carolina												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Ohio												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Oregon												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10	-	-	-	-	1	-	-	-	-	-	1
Male	7	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	1	-	-	-	-	-	1
Male	7	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	-	-	-	-	-	-	1	-	1	-	1
Male	7	-	-	-	-	-	-	1	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Texas												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Washington												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
Wisconsin												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
United States												
All races, both sexes	12	21	16	16	24	38	50	62	35	11	3	-
Male	9	17	9	11	12	25	22	21	7	1	-	-
Female	3	4	7	5	12	13	28	41	28	10	3	-
White, both sexes	12	20	10	14	20	35	44	56	34	9	3	-
Male	9	16	5	10	11	22	19	20	7	1	-	-
Female	3	4	5	4	9	13	25	36	27	8	3	-
Black, both sexes	-	1	6	2	4	2	3	5	1	1	-	-
Male	-	1	4	1	1	2	1	-	-	-	-	-
Female	-	-	2	1	3	-	2	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Arkansas												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	1	3	3	1	10	8	8	7	2	-	-
Male	-	1	3	2	1	7	4	4	1	-	-	-
Female	2	-	-	1	-	3	4	4	6	2	-	-
White, both sexes	2	1	2	3	-	9	7	7	7	1	-	-
Male	-	1	2	2	-	6	3	3	1	-	-	-
Female	2	-	-	1	-	3	4	4	6	1	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Colorado												
All races, both sexes	-	3	-	-	2	3	-	-	1	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	1	2	-	-	1	-	-	-
White, both sexes	-	3	-	-	2	3	-	-	1	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	1	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Florida												
All races, both sexes	1	2	1	-	-	3	5	2	3	-	-	-
Male	1	1	-	-	-	2	2	1	-	-	-	-
Female	-	1	1	-	-	1	3	1	3	-	-	-
White, both sexes	1	2	-	-	-	3	4	2	3	-	-	-
Male	1	1	-	-	-	2	1	1	-	-	-	-
Female	-	1	-	-	-	1	3	1	3	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	-	-	-	-	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Illinois												
All races, both sexes	1	-	2	-	3	1	1	5	1	-	-	-
Male	1	-	1	-	2	1	-	2	-	-	-	-
Female	-	-	1	-	1	-	1	3	1	-	-	-
White, both sexes	1	-	2	-	2	1	1	5	1	-	-	-
Male	1	-	1	-	2	1	-	2	-	-	-	-
Female	-	-	1	-	-	-	1	3	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Iowa												
All races, both sexes	2	-	1	-	-	1	1	3	3	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	3	2	-	-	-
White, both sexes	2	-	1	-	-	1	1	3	3	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	1	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Kentucky												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	1	-	1	-	-	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Maine												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	1	-	1	4	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	3	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	-	-	-	5	4	-	1	-	-	-
Male	1	-	-	-	-	3	3	-	1	-	-	-
Female	-	1	-	-	-	2	1	-	-	-	-	-
White, both sexes	1	1	-	-	-	5	4	-	1	-	-	-
Male	1	-	-	-	-	3	3	-	1	-	-	-
Female	-	1	-	-	-	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Minnesota												
All races, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Missouri												
All races, both sexes	-	-	-	1	1	1	1	2	-	-	-	-
Male	-	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	1	1	1	2	-	-	-	-
Male	-	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	-	2	-	4	4	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	3	4	1	-	-	-
White, both sexes	-	-	1	-	2	-	2	2	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
New Mexico												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	2	2	2	1	2	5	-	3	1	-	-
Male	-	2	1	1	-	1	1	-	1	-	-	-
Female	1	-	1	1	1	1	4	-	2	1	-	-
White, both sexes	1	2	1	2	1	2	5	-	2	1	-	-
Male	-	2	-	1	-	1	1	-	1	-	-	-
Female	1	-	1	1	1	1	4	-	1	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
North Carolina												
All races, both sexes	1	3	1	-	-	1	1	-	2	1	-	-
Male	1	2	1	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	2	1	-	-
White, both sexes	1	2	1	-	-	1	1	-	2	-	-	-
Male	1	1	1	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Ohio												
All races, both sexes	1	-	-	-	2	1	-	4	-	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
White, both sexes	1	-	-	-	2	1	-	4	-	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Oregon												
All races, both sexes	-	-	-	1	-	2	2	3	-	1	-	-
Male	-	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	1	-	-
White, both sexes	-	-	-	1	-	2	2	3	-	1	-	-
Male	-	-	-	-	-	2	1	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	1	1	-	3	2	-	-	-	-
Male	-	1	-	1	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	1	1	-	2	2	-	-	-	-
Male	-	1	-	1	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Rhode Island												
All races, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	1	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	1	-	2	-	1	1	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Texas												
All races, both sexes	-	4	-	1	1	1	2	2	2	1	-	-
Male	-	4	-	1	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	-	1	1	1	-	-
White, both sexes	-	4	-	1	1	1	2	2	2	1	-	-
Male	-	4	-	1	-	1	2	1	1	-	-	-
Female	-	-	-	-	1	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Wisconsin												
All races, both sexes	-	-	-	-	1	1	1	2	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
United States												
All races, both sexes	554	2	1	1	2	2	-	4	7	7	10	20
Male	231	1	1	1	1	1	-	1	3	2	7	7
Female	323	1	-	-	1	1	-	3	4	5	3	13
White, both sexes	473	1	1	-	1	2	-	2	6	5	8	13
Male	188	1	1	-	-	1	-	-	2	1	6	2
Female	285	-	-	-	1	1	-	2	4	4	2	11
Black, both sexes	63	1	-	1	1	-	-	2	1	1	1	6
Male	33	-	-	1	1	-	-	1	1	-	1	4
Female	30	1	-	-	-	-	-	1	-	1	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Arkansas												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	33	-	-	-	-	-	-	-	1	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	24	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Colorado												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Florida												
All races, both sexes	33	-	-	-	-	-	-	1	1	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	30	-	-	-	-	-	-	1	1	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Illinois												
All races, both sexes	32	-	-	1	-	-	-	-	-	-	1	2
Male	8	-	-	1	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	1	-	-	-	-	-	-	-	1
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Iowa												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Kentucky												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Maine												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	1	-	-	-	-	-	-
Male	5	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Massachusetts												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Minnesota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Missouri												
All races, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Nebraska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	-	-	-	-	1	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	12	-	-	-	-	1	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
New Mexico												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	26	-	-	-	1	-	-	-	-	1	2	1
Male	10	-	-	-	1	-	-	-	-	1	2	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	2	1
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
North Carolina												
All races, both sexes	22	-	-	-	-	-	-	-	1	1	1	1
Male	15	-	-	-	-	-	-	-	1	-	1	1
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	1	-	-	-	-	-	-	1	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	1	-	-	-	-	-	-	1	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Ohio												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Oregon												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	33	-	-	-	-	-	-	-	-	1	1	2
Male	15	-	-	-	-	-	-	-	-	1	-	2
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	28	-	-	-	-	-	-	-	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	-	-	-	1	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	1	-	-	-	-	1	-	2
White, both sexes	17	-	-	-	1	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	1	-	-	-	-	1	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Texas												
All races, both sexes	31	-	-	-	-	-	-	1	-	-	2	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	1	-	-	-	-	-	1	2	-	-	-
Male	4	-	-	-	-	-	-	1	2	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	2	-	-	-
Male	3	-	-	-	-	-	-	-	2	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Washington												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Wisconsin												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
United States												
All races, both sexes	32	39	41	39	46	63	72	93	47	24	2	-
Male	15	23	19	14	22	25	36	27	16	8	1	-
Female	17	16	22	25	24	38	36	66	31	16	1	-
White, both sexes	26	34	34	33	39	54	64	86	40	23	1	-
Male	12	21	15	12	17	20	30	26	14	7	-	-
Female	14	13	19	21	22	34	34	60	26	16	1	-
Black, both sexes	5	5	6	6	5	6	7	4	5	-	-	-
Male	2	2	4	2	4	4	5	-	1	-	-	-
Female	3	3	2	4	1	2	2	4	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	2	1	3	2	1	1	-
Male	1	-	-	-	1	1	1	1	1	1	1	-
Female	-	-	1	-	1	1	-	2	1	-	-	-
Alabama												
All races, both sexes	-	-	1	2	3	-	-	1	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	2	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	1	-	-	-	-
Black, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Alaska												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	1	-	-	-	2	1	2	1	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	1	-	-	-	2	1	2	1	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Arkansas												
All races, both sexes	1	-	2	-	1	1	1	2	-	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	1	1	1	-	-	-	-
White, both sexes	1	-	2	-	1	1	1	2	-	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	2	2	5	3	6	2	4	1	1	-
Male	2	-	2	-	3	1	4	-	1	1	1	-
Female	2	-	-	2	2	2	2	2	3	-	-	-
White, both sexes	4	-	2	1	3	2	5	1	3	1	-	-
Male	2	-	2	-	2	1	3	-	1	1	-	-
Female	2	-	-	1	1	1	2	1	2	-	-	-
Black, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Colorado												
All races, both sexes	1	1	-	1	2	4	1	2	1	-	-	-
Male	-	1	-	-	1	2	1	1	-	-	-	-
Female	1	-	-	1	1	2	-	1	1	-	-	-
White, both sexes	1	1	-	1	2	4	1	2	1	-	-	-
Male	-	1	-	-	1	2	1	1	-	-	-	-
Female	1	-	-	1	1	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Florida												
All races, both sexes	1	6	2	1	2	5	3	3	4	3	-	-
Male	1	5	1	-	1	2	3	2	2	1	-	-
Female	-	1	1	1	1	3	-	1	2	2	-	-
White, both sexes	1	5	2	1	2	4	3	3	3	3	-	-
Male	1	5	1	-	1	1	3	2	2	1	-	-
Female	-	-	1	1	1	3	-	1	1	2	-	-
Black, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	2	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	2	-	-	1	2	1	-	-	-
White, both sexes	1	-	-	2	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	2	-	-	1	2	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Hawaii												
All races, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	1	2	-	2	-	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
White, both sexes	1	-	-	1	2	-	2	-	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Illinois												
All races, both sexes	1	2	2	-	5	4	4	7	2	1	-	-
Male	-	1	2	-	-	2	-	-	1	-	-	-
Female	1	1	-	-	5	2	4	7	1	1	-	-
White, both sexes	1	2	1	-	5	3	4	7	1	1	-	-
Male	-	1	1	-	-	1	-	-	1	-	-	-
Female	1	1	-	-	5	2	4	7	-	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	2	4	-	-	3	1	2	1	-	-	-
Male	-	1	2	-	-	1	-	-	-	-	-	-
Female	-	1	2	-	-	2	1	2	1	-	-	-
White, both sexes	-	2	4	-	-	3	1	2	1	-	-	-
Male	-	1	2	-	-	1	-	-	-	-	-	-
Female	-	1	2	-	-	2	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Iowa												
All races, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	-	1	-	2	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	-	1	-	1	-	2	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Kentucky												
All races, both sexes	1	-	-	1	1	2	3	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	2	3	-	-	-	-	-
White, both sexes	1	-	-	1	1	2	3	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	2	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	-	2	-	2	2	1	2	-	-	-
Male	-	1	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	2	-	-	-
White, both sexes	-	1	-	1	-	1	2	1	2	-	-	-
Male	-	1	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Maine												
All races, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	1	1	-	1	1	3	-	-	-	-
Male	-	1	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	1	1	-	-	1	1	3	-	-	-	-
Male	-	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Massachusetts												
All races, both sexes	-	-	1	-	-	1	3	6	2	1	-	-
Male	-	-	-	-	-	1	2	4	-	-	-	-
Female	-	-	1	-	-	-	1	2	2	1	-	-
White, both sexes	-	-	1	-	-	1	3	6	2	1	-	-
Male	-	-	-	-	-	1	2	4	-	-	-	-
Female	-	-	1	-	-	-	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	3	1	2	2	4	3	2	2	-	-	-
Male	1	2	-	-	1	-	2	-	-	-	-	-
Female	1	1	1	2	1	4	1	2	2	-	-	-
White, both sexes	2	3	1	2	2	4	2	2	1	-	-	-
Male	1	2	-	-	1	-	1	-	-	-	-	-
Female	1	1	1	2	1	4	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Minnesota												
All races, both sexes	-	-	2	1	-	1	1	2	2	-	-	-
Male	-	-	1	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	2	1	-	-	-
White, both sexes	-	-	1	1	-	-	1	2	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Missouri												
All races, both sexes	-	-	-	2	-	2	3	1	1	-	1	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	2	-	1	1	1	1	-	1	-
White, both sexes	-	-	-	2	-	2	2	1	1	-	1	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	2	-	1	1	1	1	-	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	-	-	-	-	1	1	-	-	-
Male	1	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	1	1	-	-	-
Male	1	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	2	4	2	1	-	-	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	-	1	-	-	3	2	-	-	-	-	-	-
White, both sexes	-	1	-	2	4	2	1	-	-	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	-	1	-	-	3	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	1	2	2	2	-	4	1	1	-	-
Male	-	-	-	2	2	-	-	1	-	-	-	-
Female	1	-	1	-	-	2	-	3	1	1	-	-
White, both sexes	1	-	1	2	1	2	-	4	-	1	-	-
Male	-	-	-	2	1	-	-	1	-	-	-	-
Female	1	-	1	-	-	2	-	3	-	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
New Mexico												
All races, both sexes	-	1	-	-	-	1	2	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	2	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	1	1	-	-	4	2	9	3	-	-	-
Male	-	-	-	-	-	1	1	2	2	-	-	-
Female	1	1	1	-	-	3	1	7	1	-	-	-
White, both sexes	1	1	-	-	-	3	1	8	3	-	-	-
Male	-	-	-	-	-	1	1	2	2	-	-	-
Female	1	1	-	-	-	2	-	6	1	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
North Carolina												
All races, both sexes	1	1	4	1	1	1	1	3	2	3	-	-
Male	1	1	2	-	1	-	1	3	2	1	-	-
Female	-	-	2	1	-	1	-	-	-	2	-	-
White, both sexes	-	1	4	-	-	1	-	3	2	3	-	-
Male	-	1	2	-	-	-	-	3	2	1	-	-
Female	-	-	2	-	-	1	-	-	-	2	-	-
Black, both sexes	1	-	-	1	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Ohio												
All races, both sexes	3	3	3	2	-	-	2	10	3	1	-	-
Male	2	1	1	-	-	-	2	2	1	-	-	-
Female	1	2	2	2	-	-	-	8	2	1	-	-
White, both sexes	3	3	3	2	-	-	2	10	2	1	-	-
Male	2	1	1	-	-	-	2	2	-	-	-	-
Female	1	2	2	2	-	-	-	8	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	1	-	3	-	1	-	1	-	-	-
Male	-	2	-	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	-	-
White, both sexes	-	2	1	-	3	-	1	-	1	-	-	-
Male	-	2	-	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Oregon												
All races, both sexes	-	-	3	-	1	-	2	2	1	-	-	-
Male	-	-	2	-	-	-	1	2	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	-	-
White, both sexes	-	-	3	-	1	-	2	1	1	-	-	-
Male	-	-	2	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	-	2	2	4	6	6	3	4	-	-
Male	1	-	-	1	1	2	3	2	1	1	-	-
Female	-	1	-	1	1	2	3	4	2	3	-	-
White, both sexes	-	1	-	1	2	3	6	6	3	4	-	-
Male	-	-	-	1	1	1	3	2	1	1	-	-
Female	-	1	-	-	1	2	3	4	2	3	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	1	2	3	2	1	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	1	1	1	1	2	2	2	1	-	-	-	-
White, both sexes	-	-	1	1	1	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	1	1	2	-	-	-	-	-
Black, both sexes	1	1	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	2	1	4	1	1	2	2	-	-	-	-
Male	1	2	-	2	1	-	1	-	-	-	-	-
Female	1	-	1	2	-	1	1	2	-	-	-	-
White, both sexes	1	2	1	4	1	1	2	1	-	-	-	-
Male	-	2	-	2	1	-	1	-	-	-	-	-
Female	1	-	1	2	-	1	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Texas												
All races, both sexes	5	5	1	4	1	3	1	4	1	3	-	-
Male	2	3	1	1	1	2	1	1	-	2	-	-
Female	3	2	-	3	-	1	-	3	1	1	-	-
White, both sexes	3	3	-	4	1	2	1	3	1	3	-	-
Male	2	1	-	1	1	1	1	1	-	2	-	-
Female	1	2	-	3	-	1	-	2	1	1	-	-
Black, both sexes	2	2	1	-	-	-	-	1	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Washington												
All races, both sexes	1	1	1	-	1	-	3	1	3	2	-	-
Male	-	1	-	-	1	-	1	-	-	1	-	-
Female	1	-	1	-	-	-	2	1	3	1	-	-
White, both sexes	1	1	1	-	1	-	3	-	3	2	-	-
Male	-	1	-	-	1	-	1	-	-	1	-	-
Female	1	-	1	-	-	-	2	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Wisconsin												
All races, both sexes	-	-	-	1	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
United States												
All races, both sexes	439	2	4	5	7	3	6	6	3	6	13	12
Male	247	-	2	5	4	3	4	2	2	5	10	5
Female	192	2	2	-	3	-	2	4	1	1	3	7
White, both sexes	373	2	4	5	2	1	3	5	3	4	10	10
Male	208	-	2	5	1	1	2	2	2	3	7	4
Female	165	2	2	-	1	-	1	3	1	1	3	6
Black, both sexes	55	-	-	-	4	2	3	1	-	2	3	2
Male	33	-	-	-	3	2	2	-	-	2	3	1
Female	22	-	-	-	1	-	1	1	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	1	-	-	-	-	-	-	-
Alabama												
All races, both sexes	5	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Arkansas												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	53	1	2	-	2	-	1	1	1	-	-	-
Male	32	-	1	-	2	-	1	1	1	-	-	-
Female	21	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	39	1	2	-	1	-	-	1	1	-	-	-
Male	25	-	1	-	1	-	-	1	1	-	-	-
Female	14	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	1	-	1	-	-	-	-	-
Male	5	-	-	-	1	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Colorado												
All races, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	9	-	-	-	-	1	-	-	1	-	-	-
Male	5	-	-	-	-	1	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Florida												
All races, both sexes	22	-	-	1	-	-	-	-	-	-	1	1
Male	12	-	-	1	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	20	-	-	1	-	-	-	-	-	-	1	-
Male	11	-	-	1	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Illinois												
All races, both sexes	18	-	-	1	-	-	1	1	-	-	-	1
Male	12	-	-	1	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	18	-	-	1	-	-	1	1	-	-	-	1
Male	12	-	-	1	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Iowa												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Kentucky												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Maine												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Massachusetts												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	18	-	1	-	-	-	1	-	-	-	2	-
Male	8	-	1	-	-	-	-	-	-	-	2	-
Female	10	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	16	-	1	-	-	-	-	-	-	-	2	-
Male	8	-	1	-	-	-	-	-	-	-	2	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Minnesota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Missouri												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Nebraska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	-	-	1	1	-	-	-	-	-	1	-
Male	5	-	-	1	-	-	-	-	-	-	1	-
Female	6	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	8	-	-	1	-	-	-	-	-	-	-	-
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
New Mexico												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
North Carolina												
All races, both sexes	14	-	1	-	1	-	2	-	-	-	1	-
Male	7	-	-	-	1	-	2	-	-	-	1	-
Female	7	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	1	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	1	-	1	-	-	-	1	-
Male	3	-	-	-	1	-	1	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Ohio												
All races, both sexes	17	-	-	-	1	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	1	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Oregon												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15	-	-	1	1	-	-	-	1	-	-	-
Male	8	-	-	1	1	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	12	-	-	1	-	-	-	-	1	-	-	-
Male	6	-	-	1	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Texas												
All races, both sexes	34	1	-	-	-	1	-	-	-	3	1	1
Male	22	-	-	-	-	1	-	-	-	3	-	-
Female	12	1	-	-	-	-	-	-	-	-	1	1
White, both sexes	29	1	-	-	-	-	-	-	-	1	1	1
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	12	1	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	1	-	-	-	2	-	-
Male	5	-	-	-	-	1	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	13	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Washington												
All races, both sexes	10	-	-	1	-	-	-	-	-	-	1	1
Male	5	-	-	1	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	1	-	-	-	-	-	-	1	1
Male	5	-	-	1	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Wisconsin												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
United States												
All races, both sexes	21	27	30	39	37	58	52	63	33	10	2	-
Male	13	22	15	27	26	29	28	28	13	3	1	-
Female	8	5	15	12	11	29	24	35	20	7	1	-
White, both sexes	13	23	23	34	31	48	50	61	31	8	2	-
Male	8	18	13	24	22	24	28	27	11	3	1	-
Female	5	5	10	10	9	24	22	34	20	5	1	-
Black, both sexes	7	4	5	4	5	8	1	2	-	2	-	-
Male	4	4	1	3	3	4	-	1	-	-	-	-
Female	3	-	4	1	2	4	1	1	-	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	1	1	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
Alabama												
All races, both sexes	-	-	1	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	2	-	-	-	3	3	1	-	-	-
Male	-	1	2	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	1	1	-	-	-	3	3	1	-	-	-
Male	-	1	1	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Arkansas												
All races, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	4	5	4	4	8	5	5	5	2	-	-
Male	1	4	3	4	3	5	2	2	2	-	-	-
Female	2	-	2	-	1	3	3	3	3	2	-	-
White, both sexes	2	3	4	3	3	6	3	5	4	-	-	-
Male	-	3	3	3	2	5	2	2	1	-	-	-
Female	2	-	1	-	1	1	1	3	3	-	-	-
Black, both sexes	1	1	1	1	-	1	1	-	-	2	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Colorado												
All races, both sexes	-	-	1	-	-	2	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	3	-	2	2	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	3	-	2	2	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Florida												
All races, both sexes	3	1	1	-	1	4	4	4	1	-	-	-
Male	2	-	1	-	-	2	4	1	1	-	-	-
Female	1	1	-	-	1	2	-	3	-	-	-	-
White, both sexes	2	1	1	-	1	4	4	4	1	-	-	-
Male	1	-	1	-	-	2	4	1	1	-	-	-
Female	1	1	-	-	1	2	-	3	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	1	-	2	1	2	2	3	-	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	3	-	-	-
White, both sexes	-	1	1	-	2	1	2	2	3	-	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Illinois												
All races, both sexes	-	1	-	1	2	4	1	5	-	-	-	-
Male	-	1	-	1	2	1	1	4	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	-	-
White, both sexes	-	1	-	1	2	4	1	5	-	-	-	-
Male	-	1	-	1	2	1	1	4	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	2	-	1	2	1	3	2	-	-	-	-
Male	-	2	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
White, both sexes	-	1	-	1	2	1	3	1	-	-	-	-
Male	-	1	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Iowa												
All races, both sexes	-	-	-	1	-	1	1	-	2	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	1	-	1	1	-	2	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Kentucky												
All races, both sexes	1	-	-	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	3	1	1	1	-	-	-	-	-	-
Male	1	-	3	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	3	-	1	1	-	-	-	-	-	-
Male	-	-	3	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Maine												
All races, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	-	-	-	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Massachusetts												
All races, both sexes	-	-	1	1	-	1	4	-	-	-	1	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	1	-
White, both sexes	-	-	1	1	-	1	4	-	-	-	1	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	3	2	2	2	2	-	2	-	-	-
Male	-	1	1	1	1	-	-	-	1	-	-	-
Female	-	-	2	1	1	2	2	-	1	-	-	-
White, both sexes	-	1	3	1	2	2	2	-	2	-	-	-
Male	-	1	1	1	1	-	-	-	1	-	-	-
Female	-	-	2	-	1	2	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Minnesota												
All races, both sexes	-	-	-	1	1	-	1	-	2	1	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	1	-	1	-	2	1	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Missouri												
All races, both sexes	-	1	-	-	1	1	1	2	2	-	-	-
Male	-	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
White, both sexes	-	1	-	-	-	1	1	2	2	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Nebraska												
All races, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
New Hampshire												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	3	2	1	1	-	-	1	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	1	-	-	1	-	-
White, both sexes	-	-	-	2	2	1	1	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	1	-	-	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
New Mexico												
All races, both sexes	-	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	3	3	4	3	4	1	4	4	-	-	-
Male	1	3	-	2	-	1	-	2	1	-	-	-
Female	1	-	3	2	3	3	1	2	3	-	-	-
White, both sexes	1	3	3	3	2	3	1	4	4	-	-	-
Male	1	3	-	2	-	1	-	2	1	-	-	-
Female	-	-	3	1	2	2	1	2	3	-	-	-
Black, both sexes	1	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
North Carolina												
All races, both sexes	2	1	1	1	1	-	2	1	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	2	1	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	1	1	1	-	-	2	1	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Ohio												
All races, both sexes	-	-	1	2	-	4	2	4	-	-	-	-
Male	-	-	1	2	-	1	2	1	-	-	-	-
Female	-	-	-	-	-	3	-	3	-	-	-	-
White, both sexes	-	-	1	2	-	3	2	3	-	-	-	-
Male	-	-	1	2	-	1	2	1	-	-	-	-
Female	-	-	-	-	-	2	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Oregon												
All races, both sexes	-	-	-	1	-	-	2	2	-	2	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	-	-	-	1	-	-	2	2	-	2	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	1	1	2	1	1	-	3	-	-	1	-
Male	1	1	-	1	-	1	-	1	-	-	1	-
Female	1	-	1	1	1	-	-	2	-	-	-	-
White, both sexes	2	-	-	2	1	1	-	3	-	-	1	-
Male	1	-	-	1	-	1	-	1	-	-	1	-
Female	1	-	-	1	1	-	-	2	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Rhode Island												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	1	1	1	1	-	-	-	-	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
South Dakota												
All races, both sexes	1	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	2	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	2	-	-	2	-	3	-	1	-	-
Male	-	-	1	-	-	2	-	2	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	2	-	3	-	1	-	-
Male	-	-	-	-	-	2	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Texas												
All races, both sexes	2	2	1	5	6	4	3	4	-	-	-	-
Male	2	1	-	2	6	3	2	2	-	-	-	-
Female	-	1	1	3	-	1	1	2	-	-	-	-
White, both sexes	2	2	1	5	6	2	3	4	-	-	-	-
Male	2	1	-	2	6	1	2	2	-	-	-	-
Female	-	1	1	3	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	1	-	-	2	2	2	3	1	1	-	-
Male	-	-	-	-	1	1	1	1	-	1	-	-
Female	-	1	-	-	1	1	1	2	1	-	-	-
White, both sexes	-	1	-	-	2	1	2	3	1	1	-	-
Male	-	-	-	-	1	-	1	1	-	1	-	-
Female	-	1	-	-	1	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Washington												
All races, both sexes	-	-	1	1	-	-	2	3	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	2	2	-	-	-	-
White, both sexes	-	-	1	1	-	-	2	3	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	1	-	-	2	-	-	-	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	2	-	-	-	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Wisconsin												
All races, both sexes	-	1	-	-	-	2	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
White, both sexes	-	1	-	-	-	2	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
United States												
All races, both sexes	1,639	43	4	2	-	-	2	2	3	15	24	31
Male	646	28	1	1	-	-	-	-	1	6	9	16
Female	993	15	3	1	-	-	2	2	2	9	15	15
White, both sexes	1,456	35	4	1	-	-	1	1	3	11	18	21
Male	568	22	1	-	-	-	-	-	1	5	7	12
Female	888	13	3	1	-	-	1	1	2	6	11	9
Black, both sexes	152	5	-	1	-	-	1	1	-	3	5	8
Male	66	4	-	1	-	-	-	-	-	1	2	4
Female	86	1	-	-	-	-	1	1	-	2	3	4
American Indian 1/, both sexes	17	2	-	-	-	-	-	-	-	1	1	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	12	1	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	-	-	-	-	-	2
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
Alabama												
All races, both sexes	30	2	-	-	-	-	-	-	-	-	-	2
Male	13	2	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	2	-	-	-	-	-	-	-	-	-	1
Male	10	2	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	31	5	-	-	-	-	-	-	-	1	1	-
Male	17	2	-	-	-	-	-	-	-	1	-	-
Female	14	3	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	5	-	-	-	-	-	-	-	1	-	-
Male	16	2	-	-	-	-	-	-	-	1	-	-
Female	13	3	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Arkansas												
All races, both sexes	20	2	-	-	-	-	-	-	-	3	-	-
Male	7	2	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	17	2	-	-	-	-	-	-	-	3	-	-
Male	5	2	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	189	4	-	-	-	-	-	1	-	4	5	5
Male	80	1	-	-	-	-	-	-	-	3	3	2
Female	109	3	-	-	-	-	-	1	-	1	2	3
White, both sexes	166	4	-	-	-	-	-	-	-	3	5	3
Male	74	1	-	-	-	-	-	-	-	2	3	2
Female	92	3	-	-	-	-	-	-	-	1	2	1
Black, both sexes	14	-	-	-	-	-	-	1	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Colorado												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	21	1	-	1	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	20	1	-	1	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Florida												
All races, both sexes	78	2	-	-	-	-	1	-	-	2	-	-
Male	37	2	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	1	-	-	2	-	-
White, both sexes	70	2	-	-	-	-	1	-	-	1	-	-
Male	34	2	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	36	1	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	23	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Hawaii												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Illinois												
All races, both sexes	54	1	-	-	-	-	-	-	1	-	-	2
Male	29	1	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	49	1	-	-	-	-	-	-	1	-	-	1
Male	26	1	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	34	4	-	-	-	-	-	-	-	-	-	1
Male	13	2	-	-	-	-	-	-	-	-	-	1
Female	21	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	3	-	-	-	-	-	-	-	-	-	1
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	21	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Iowa												
All races, both sexes	22	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Kentucky												
All races, both sexes	26	-	-	-	-	-	-	1	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	26	-	-	-	-	-	-	1	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Maine												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	29	-	-	1	-	-	-	-	-	-	1	-
Male	11	-	-	1	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Massachusetts												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	62	1	-	-	-	-	-	-	-	-	1	3
Male	26	-	-	-	-	-	-	-	-	-	-	2
Female	36	1	-	-	-	-	-	-	-	-	1	1
White, both sexes	58	1	-	-	-	-	-	-	-	-	1	3
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	35	1	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Minnesota												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Missouri												
All races, both sexes	44	2	-	-	-	-	1	-	-	-	1	1
Male	20	2	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	41	2	-	-	-	-	-	-	-	-	1	1
Male	19	2	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
New Hampshire												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
New Mexico												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	105	2	-	-	-	-	-	-	-	-	1	-
Male	50	2	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	83	2	-	-	-	-	-	-	-	-	1	-
Male	40	2	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
North Carolina												
All races, both sexes	39	1	-	-	-	-	-	-	-	1	-	-
Male	18	1	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	-	-	1	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Ohio												
All races, both sexes	81	2	-	-	-	-	-	-	-	1	2	-
Male	29	1	-	-	-	-	-	-	-	-	-	-
Female	52	1	-	-	-	-	-	-	-	1	2	-
White, both sexes	78	2	-	-	-	-	-	-	-	1	1	-
Male	27	1	-	-	-	-	-	-	-	-	-	-
Female	51	1	-	-	-	-	-	-	-	1	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Oregon												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	94	2	-	-	-	-	-	-	-	-	1	-
Male	31	2	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	2	-	-	-	-	-	-	-	-	1	-
Male	30	2	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Rhode Island												
All races, both sexes	9	-	1	-	-	-	-	-	-	-	-	1
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	1	-	-	-	-	-	-	-	-	1
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	27	2	-	-	-	-	-	-	-	-	-	2
Male	11	2	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	2	-	-	-	-	-	-	-	-	-	1
Male	4	2	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	44	-	1	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	1	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Texas												
All races, both sexes	104	3	1	-	-	-	-	-	-	-	2	2
Male	40	1	-	-	-	-	-	-	-	-	-	-
Female	64	2	1	-	-	-	-	-	-	-	2	2
White, both sexes	95	3	1	-	-	-	-	-	-	-	1	2
Male	37	1	-	-	-	-	-	-	-	-	-	-
Female	58	2	1	-	-	-	-	-	-	-	1	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	1	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Vermont												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Washington												
All races, both sexes	37	4	-	-	-	-	-	-	-	-	-	1
Male	14	3	-	-	-	-	-	-	-	-	-	-
Female	23	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	2	-	-	-	-	-	-	-	-	-	-
Male	12	2	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
Wisconsin												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
United States												
All races, both sexes	52	69	112	95	115	182	272	296	246	65	9	-
Male	29	35	56	35	51	76	99	114	71	14	4	-
Female	23	34	56	60	64	106	173	182	175	51	5	-
White, both sexes	44	53	96	79	103	163	251	277	231	58	6	-
Male	26	26	50	30	46	67	90	105	66	13	1	-
Female	18	27	46	49	57	96	161	172	165	45	5	-
Black, both sexes	8	12	16	13	12	17	16	15	12	4	3	-
Male	3	6	6	5	5	9	7	6	4	-	3	-
Female	5	6	10	8	7	8	9	9	8	4	-	-
American Indian 1/, both sexes	-	3	-	3	-	1	3	1	2	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	3	-	1	2	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	2	3	1	3	-	-
Male	-	1	-	-	-	-	1	2	1	1	-	-
Female	-	-	-	-	-	1	1	1	-	2	-	-
Alabama												
All races, both sexes	2	-	-	6	3	3	4	4	3	-	1	-
Male	1	-	-	3	2	1	-	1	-	-	1	-
Female	1	-	-	3	1	2	4	3	3	-	-	-
White, both sexes	2	-	-	5	1	3	4	4	3	-	-	-
Male	1	-	-	3	1	1	-	1	-	-	-	-
Female	1	-	-	2	-	2	4	3	3	-	-	-
Black, both sexes	-	-	-	1	2	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	1	2	2	3	4	8	4	-	-	-
Male	-	-	1	1	1	2	2	5	2	-	-	-
Female	-	-	-	1	1	1	2	3	2	-	-	-
White, both sexes	-	-	1	2	2	3	4	7	4	-	-	-
Male	-	-	1	1	1	2	2	4	2	-	-	-
Female	-	-	-	1	1	1	2	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Arkansas												
All races, both sexes	-	1	-	1	1	1	3	5	3	-	-	-
Male	-	-	-	-	-	1	-	2	1	-	-	-
Female	-	1	-	1	1	-	3	3	2	-	-	-
White, both sexes	-	-	-	1	1	-	3	4	3	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	1	1	-	3	3	2	-	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	8	15	13	9	29	23	37	17	10	1	-
Male	4	4	7	4	4	12	11	18	4	2	1	-
Female	4	4	8	9	5	17	12	19	13	8	-	-
White, both sexes	7	8	12	11	8	24	19	36	16	9	1	-
Male	4	4	6	4	4	11	11	17	3	1	1	-
Female	3	4	6	7	4	13	8	19	13	8	-	-
Black, both sexes	1	-	3	-	1	4	2	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	1	-	2	-	1	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Colorado												
All races, both sexes	-	-	1	1	2	2	4	3	3	1	-	-
Male	-	-	-	-	2	-	1	2	1	-	-	-
Female	-	-	1	1	-	2	3	1	2	1	-	-
White, both sexes	-	-	1	1	2	2	3	3	3	1	-	-
Male	-	-	-	-	2	-	1	2	1	-	-	-
Female	-	-	1	1	-	2	2	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	1	-	3	1	2	4	5	2	-	-
Male	-	-	1	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	3	1	1	3	4	2	-	-
White, both sexes	-	1	1	-	3	1	2	3	5	2	-	-
Male	-	-	1	-	-	-	1	1	1	-	-	-
Female	-	1	-	-	3	1	1	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Delaware												
All races, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Florida												
All races, both sexes	6	6	1	6	4	11	14	13	10	2	-	-
Male	4	3	1	1	4	6	3	6	6	1	-	-
Female	2	3	-	5	-	5	11	7	4	1	-	-
White, both sexes	5	6	1	4	4	10	13	11	10	2	-	-
Male	3	3	1	-	4	6	3	5	6	1	-	-
Female	2	3	-	4	-	4	10	6	4	1	-	-
Black, both sexes	1	-	-	2	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	4	3	5	5	2	5	1	6	-	-	-
Male	1	3	-	1	1	2	2	-	2	-	-	-
Female	1	1	3	4	4	-	3	1	4	-	-	-
White, both sexes	1	2	3	5	3	1	5	-	6	-	-	-
Male	-	2	-	1	1	1	2	-	2	-	-	-
Female	1	-	3	4	2	-	3	-	4	-	-	-
Black, both sexes	1	2	-	-	2	1	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Hawaii												
All races, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	1	-	-	-	1	1	1	2	2	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	2	-	-	-
White, both sexes	1	-	-	-	1	1	1	2	2	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Illinois												
All races, both sexes	-	4	6	2	4	5	6	7	11	3	2	-
Male	-	1	2	2	2	3	4	4	6	-	2	-
Female	-	3	4	-	2	2	2	3	5	3	-	-
White, both sexes	-	4	5	2	4	5	6	7	11	2	-	-
Male	-	1	2	2	2	3	4	4	6	-	-	-
Female	-	3	3	-	2	2	2	3	5	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	2	-
Male	-	-	-	-	-	-	-	-	-	-	2	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	3	1	7	7	6	5	-	-	-
Male	-	-	-	-	-	2	5	3	-	-	-	-
Female	-	-	-	3	1	5	2	3	5	-	-	-
White, both sexes	-	-	-	3	1	7	7	6	5	-	-	-
Male	-	-	-	-	-	2	5	3	-	-	-	-
Female	-	-	-	3	1	5	2	3	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Iowa												
All races, both sexes	-	-	1	-	3	-	5	4	4	3	-	-
Male	-	-	-	-	1	-	2	1	1	-	-	-
Female	-	-	1	-	2	-	3	3	3	3	-	-
White, both sexes	-	-	1	-	3	-	5	4	4	3	-	-
Male	-	-	-	-	1	-	2	1	1	-	-	-
Female	-	-	1	-	2	-	3	3	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	2	2	3	2	5	1	-	-
Male	-	-	-	1	-	-	-	1	2	-	-	-
Female	-	-	-	-	2	2	3	1	3	1	-	-
White, both sexes	-	-	-	1	2	2	3	2	5	1	-	-
Male	-	-	-	1	-	-	-	1	2	-	-	-
Female	-	-	-	-	2	2	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Kentucky												
All races, both sexes	1	-	3	-	2	4	8	3	3	-	-	-
Male	1	-	3	-	-	2	4	1	2	-	-	-
Female	-	-	-	-	2	2	4	2	1	-	-	-
White, both sexes	1	-	3	-	2	4	8	3	3	-	-	-
Male	1	-	3	-	-	2	4	1	2	-	-	-
Female	-	-	-	-	2	2	4	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	1	2	2	-	2	4	6	3	-	-	-
Male	1	1	2	1	-	-	-	2	-	-	-	-
Female	1	-	-	1	-	2	4	4	3	-	-	-
White, both sexes	1	-	2	-	-	2	4	5	3	-	-	-
Male	1	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	4	4	3	-	-	-
Black, both sexes	1	1	-	2	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Maine												
All races, both sexes	1	1	-	-	-	-	4	3	-	-	-	-
Male	1	1	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
White, both sexes	1	1	-	-	-	-	4	3	-	-	-	-
Male	1	1	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	1	1	1	1	8	7	5	1	-	-
Male	1	1	-	-	-	-	2	3	3	-	-	-
Female	-	-	1	1	1	1	6	4	2	1	-	-
White, both sexes	1	-	-	1	1	1	5	7	5	-	-	-
Male	1	-	-	-	-	-	2	3	3	-	-	-
Female	-	-	-	1	1	1	3	4	2	-	-	-
Black, both sexes	-	1	1	-	-	-	3	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Massachusetts												
All races, both sexes	-	1	-	1	6	4	3	6	8	2	-	-
Male	-	-	-	-	2	2	2	1	4	1	-	-
Female	-	1	-	1	4	2	1	5	4	1	-	-
White, both sexes	-	1	-	1	6	4	3	6	8	2	-	-
Male	-	-	-	-	2	2	2	1	4	1	-	-
Female	-	1	-	1	4	2	1	5	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	2	8	4	4	6	10	9	8	1	1	-
Male	-	1	5	3	2	2	4	5	1	1	-	-
Female	4	1	3	1	2	4	6	4	7	-	1	-
White, both sexes	3	2	8	3	4	6	8	9	8	1	1	-
Male	-	1	5	2	2	2	2	5	1	1	-	-
Female	3	1	3	1	2	4	6	4	7	-	1	-
Black, both sexes	1	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Minnesota												
All races, both sexes	1	-	2	2	1	1	4	7	9	-	-	-
Male	-	-	1	2	1	-	1	2	3	-	-	-
Female	1	-	1	-	-	1	3	5	6	-	-	-
White, both sexes	1	-	2	2	1	1	4	7	9	-	-	-
Male	-	-	1	2	1	-	1	2	3	-	-	-
Female	1	-	1	-	-	1	3	5	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	1	-	-	5	2	5	1	1	-	-
Male	1	-	-	-	-	4	1	1	-	-	-	-
Female	-	1	1	-	-	1	1	4	1	1	-	-
White, both sexes	1	1	1	-	-	3	2	4	-	1	-	-
Male	1	-	-	-	-	3	1	1	-	-	-	-
Female	-	1	1	-	-	-	1	3	-	1	-	-
Black, both sexes	-	-	-	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Missouri												
All races, both sexes	1	-	3	1	2	5	12	7	6	2	-	-
Male	-	-	1	-	1	2	7	4	2	1	-	-
Female	1	-	2	1	1	3	5	3	4	1	-	-
White, both sexes	1	-	3	1	2	5	12	7	4	2	-	-
Male	-	-	1	-	1	2	7	4	1	1	-	-
Female	1	-	2	1	1	3	5	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Nebraska												
All races, both sexes	-	-	1	-	1	2	1	3	-	2	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	3	-	2	-	-
White, both sexes	-	-	1	-	-	2	1	3	-	2	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	3	-	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	2	-	1	1	2	1	-	-	-	-
Male	1	-	2	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
White, both sexes	1	-	2	-	1	1	2	-	-	-	-	-
Male	1	-	2	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	1	-	5	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	4	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	5	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	5	-	2	4	6	7	5	3	-	-
Male	-	1	3	-	2	4	2	-	1	2	-	-
Female	1	1	2	-	-	-	4	7	4	1	-	-
White, both sexes	-	2	4	-	2	3	6	6	5	3	-	-
Male	-	1	2	-	2	3	2	-	1	2	-	-
Female	-	1	2	-	-	-	4	6	4	1	-	-
Black, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
New Mexico												
All races, both sexes	-	1	1	1	-	1	2	1	2	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	1	-	-	1	1	1	2	-	-	-
White, both sexes	-	1	1	1	-	1	1	1	2	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	1	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	5	10	10	9	15	16	15	14	6	-	-
Male	2	4	7	4	8	3	4	9	4	3	-	-
Female	-	1	3	6	1	12	12	6	10	3	-	-
White, both sexes	1	2	6	6	7	13	15	13	13	4	-	-
Male	1	2	5	3	6	3	4	7	4	3	-	-
Female	-	-	1	3	1	10	11	6	9	1	-	-
Black, both sexes	1	2	4	3	2	2	1	2	1	1	-	-
Male	1	1	2	1	2	-	-	2	-	-	-	-
Female	-	1	2	2	-	2	1	-	1	1	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
North Carolina												
All races, both sexes	1	5	4	1	3	8	3	7	3	2	-	-
Male	1	3	3	-	1	3	1	2	3	-	-	-
Female	-	2	1	1	2	5	2	5	-	2	-	-
White, both sexes	1	2	3	1	3	8	3	6	1	1	-	-
Male	1	1	2	-	1	3	1	2	1	-	-	-
Female	-	1	1	1	2	5	2	4	-	1	-	-
Black, both sexes	-	2	1	-	-	-	-	1	2	1	-	-
Male	-	2	1	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	-	1	1	2	1	-	-	-
Male	-	1	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Ohio												
All races, both sexes	1	4	5	4	6	10	15	18	10	3	-	-
Male	-	3	1	1	4	6	4	7	2	-	-	-
Female	1	1	4	3	2	4	11	11	8	3	-	-
White, both sexes	1	4	5	4	6	10	14	18	9	3	-	-
Male	-	3	1	1	4	6	3	7	1	-	-	-
Female	1	1	4	3	2	4	11	11	8	3	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	4	2	2	7	2	2	5	3	2	-	-
Male	-	2	1	1	-	1	1	3	-	-	-	-
Female	-	2	1	1	7	1	1	2	3	2	-	-
White, both sexes	-	4	2	1	7	2	1	5	2	2	-	-
Male	-	2	1	-	-	1	-	3	-	-	-	-
Female	-	2	1	1	7	1	1	2	2	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Oregon												
All races, both sexes	-	-	2	3	1	2	6	6	6	3	-	-
Male	-	-	1	-	1	1	2	-	1	-	-	-
Female	-	-	1	3	-	1	4	6	5	3	-	-
White, both sexes	-	-	2	3	1	2	6	6	6	3	-	-
Male	-	-	1	-	1	1	2	-	1	-	-	-
Female	-	-	1	3	-	1	4	6	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	1	7	2	5	9	22	18	18	4	1	-
Male	3	1	3	1	3	4	5	3	4	1	-	-
Female	1	-	4	1	2	5	17	15	14	3	1	-
White, both sexes	4	1	7	2	5	8	22	18	18	4	1	-
Male	3	1	3	1	3	3	5	3	4	1	-	-
Female	1	-	4	1	2	5	17	15	14	3	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Rhode Island												
All races, both sexes	-	-	-	-	2	2	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	2	1	-	1	-	-	-
White, both sexes	-	-	-	-	2	2	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	2	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	1	2	4	3	6	3	1	-	-
Male	-	1	-	1	-	1	1	1	2	-	-	-
Female	1	-	1	-	2	3	2	5	1	1	-	-
White, both sexes	1	1	-	1	2	2	3	5	2	1	-	-
Male	-	1	-	1	-	-	1	1	2	-	-	-
Female	1	-	-	-	2	2	2	4	-	1	-	-
Black, both sexes	-	-	1	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	3	2	1	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	1	1	-	2	2	1	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	3	2	5	4	3	11	5	5	1	1	-
Male	2	1	-	2	2	1	3	3	-	-	-	-
Female	-	2	2	3	2	2	8	2	5	1	1	-
White, both sexes	2	2	2	3	2	2	9	5	2	1	1	-
Male	2	-	-	2	1	-	2	3	-	-	-	-
Female	-	2	2	1	1	2	7	2	2	1	1	-
Black, both sexes	-	-	-	2	2	1	2	-	2	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	2	1	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Texas												
All races, both sexes	3	3	10	7	10	13	9	18	17	4	2	-
Male	2	1	7	2	3	6	2	8	6	2	-	-
Female	1	2	3	5	7	7	7	10	11	2	2	-
White, both sexes	3	1	8	7	9	12	8	18	16	4	2	-
Male	2	1	6	2	3	5	1	8	6	2	-	-
Female	1	-	2	5	6	7	7	10	10	2	2	-
Black, both sexes	-	2	2	-	1	1	1	-	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	2	1	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	2	3	2	1	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	2	3	2	1	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Vermont												
All races, both sexes	-	-	-	1	-	2	1	2	2	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	1	-	1	-	-	2	-	-	-
White, both sexes	-	-	-	1	-	2	1	2	2	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	1	-	1	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	3	3	1	3	1	7	5	7	-	-	-
Male	-	-	-	-	1	1	3	1	1	-	-	-
Female	1	3	3	1	2	-	4	4	6	-	-	-
White, both sexes	-	2	1	-	2	1	6	3	7	-	-	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	-	2	1	-	1	-	4	2	6	-	-	-
Black, both sexes	1	1	2	1	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	2	1	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Washington												
All races, both sexes	1	2	3	3	1	3	5	8	5	1	-	-
Male	-	1	2	2	-	1	2	1	2	-	-	-
Female	1	1	1	1	1	2	3	7	3	1	-	-
White, both sexes	1	2	3	3	1	3	5	7	5	1	-	-
Male	-	1	2	2	-	1	2	-	2	-	-	-
Female	1	1	1	1	1	2	3	7	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	1	1	-	-	5	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	1	-	-	4	-	2	1	-	-
White, both sexes	-	1	1	1	-	-	5	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	1	-	-	4	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Wisconsin												
All races, both sexes	1	2	2	1	1	1	6	11	10	2	-	-
Male	-	1	-	1	-	-	2	3	2	-	-	-
Female	1	1	2	-	1	1	4	8	8	2	-	-
White, both sexes	1	2	2	1	1	1	6	11	9	2	-	-
Male	-	1	-	1	-	-	2	3	2	-	-	-
Female	1	1	2	-	1	1	4	8	7	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	1	-	-	-	1	1	1	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	1	-	1	-	-	-	1	1	1	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
United States												
All races, both sexes	897	-	-	-	3	4	6	10	13	12	24	48
Male	351	-	-	-	3	2	3	6	5	4	12	24
Female	546	-	-	-	-	2	3	4	8	8	12	24
White, both sexes	835	-	-	-	3	3	4	8	8	10	19	45
Male	330	-	-	-	3	1	2	4	2	4	10	22
Female	505	-	-	-	-	2	2	4	6	6	9	23
Black, both sexes	50	-	-	-	-	1	2	1	3	1	5	2
Male	16	-	-	-	-	1	1	1	2	-	2	2
Female	34	-	-	-	-	-	1	-	1	1	3	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	2	1	-	1
Male	3	-	-	-	-	-	-	1	1	-	-	-
Female	5	-	-	-	-	-	-	-	1	1	-	1
Alabama												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	14	-	-	-	-	-	-	-	-	1	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Arkansas												
All races, both sexes	8	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	83	-	-	-	-	-	1	1	2	3	4	8
Male	32	-	-	-	-	-	1	1	1	-	2	4
Female	51	-	-	-	-	-	-	-	1	3	2	4
White, both sexes	73	-	-	-	-	-	-	1	2	2	3	7
Male	30	-	-	-	-	-	-	1	1	-	2	3
Female	43	-	-	-	-	-	-	-	1	2	1	4
Black, both sexes	6	-	-	-	-	-	1	-	-	-	1	1
Male	2	-	-	-	-	-	1	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Colorado												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	1	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Florida												
All races, both sexes	77	-	-	-	-	1	-	-	1	1	3	5
Male	31	-	-	-	-	-	-	-	-	-	1	3
Female	46	-	-	-	-	1	-	-	1	1	2	2
White, both sexes	75	-	-	-	-	1	-	-	1	1	2	5
Male	30	-	-	-	-	-	-	-	-	-	1	3
Female	45	-	-	-	-	1	-	-	1	1	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	21	-	-	-	-	-	-	-	2	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	2	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Illinois												
All races, both sexes	42	-	-	-	1	-	-	2	-	-	1	2
Male	14	-	-	-	1	-	-	1	-	-	1	1
Female	28	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	37	-	-	-	1	-	-	2	-	-	1	2
Male	13	-	-	-	1	-	-	1	-	-	1	1
Female	24	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Iowa												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	1	-	-	-	-	-	-
Male	7	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Kentucky												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	-	-	-	-	-	-	1	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Massachusetts												
All races, both sexes	32	-	-	-	-	-	-	2	-	1	1	-
Male	14	-	-	-	-	-	-	2	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	31	-	-	-	-	-	-	1	-	1	1	-
Male	13	-	-	-	-	-	-	1	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	27	-	-	-	-	1	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	19	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	1	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Minnesota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Missouri												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Nebraska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	30	-	-	-	-	-	1	1	1	1	1	1
Male	16	-	-	-	-	-	-	1	1	-	1	1
Female	14	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	25	-	-	-	-	-	1	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	1	1	-	1	-
Male	4	-	-	-	-	-	-	1	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
New Mexico												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	55	-	-	-	-	-	1	-	-	1	2	2
Male	25	-	-	-	-	-	1	-	-	1	1	-
Female	30	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	52	-	-	-	-	-	1	-	-	1	1	1
Male	24	-	-	-	-	-	1	-	-	1	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
North Carolina												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Ohio												
All races, both sexes	44	-	-	-	1	-	1	-	2	-	-	1
Male	17	-	-	-	1	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	42	-	-	-	1	-	1	-	2	-	-	1
Male	16	-	-	-	1	-	-	-	1	-	-	1
Female	26	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	11	-	-	-	-	-	1	-	-	-	-	2
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	11	-	-	-	-	-	1	-	-	-	-	2
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Oregon												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	4
Male	10	-	-	-	-	-	-	-	-	-	-	3
Female	25	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	4
Male	10	-	-	-	-	-	-	-	-	-	-	3
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Rhode Island												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Texas												
All races, both sexes	48	-	-	-	-	1	-	1	1	-	2	2
Male	19	-	-	-	-	1	-	-	1	-	2	1
Female	29	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	44	-	-	-	-	-	-	1	-	-	2	2
Male	17	-	-	-	-	-	-	-	-	-	2	1
Female	27	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	4	-	-	-	-	1	-	-	1	-	-	-
Male	2	-	-	-	-	1	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	20	-	-	-	-	-	-	-	2	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	1	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Washington												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Crohn's disease and ulcerative colitis (K50-K51)												
Wisconsin												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
United States												
All races, both sexes	56	52	60	75	90	124	142	112	53	10	3	-
Male	23	25	30	39	38	51	48	26	9	3	-	-
Female	33	27	30	36	52	73	94	86	44	7	3	-
White, both sexes	45	47	57	69	87	123	139	103	53	10	2	-
Male	19	23	29	36	38	51	48	26	9	3	-	-
Female	26	24	28	33	49	72	91	77	44	7	2	-
Black, both sexes	8	4	3	5	3	1	3	7	-	-	1	-
Male	2	2	1	2	-	-	-	-	-	-	-	-
Female	6	2	2	3	3	1	3	7	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	2	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	3	1	-	2	1	2	-	3	1	-	-	-
Male	1	1	-	1	-	1	-	1	-	-	-	-
Female	2	-	-	1	1	1	-	2	1	-	-	-
White, both sexes	3	1	-	1	1	2	-	2	1	-	-	-
Male	1	1	-	-	-	1	-	1	-	-	-	-
Female	2	-	-	1	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	1	-	2	3	3	1	2	-	-	-
Male	2	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	3	2	1	2	-	-	-
White, both sexes	-	-	1	-	2	3	3	1	2	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	3	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Arkansas												
All races, both sexes	1	-	1	-	3	2	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	2	2	-	-	-	-	-	-
White, both sexes	1	-	1	-	3	2	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	2	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	8	5	3	9	8	9	18	-	2	-	-
Male	1	2	1	1	5	4	3	5	-	1	-	-
Female	1	6	4	2	4	4	6	13	-	1	-	-
White, both sexes	1	7	5	3	9	8	8	15	-	2	-	-
Male	1	2	1	1	5	4	3	5	-	1	-	-
Female	-	5	4	2	4	4	5	10	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Colorado												
All races, both sexes	1	1	-	3	1	3	2	2	4	-	-	-
Male	-	1	-	2	1	2	1	-	2	-	-	-
Female	1	-	-	1	-	1	1	2	2	-	-	-
White, both sexes	1	1	-	3	1	3	2	2	4	-	-	-
Male	-	1	-	2	1	2	1	-	2	-	-	-
Female	1	-	-	1	-	1	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	1	1	2	3	-	2	2	-	-	-	-
Male	-	-	1	2	2	-	-	2	-	-	-	-
Female	2	1	-	-	1	-	2	-	-	-	-	-
White, both sexes	1	1	1	2	3	-	2	2	-	-	-	-
Male	-	-	1	2	2	-	-	2	-	-	-	-
Female	1	1	-	-	1	-	2	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Delaware												
All races, both sexes	-	-	-	2	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Florida												
All races, both sexes	2	2	11	10	11	9	13	4	3	-	1	-
Male	1	2	7	6	3	2	4	1	1	-	-	-
Female	1	-	4	4	8	7	9	3	2	-	1	-
White, both sexes	2	2	11	9	11	9	13	4	3	-	1	-
Male	1	2	7	5	3	2	4	1	1	-	-	-
Female	1	-	4	4	8	7	9	3	2	-	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	1	2	2	3	2	2	3	-	-	-	-
Male	1	1	1	1	-	1	1	-	-	-	-	-
Female	2	-	1	1	3	1	1	3	-	-	-	-
White, both sexes	3	-	1	2	3	2	2	2	-	-	-	-
Male	1	-	1	1	-	1	1	-	-	-	-	-
Female	2	-	-	1	3	1	1	2	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	1	3	-	-	2	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	2	1	-	-	-	-	-
White, both sexes	1	1	3	-	-	2	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	2	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Illinois												
All races, both sexes	5	2	2	2	4	8	4	6	3	-	-	-
Male	3	-	-	1	-	3	1	1	1	-	-	-
Female	2	2	2	1	4	5	3	5	2	-	-	-
White, both sexes	3	2	2	2	2	8	4	5	3	-	-	-
Male	2	-	-	1	-	3	1	1	1	-	-	-
Female	1	2	2	1	2	5	3	4	2	-	-	-
Black, both sexes	2	-	-	-	2	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	2	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	2	2	1	2	4	-	1	-	-	-
Male	1	-	1	1	-	-	1	-	-	-	-	-
Female	1	-	1	1	1	2	3	-	1	-	-	-
White, both sexes	2	-	1	2	1	2	4	-	1	-	-	-
Male	1	-	1	1	-	-	1	-	-	-	-	-
Female	1	-	-	1	1	2	3	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Iowa												
All races, both sexes	1	-	-	1	-	-	4	2	1	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	1	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	1	-	-	1	-	-	4	2	1	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	1	-	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	2	1	1	1	2	1	-	-	-
Male	-	-	1	2	1	1	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	1	1	1	1	1	2	1	-	-	-
Male	-	-	1	1	1	1	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Kentucky												
All races, both sexes	2	-	1	-	2	1	4	1	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	-	1	1	3	1	-	-	-	-
White, both sexes	2	-	1	-	2	1	4	1	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	-	1	1	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	2	4	2	2	-	-	-	-
Male	-	-	-	-	1	3	-	-	-	-	-	-
Female	1	-	-	-	1	1	2	2	-	-	-	-
White, both sexes	1	-	-	-	2	4	2	2	-	-	-	-
Male	-	-	-	-	1	3	-	-	-	-	-	-
Female	1	-	-	-	1	1	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Maine												
All races, both sexes	1	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	1	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	1	1	1	1	2	1	-	-	-	-
Male	1	2	1	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	1	2	-	1	1	1	2	1	-	-	-	-
Male	1	2	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Massachusetts												
All races, both sexes	1	1	-	3	4	4	6	7	2	-	-	-
Male	1	-	-	3	3	2	1	2	-	-	-	-
Female	-	1	-	-	1	2	5	5	2	-	-	-
White, both sexes	1	1	-	3	4	4	6	7	2	-	-	-
Male	1	-	-	3	3	2	1	2	-	-	-	-
Female	-	1	-	-	1	2	5	5	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	1	1	1	3	5	5	2	4	-	-	-
Male	1	-	1	-	2	3	-	-	-	-	-	-
Female	1	1	-	1	1	2	5	2	4	-	-	-
White, both sexes	2	1	1	1	3	5	5	2	4	-	-	-
Male	1	-	1	-	2	3	-	-	-	-	-	-
Female	1	1	-	1	1	2	5	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Minnesota												
All races, both sexes	-	-	1	3	-	2	3	2	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	3	-	1	1	2	-	-	-	-
White, both sexes	-	-	1	3	-	2	3	2	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	3	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Missouri												
All races, both sexes	2	1	1	2	1	2	1	2	2	2	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	2	-	1	2	-	2	1	1	2	2	-	-
White, both sexes	2	1	1	2	1	2	1	2	2	2	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	2	-	1	2	-	2	1	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Nebraska												
All races, both sexes	-	1	-	1	1	-	4	2	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	1	-	1	1	-	2	2	-	1	-	-
White, both sexes	-	1	-	1	1	-	4	2	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	1	-	1	1	-	2	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	-	1	1	3	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	3	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
New Hampshire												
All races, both sexes	-	-	-	1	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	3	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	3	3	3	1	5	5	2	1	-	-	-
Male	1	2	2	2	-	2	3	-	-	-	-	-
Female	-	1	1	1	1	3	2	2	1	-	-	-
White, both sexes	1	1	3	3	1	5	5	2	1	-	-	-
Male	1	1	2	2	-	2	3	-	-	-	-	-
Female	-	-	1	1	1	3	2	2	1	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
New Mexico												
All races, both sexes	-	1	1	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	1	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	2	1	3	10	14	8	6	4	-	-	-
Male	1	2	-	2	5	7	3	1	1	-	-	-
Female	-	-	1	1	5	7	5	5	3	-	-	-
White, both sexes	1	2	1	3	10	14	7	6	4	-	-	-
Male	1	2	-	2	5	7	3	1	1	-	-	-
Female	-	-	1	1	5	7	4	5	3	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
North Carolina												
All races, both sexes	2	-	-	4	6	2	4	2	2	-	-	-
Male	1	-	-	2	2	1	1	2	-	-	-	-
Female	1	-	-	2	4	1	3	-	2	-	-	-
White, both sexes	2	-	-	3	6	2	4	2	2	-	-	-
Male	1	-	-	2	2	1	1	2	-	-	-	-
Female	1	-	-	1	4	1	3	-	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Ohio												
All races, both sexes	2	3	3	2	4	9	7	7	2	-	-	-
Male	1	3	2	1	1	4	1	1	-	-	-	-
Female	1	-	1	1	3	5	6	6	2	-	-	-
White, both sexes	1	3	3	2	4	9	6	7	2	-	-	-
Male	-	3	2	1	1	4	1	1	-	-	-	-
Female	1	-	1	1	3	5	5	6	2	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	1	2	3	-	1	-	-	-	-
Male	-	1	-	1	-	3	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
White, both sexes	-	1	-	1	2	3	-	1	-	-	-	-
Male	-	1	-	1	-	3	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Oregon												
All races, both sexes	1	2	4	1	-	3	-	1	4	-	-	-
Male	1	1	3	-	-	1	-	-	1	-	-	-
Female	-	1	1	1	-	2	-	1	3	-	-	-
White, both sexes	1	2	4	1	-	3	-	1	4	-	-	-
Male	1	1	3	-	-	1	-	-	1	-	-	-
Female	-	1	1	1	-	2	-	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	2	2	3	13	5	4	1	-	-
Male	-	-	-	1	2	-	3	1	-	-	-	-
Female	-	1	-	1	-	3	10	4	4	1	-	-
White, both sexes	-	1	-	2	2	3	13	5	4	1	-	-
Male	-	-	-	1	2	-	3	1	-	-	-	-
Female	-	1	-	1	-	3	10	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	1	-	-	-
White, both sexes	-	1	-	-	-	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	2	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	1	-	1	1	-	-	-	-	-
White, both sexes	-	2	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	1	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
South Dakota												
All races, both sexes	-	1	1	-	-	-	2	1	-	-	-	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	1	-	-	-	2	1	-	-	-	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	1	1	4	2	2	-	4	1	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	3	-	1	3	2	1	-	4	1	-	-	-
White, both sexes	1	1	1	3	1	2	-	3	1	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	2	1	1	-	3	1	-	-	-
Black, both sexes	2	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Texas												
All races, both sexes	5	7	5	3	3	6	6	2	3	-	1	-
Male	-	2	3	1	1	3	3	1	-	-	-	-
Female	5	5	2	2	2	3	3	1	3	-	1	-
White, both sexes	4	7	5	3	3	6	6	2	3	-	-	-
Male	-	2	3	1	1	3	3	1	-	-	-	-
Female	4	5	2	2	2	3	3	1	3	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	2	2	-	1	1	-	-	-
Male	-	-	1	1	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes	-	-	1	1	2	2	-	1	1	-	-	-
Male	-	-	1	1	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Vermont												
All races, both sexes	-	-	-	1	-	2	-	2	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	2	-	1	-	-
White, both sexes	-	-	-	1	-	2	-	2	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	2	1	2	1	4	2	1	1	-	-	-
Male	1	-	-	1	1	1	2	-	-	-	-	-
Female	1	2	1	1	-	3	-	1	1	-	-	-
White, both sexes	1	1	1	2	1	3	2	1	1	-	-	-
Male	1	-	-	1	1	1	2	-	-	-	-	-
Female	-	1	1	1	-	2	-	1	1	-	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Washington												
All races, both sexes	-	1	1	2	2	1	4	3	2	1	-	-
Male	-	-	-	2	1	-	2	1	1	1	-	-
Female	-	1	1	-	1	1	2	2	1	-	-	-
White, both sexes	-	1	1	2	2	1	4	2	2	1	-	-
Male	-	-	-	2	1	-	2	1	1	1	-	-
Female	-	1	1	-	1	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Wisconsin												
All races, both sexes	3	-	1	1	-	3	2	1	1	1	-	-
Male	2	-	-	-	-	2	-	-	1	1	-	-
Female	1	-	1	1	-	1	2	1	-	-	-	-
White, both sexes	3	-	1	1	-	3	2	1	1	1	-	-
Male	2	-	-	-	-	2	-	-	1	1	-	-
Female	1	-	1	1	-	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
United States												
All races, both sexes	14,442	129	25	8	13	16	23	24	30	50	95	193
Male	5,167	82	17	1	10	10	9	13	17	23	44	104
Female	9,275	47	8	7	3	6	14	11	13	27	51	89
White, both sexes	12,645	78	16	8	7	12	16	22	23	34	72	147
Male	4,402	47	10	1	5	7	6	11	13	16	35	77
Female	8,243	31	6	7	2	5	10	11	10	18	37	70
Black, both sexes	1,514	45	7	-	5	3	7	2	5	14	23	44
Male	641	33	6	-	5	2	3	2	3	6	9	25
Female	873	12	1	-	-	1	4	-	2	8	14	19
American Indian 1/, both sexes	90	3	1	-	-	-	-	-	2	1	-	1
Male	36	-	-	-	-	-	-	-	1	-	-	1
Female	54	3	1	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	193	3	1	-	1	1	-	-	-	1	-	1
Male	88	2	1	-	-	1	-	-	-	1	-	1
Female	105	1	-	-	1	-	-	-	-	-	-	-
Alabama												
All races, both sexes	243	2	-	-	-	-	1	1	-	-	3	4
Male	79	1	-	-	-	-	1	-	-	-	-	2
Female	164	1	-	-	-	-	-	1	-	-	3	2
White, both sexes	185	-	-	-	-	-	-	1	-	-	1	3
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	128	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	57	2	-	-	-	-	1	-	-	-	2	1
Male	21	1	-	-	-	-	1	-	-	-	-	-
Female	36	1	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Alaska												
All races, both sexes	24	1	-	-	-	-	-	-	-	1	1	-
Male	7	1	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	14	1	-	-	-	-	-	-	-	1	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	253	4	-	1	-	-	-	-	1	-	2	4
Male	102	2	-	-	-	-	-	-	1	-	2	3
Female	151	2	-	1	-	-	-	-	-	-	-	1
White, both sexes	233	4	-	1	-	-	-	-	1	-	1	4
Male	91	2	-	-	-	-	-	-	1	-	1	3
Female	142	2	-	1	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Arkansas												
All races, both sexes	186	1	1	-	-	-	1	-	-	1	1	-
Male	63	1	-	-	-	-	-	-	-	-	-	-
Female	123	-	1	-	-	-	1	-	-	1	1	-
White, both sexes	164	-	1	-	-	-	1	-	-	-	1	-
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	111	-	1	-	-	-	1	-	-	-	1	-
Black, both sexes	22	1	-	-	-	-	-	-	-	1	-	-
Male	10	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,373	9	8	1	4	4	1	1	4	4	8	20
Male	508	4	5	-	3	2	1	1	2	1	3	12
Female	865	5	3	1	1	2	-	-	2	3	5	8
White, both sexes	1,160	7	5	1	2	4	1	1	3	2	5	15
Male	405	2	3	-	2	2	1	1	1	1	2	8
Female	755	5	2	1	-	2	-	-	2	1	3	7
Black, both sexes	105	2	1	-	1	-	-	-	-	2	3	5
Male	55	2	1	-	1	-	-	-	-	-	1	4
Female	50	-	-	-	-	-	-	-	-	2	2	1
American Indian 1/, both sexes	11	-	1	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	97	-	1	-	1	-	-	-	-	-	-	-
Male	45	-	1	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Colorado												
All races, both sexes	176	2	-	2	-	-	2	1	1	-	1	-
Male	64	2	-	-	-	-	1	1	-	-	1	-
Female	112	-	-	2	-	-	1	-	1	-	-	-
White, both sexes	172	2	-	2	-	-	2	1	1	-	1	-
Male	63	2	-	-	-	-	1	1	-	-	1	-
Female	109	-	-	2	-	-	1	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	206	2	-	-	-	-	2	-	-	-	1	2
Male	62	1	-	-	-	-	-	-	-	-	1	1
Female	144	1	-	-	-	-	2	-	-	-	-	1
White, both sexes	194	1	-	-	-	-	2	-	-	-	1	1
Male	61	-	-	-	-	-	-	-	-	-	1	1
Female	133	1	-	-	-	-	2	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	-	-	-	-	1
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Delaware												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Florida												
All races, both sexes	874	7	-	-	-	-	-	-	4	5	5	13
Male	313	5	-	-	-	-	-	-	2	2	1	3
Female	561	2	-	-	-	-	-	-	2	3	4	10
White, both sexes	791	3	-	-	-	-	-	-	3	3	4	11
Male	282	2	-	-	-	-	-	-	2	1	1	3
Female	509	1	-	-	-	-	-	-	1	2	3	8
Black, both sexes	77	3	-	-	-	-	-	-	1	2	1	2
Male	30	2	-	-	-	-	-	-	-	1	-	-
Female	47	1	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	382	6	2	1	1	-	-	-	-	2	5	5
Male	132	4	2	-	-	-	-	-	-	1	1	1
Female	250	2	-	1	1	-	-	-	-	1	4	4
White, both sexes	291	-	2	1	1	-	-	-	-	1	2	3
Male	98	-	2	-	-	-	-	-	-	1	-	1
Female	193	-	-	1	1	-	-	-	-	-	2	2
Black, both sexes	89	5	-	-	-	-	-	-	-	1	3	2
Male	34	4	-	-	-	-	-	-	-	-	1	-
Female	55	1	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Hawaii												
All races, both sexes	46	-	-	-	-	-	1	-	-	1	-	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	1	-	-
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	77	2	-	-	-	1	-	-	-	-	-	1
Male	31	1	-	-	-	1	-	-	-	-	-	-
Female	46	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	77	2	-	-	-	1	-	-	-	-	-	1
Male	31	1	-	-	-	1	-	-	-	-	-	-
Female	46	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Illinois												
All races, both sexes	651	7	-	-	-	1	-	2	3	2	5	8
Male	237	6	-	-	-	1	-	1	3	1	4	6
Female	414	1	-	-	-	-	-	1	-	1	1	2
White, both sexes	577	6	-	-	-	1	-	2	2	2	3	4
Male	204	5	-	-	-	1	-	1	2	1	3	3
Female	373	1	-	-	-	-	-	1	-	1	-	1
Black, both sexes	68	1	-	-	-	-	-	-	1	-	2	4
Male	29	1	-	-	-	-	-	-	1	-	1	3
Female	39	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	346	7	1	-	-	-	-	-	-	1	-	6
Male	113	4	1	-	-	-	-	-	-	-	-	3
Female	233	3	-	-	-	-	-	-	-	1	-	3
White, both sexes	323	6	1	-	-	-	-	-	-	1	-	6
Male	104	3	1	-	-	-	-	-	-	-	-	3
Female	219	3	-	-	-	-	-	-	-	1	-	3
Black, both sexes	23	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Iowa												
All races, both sexes	171	1	-	-	1	-	1	-	-	-	-	1
Male	55	1	-	-	-	-	-	-	-	-	-	1
Female	116	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	168	1	-	-	1	-	1	-	-	-	-	1
Male	55	1	-	-	-	-	-	-	-	-	-	1
Female	113	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	176	3	1	-	-	-	-	2	-	1	1	2
Male	64	1	1	-	-	-	-	1	-	1	-	1
Female	112	2	-	-	-	-	-	1	-	-	1	1
White, both sexes	166	3	-	-	-	-	-	2	-	1	1	1
Male	58	1	-	-	-	-	-	1	-	1	-	-
Female	108	2	-	-	-	-	-	1	-	-	1	1
Black, both sexes	6	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Kentucky												
All races, both sexes	217	1	-	-	-	-	-	1	3	-	-	6
Male	81	1	-	-	-	-	-	-	1	-	-	4
Female	136	-	-	-	-	-	-	1	2	-	-	2
White, both sexes	203	-	-	-	-	-	-	1	3	-	-	6
Male	74	-	-	-	-	-	-	-	1	-	-	4
Female	129	-	-	-	-	-	-	1	2	-	-	2
Black, both sexes	12	1	-	-	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	213	3	1	-	1	1	1	1	-	-	4	3
Male	77	2	-	-	1	-	-	1	-	-	1	2
Female	136	1	1	-	-	1	1	-	-	-	3	1
White, both sexes	141	-	1	-	-	-	-	-	-	-	3	1
Male	45	-	-	-	-	-	-	-	-	-	-	1
Female	96	-	1	-	-	-	-	-	-	-	3	-
Black, both sexes	70	3	-	-	1	1	1	1	-	-	1	2
Male	32	2	-	-	1	-	-	1	-	-	1	1
Female	38	1	-	-	-	1	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Maine												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	275	2	-	-	-	-	-	-	1	1	1	6
Male	97	2	-	-	-	-	-	-	-	-	-	3
Female	178	-	-	-	-	-	-	-	1	1	1	3
White, both sexes	215	1	-	-	-	-	-	-	1	-	-	6
Male	71	1	-	-	-	-	-	-	-	-	-	3
Female	144	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	60	1	-	-	-	-	-	-	-	1	1	-
Male	26	1	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Massachusetts												
All races, both sexes	381	4	-	-	-	1	-	-	-	1	3	8
Male	150	4	-	-	-	1	-	-	-	1	1	7
Female	231	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	358	4	-	-	-	-	-	-	-	1	3	7
Male	137	4	-	-	-	-	-	-	-	1	1	6
Female	221	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	1	-	-	-	-	-	1
Male	5	-	-	-	-	1	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	513	4	-	-	1	-	2	1	-	-	5	9
Male	186	2	-	-	1	-	-	1	-	-	5	6
Female	327	2	-	-	-	-	2	-	-	-	-	3
White, both sexes	420	1	-	-	-	-	1	1	-	-	5	5
Male	146	-	-	-	-	-	-	1	-	-	5	3
Female	274	1	-	-	-	-	1	-	-	-	-	2
Black, both sexes	87	3	-	-	1	-	1	-	-	-	-	4
Male	39	2	-	-	1	-	-	-	-	-	-	3
Female	48	1	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Minnesota												
All races, both sexes	246	2	-	-	-	-	1	-	2	1	-	3
Male	94	-	-	-	-	-	1	-	2	1	-	1
Female	152	2	-	-	-	-	-	-	-	-	-	2
White, both sexes	236	1	-	-	-	-	-	-	2	1	-	3
Male	87	-	-	-	-	-	-	-	2	1	-	1
Female	149	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	141	3	2	-	-	-	-	-	1	3	2	2
Male	62	2	2	-	-	-	-	-	-	2	1	2
Female	79	1	-	-	-	-	-	-	1	1	1	-
White, both sexes	91	2	-	-	-	-	-	-	-	2	2	2
Male	38	1	-	-	-	-	-	-	-	2	1	2
Female	53	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	49	1	2	-	-	-	-	-	-	1	-	-
Male	24	1	2	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Missouri												
All races, both sexes	334	1	-	-	-	-	-	-	3	1	-	4
Male	121	1	-	-	-	-	-	-	2	1	-	2
Female	213	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	302	1	-	-	-	-	-	-	1	1	-	3
Male	109	1	-	-	-	-	-	-	1	1	-	1
Female	193	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	30	-	-	-	-	-	-	-	2	-	-	1
Male	12	-	-	-	-	-	-	-	1	-	-	1
Female	18	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	50	1	-	-	-	-	1	-	-	-	-	-
Male	28	-	-	-	-	-	1	-	-	-	-	-
Female	22	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	1	-	-	-	-	-
Male	27	-	-	-	-	-	1	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Nebraska												
All races, both sexes	116	-	-	-	2	-	1	-	-	-	2	-
Male	45	-	-	-	2	-	-	-	-	-	-	-
Female	71	-	-	-	-	-	1	-	-	-	2	-
White, both sexes	109	-	-	-	1	-	-	-	-	-	2	-
Male	41	-	-	-	1	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	6	-	-	-	1	-	1	-	-	-	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	75	-	-	-	-	-	-	1	-	1	3	-
Male	28	-	-	-	-	-	-	1	-	-	2	-
Female	47	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	68	-	-	-	-	-	-	1	-	1	3	-
Male	25	-	-	-	-	-	-	1	-	-	2	-
Female	43	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
New Hampshire												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	77	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	317	4	-	-	-	1	-	-	-	-	2	1
Male	115	2	-	-	-	1	-	-	-	-	1	-
Female	202	2	-	-	-	-	-	-	-	-	1	1
White, both sexes	285	1	-	-	-	-	-	-	-	-	2	-
Male	101	1	-	-	-	-	-	-	-	-	1	-
Female	184	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	30	3	-	-	-	1	-	-	-	-	-	1
Male	13	1	-	-	-	1	-	-	-	-	-	-
Female	17	2	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
New Mexico												
All races, both sexes	76	2	1	-	-	1	-	-	-	-	1	-
Male	35	2	1	-	-	1	-	-	-	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	2	1	-	-	1	-	-	-	-	1	-
Male	30	2	1	-	-	1	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	810	3	2	-	1	1	-	-	-	2	5	13
Male	264	2	2	-	1	-	-	-	-	2	2	6
Female	546	1	-	-	-	1	-	-	-	-	3	7
White, both sexes	710	2	1	-	-	1	-	-	-	-	4	10
Male	220	1	1	-	-	-	-	-	-	-	2	6
Female	490	1	-	-	-	1	-	-	-	-	2	4
Black, both sexes	93	1	1	-	1	-	-	-	-	2	1	3
Male	40	1	1	-	1	-	-	-	-	2	-	-
Female	53	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
North Carolina												
All races, both sexes	507	14	1	-	-	-	1	-	-	4	1	10
Male	175	11	-	-	-	-	-	-	-	3	1	7
Female	332	3	1	-	-	-	1	-	-	1	-	3
White, both sexes	394	6	1	-	-	-	1	-	-	2	-	7
Male	128	4	-	-	-	-	-	-	-	1	-	4
Female	266	2	1	-	-	-	1	-	-	1	-	3
Black, both sexes	111	8	-	-	-	-	-	-	-	2	1	3
Male	46	7	-	-	-	-	-	-	-	2	1	3
Female	65	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	46	-	-	-	-	1	-	-	-	-	-	1
Male	17	-	-	-	-	1	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	44	-	-	-	-	1	-	-	-	-	-	1
Male	16	-	-	-	-	1	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Ohio												
All races, both sexes	671	1	-	2	1	1	1	1	-	4	3	6
Male	253	-	-	1	1	1	1	1	-	2	1	3
Female	418	1	-	1	-	-	-	-	-	2	2	3
White, both sexes	600	1	-	2	1	-	1	-	-	4	2	6
Male	226	-	-	1	1	-	1	-	-	2	-	3
Female	374	1	-	1	-	-	-	-	-	2	2	3
Black, both sexes	71	-	-	-	-	1	-	1	-	-	1	-
Male	27	-	-	-	-	1	-	1	-	-	1	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	245	-	-	-	-	-	-	1	1	2	2	5
Male	75	-	-	-	-	-	-	-	-	1	-	3
Female	170	-	-	-	-	-	-	1	1	1	2	2
White, both sexes	219	-	-	-	-	-	-	1	1	2	2	4
Male	66	-	-	-	-	-	-	-	-	1	-	3
Female	153	-	-	-	-	-	-	1	1	1	2	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Oregon												
All races, both sexes	186	3	-	-	1	1	-	-	-	-	1	2
Male	78	2	-	-	1	-	-	-	-	-	-	-
Female	108	1	-	-	-	1	-	-	-	-	1	2
White, both sexes	181	3	-	-	1	1	-	-	-	-	1	2
Male	75	2	-	-	1	-	-	-	-	-	-	-
Female	106	1	-	-	-	1	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	818	3	1	-	-	-	2	-	1	1	8	7
Male	260	2	1	-	-	-	1	-	1	1	2	4
Female	558	1	-	-	-	-	1	-	-	-	6	3
White, both sexes	747	1	1	-	-	-	1	-	-	1	6	5
Male	235	1	1	-	-	-	1	-	-	1	2	3
Female	512	-	-	-	-	-	-	-	-	-	4	2
Black, both sexes	70	2	-	-	-	-	1	-	1	-	2	2
Male	25	1	-	-	-	-	-	-	1	-	-	1
Female	45	1	-	-	-	-	1	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Rhode Island												
All races, both sexes	61	-	-	-	-	-	-	1	-	-	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	1	-	-	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	216	2	-	1	-	-	1	1	-	-	-	5
Male	86	1	-	-	-	-	1	-	-	-	-	2
Female	130	1	-	1	-	-	-	1	-	-	-	3
White, both sexes	163	2	-	1	-	-	-	1	-	-	-	2
Male	62	1	-	-	-	-	-	-	-	-	-	-
Female	101	1	-	1	-	-	-	1	-	-	-	2
Black, both sexes	52	-	-	-	-	-	1	-	-	-	-	3
Male	23	-	-	-	-	-	1	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
South Dakota												
All races, both sexes	49	1	-	-	-	-	1	-	-	1	-	1
Male	20	1	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	45	1	-	-	-	-	1	-	-	-	-	-
Male	19	1	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	325	3	-	-	-	-	1	1	1	3	4	4
Male	105	-	-	-	-	-	-	1	1	1	2	2
Female	220	3	-	-	-	-	1	-	-	2	2	2
White, both sexes	288	1	-	-	-	-	1	1	1	2	3	3
Male	89	-	-	-	-	-	-	1	1	1	2	1
Female	199	1	-	-	-	-	1	-	-	1	1	2
Black, both sexes	36	1	-	-	-	-	-	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	20	1	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Texas												
All races, both sexes	953	9	3	-	-	1	1	4	2	2	7	14
Male	329	3	1	-	-	-	1	2	2	-	4	5
Female	624	6	2	-	-	1	-	2	-	2	3	9
White, both sexes	852	6	2	-	-	1	1	4	2	2	6	11
Male	287	2	1	-	-	-	1	2	2	-	3	3
Female	565	4	1	-	-	1	-	2	-	2	3	8
Black, both sexes	95	3	1	-	-	-	-	-	-	-	1	3
Male	41	1	-	-	-	-	-	-	-	-	1	2
Female	54	2	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	92	1	-	-	-	-	-	-	-	-	1	2
Male	39	1	-	-	-	-	-	-	-	-	1	2
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	90	1	-	-	-	-	-	-	-	-	1	2
Male	38	1	-	-	-	-	-	-	-	-	1	2
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Vermont												
All races, both sexes	31	-	-	-	-	1	-	-	-	1	-	-
Male	12	-	-	-	-	1	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	31	-	-	-	-	1	-	-	-	1	-	-
Male	12	-	-	-	-	1	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	352	2	-	-	-	-	-	-	2	2	2	3
Male	140	2	-	-	-	-	-	-	-	1	1	2
Female	212	-	-	-	-	-	-	-	2	1	1	1
White, both sexes	274	1	-	-	-	-	-	-	2	1	1	2
Male	108	1	-	-	-	-	-	-	-	-	1	1
Female	166	-	-	-	-	-	-	-	2	1	-	1
Black, both sexes	76	1	-	-	-	-	-	-	-	1	1	1
Male	32	1	-	-	-	-	-	-	-	1	-	1
Female	44	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Washington												
All races, both sexes	277	2	-	-	-	-	-	2	-	2	1	3
Male	87	2	-	-	-	-	-	1	-	-	1	3
Female	190	-	-	-	-	-	-	1	-	2	-	-
White, both sexes	258	-	-	-	-	-	-	2	-	2	1	2
Male	77	-	-	-	-	-	-	1	-	-	1	2
Female	181	-	-	-	-	-	-	1	-	2	-	-
Black, both sexes	10	1	-	-	-	-	-	-	-	-	-	1
Male	5	1	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	103	-	-	-	-	-	-	1	-	-	1	3
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	100	-	-	-	-	-	-	1	-	-	1	3
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	1	-	-	1	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Wisconsin												
All races, both sexes	302	3	1	-	-	-	-	1	-	-	3	2
Male	101	2	1	-	-	-	-	-	-	-	3	2
Female	201	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	281	3	-	-	-	-	-	1	-	-	3	1
Male	96	2	-	-	-	-	-	-	-	-	3	1
Female	185	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	14	-	1	-	-	-	-	-	-	-	-	1
Male	5	-	1	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	31	1	-	-	-	-	-	-	-	-	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	1	-	-	-	-	-	-	-	-	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
United States												
All races, both sexes	304	454	600	865	1,448	2,111	2,797	2,656	1,827	653	120	1
Male	154	215	287	377	605	835	951	791	467	129	25	1
Female	150	239	313	488	843	1,276	1,846	1,865	1,360	524	95	-
White, both sexes	230	361	484	704	1,256	1,853	2,525	2,434	1,670	587	105	1
Male	113	172	224	294	515	729	849	718	424	114	21	1
Female	117	189	260	410	741	1,124	1,676	1,716	1,246	473	84	-
Black, both sexes	64	82	101	126	171	217	215	184	128	57	14	-
Male	34	38	55	71	80	85	79	57	32	12	4	-
Female	30	44	46	55	91	132	136	127	96	45	10	-
American Indian 1/, both sexes	5	5	7	18	5	11	12	9	5	4	1	-
Male	3	1	5	5	3	5	4	2	4	2	-	-
Female	2	4	2	13	2	6	8	7	1	2	1	-
Asian or Pacific Islander 2/, both sexes	5	6	8	17	16	30	45	29	24	5	-	-
Male	4	4	3	7	7	16	19	14	7	1	-	-
Female	1	2	5	10	9	14	26	15	17	4	-	-
Alabama												
All races, both sexes	12	11	14	17	18	42	47	36	23	10	2	-
Male	7	5	5	5	6	16	18	6	4	2	1	-
Female	5	6	9	12	12	26	29	30	19	8	1	-
White, both sexes	7	9	11	15	14	33	39	29	17	5	1	-
Male	4	3	5	4	4	12	15	5	2	1	-	-
Female	3	6	6	11	10	21	24	24	15	4	1	-
Black, both sexes	5	1	3	2	4	9	8	7	6	5	1	-
Male	3	1	-	1	2	4	3	1	2	1	1	-
Female	2	-	3	1	2	5	5	6	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Alaska												
All races, both sexes	1	-	2	4	1	3	4	4	2	-	-	-
Male	-	-	2	-	-	1	-	2	-	-	-	-
Female	1	-	-	4	1	2	4	2	2	-	-	-
White, both sexes	1	-	1	1	1	2	3	2	1	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	1	-	-	1	1	1	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	3	-	1	-	2	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	3	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	9	7	16	21	22	33	43	43	35	8	4	-
Male	8	4	6	11	12	15	17	11	9	1	-	-
Female	1	3	10	10	10	18	26	32	26	7	4	-
White, both sexes	8	6	15	17	20	28	40	43	34	7	4	-
Male	7	3	5	9	11	14	15	11	9	-	-	-
Female	1	3	10	8	9	14	25	32	25	7	4	-
Black, both sexes	-	-	1	1	1	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	1	1	-	2	-	4	2	-	-	1	-	-
Male	1	1	-	1	-	1	1	-	-	1	-	-
Female	-	-	-	1	-	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Arkansas												
All races, both sexes	4	8	5	13	20	24	38	36	26	4	3	-
Male	3	3	2	10	10	9	11	10	3	-	1	-
Female	1	5	3	3	10	15	27	26	23	4	2	-
White, both sexes	4	6	4	9	19	21	36	34	24	3	1	-
Male	3	3	2	7	10	6	10	10	2	-	-	-
Female	1	3	2	2	9	15	26	24	22	3	1	-
Black, both sexes	-	2	1	4	1	3	2	2	2	1	2	-
Male	-	-	-	3	-	3	1	-	1	-	1	-
Female	-	2	1	1	1	-	1	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	31	50	56	93	124	207	286	256	142	54	9	1
Male	19	21	29	34	54	81	96	86	41	9	3	1
Female	12	29	27	59	70	126	190	170	101	45	6	-
White, both sexes	23	39	42	75	103	174	242	235	122	50	8	1
Male	15	16	19	27	44	61	75	80	34	8	2	1
Female	8	23	23	48	59	113	167	155	88	42	6	-
Black, both sexes	5	8	9	7	11	14	21	7	6	2	1	-
Male	2	3	7	3	6	11	9	1	2	1	1	-
Female	3	5	2	4	5	3	12	6	4	1	-	-
American Indian 1/, both sexes	1	1	1	1	1	-	2	2	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	1	-	1	1	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	4	10	9	19	21	12	14	2	-	-
Male	2	2	2	4	4	9	11	5	5	-	-	-
Female	-	-	2	6	5	10	10	7	9	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Colorado												
All races, both sexes	6	9	9	15	17	24	22	39	12	12	2	-
Male	4	3	3	9	11	6	5	13	2	3	-	-
Female	2	6	6	6	6	18	17	26	10	9	2	-
White, both sexes	6	9	7	13	17	24	22	39	12	12	2	-
Male	4	3	3	8	11	6	5	13	2	3	-	-
Female	2	6	4	5	6	18	17	26	10	9	2	-
Black, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	3	6	5	9	26	33	49	41	22	2	-
Male	-	2	2	2	2	8	14	16	10	2	1	-
Female	3	1	4	3	7	18	19	33	31	20	1	-
White, both sexes	2	3	5	5	9	24	31	48	39	22	1	-
Male	-	2	2	2	2	8	14	16	10	2	1	-
Female	2	1	3	3	7	16	17	32	29	20	-	-
Black, both sexes	1	-	1	-	-	1	2	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	2	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Delaware												
All races, both sexes	-	2	2	-	5	5	6	4	5	1	1	-
Male	-	1	1	-	1	3	4	-	2	-	-	-
Female	-	1	1	-	4	2	2	4	3	1	1	-
White, both sexes	-	1	2	-	5	5	6	4	4	1	1	-
Male	-	-	1	-	1	3	4	-	1	-	-	-
Female	-	1	1	-	4	2	2	4	3	1	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	2	2	2	5	1	9	4	4	1	-	-
Male	1	-	-	2	3	-	1	4	1	-	-	-
Female	-	2	2	-	2	1	8	-	3	1	-	-
White, both sexes	-	-	-	1	-	-	1	2	2	1	-	-
Male	-	-	-	1	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	2	2	1	5	1	7	2	2	-	-	-
Male	-	-	-	1	3	-	-	2	-	-	-	-
Female	-	2	2	-	2	1	7	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Florida												
All races, both sexes	17	30	37	51	82	129	150	162	130	43	9	-
Male	10	17	17	22	28	52	62	49	30	10	3	-
Female	7	13	20	29	54	77	88	113	100	33	6	-
White, both sexes	13	20	34	47	70	121	143	148	125	38	8	-
Male	8	11	16	19	24	50	58	44	30	10	3	-
Female	5	9	18	28	46	71	85	104	95	28	5	-
Black, both sexes	4	10	2	3	11	8	6	14	5	4	1	-
Male	2	6	1	3	4	2	4	5	-	-	-	-
Female	2	4	1	-	7	6	2	9	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	1	-	-
Georgia												
All races, both sexes	14	17	24	27	42	65	60	65	31	11	4	-
Male	6	7	10	13	21	23	14	19	8	1	1	-
Female	8	10	14	14	21	42	46	46	23	10	3	-
White, both sexes	9	15	15	20	34	52	48	51	25	9	3	-
Male	4	6	4	9	17	18	12	16	6	1	1	-
Female	5	9	11	11	17	34	36	35	19	8	2	-
Black, both sexes	4	2	9	7	8	13	12	14	6	2	1	-
Male	2	1	6	4	4	5	2	3	2	-	-	-
Female	2	1	3	3	4	8	10	11	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Hawaii												
All races, both sexes	1	-	5	3	2	5	11	13	3	1	-	-
Male	1	-	3	3	2	1	5	5	-	1	-	-
Female	-	-	2	-	-	4	6	8	3	-	-	-
White, both sexes	-	-	3	2	1	1	5	2	1	-	-	-
Male	-	-	2	2	1	-	2	-	-	-	-	-
Female	-	-	1	-	-	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	1	4	6	11	2	1	-	-
Male	1	-	1	1	1	1	3	5	-	1	-	-
Female	-	-	1	-	-	3	3	6	2	-	-	-
Idaho												
All races, both sexes	-	3	2	2	7	15	15	13	13	3	-	-
Male	-	1	1	-	3	7	6	5	5	1	-	-
Female	-	2	1	2	4	8	9	8	8	2	-	-
White, both sexes	-	3	2	2	7	15	15	13	13	3	-	-
Male	-	1	1	-	3	7	6	5	5	1	-	-
Female	-	2	1	2	4	8	9	8	8	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Illinois												
All races, both sexes	15	16	24	38	54	91	141	134	84	22	4	-
Male	6	5	13	21	24	38	43	41	18	4	2	-
Female	9	11	11	17	30	53	98	93	66	18	2	-
White, both sexes	9	13	19	34	51	82	127	124	74	20	4	-
Male	2	4	11	19	23	35	37	37	14	4	2	-
Female	7	9	8	15	28	47	90	87	60	16	2	-
Black, both sexes	6	3	5	4	2	7	12	9	10	2	-	-
Male	4	1	2	2	1	1	5	3	4	-	-	-
Female	2	2	3	2	1	6	7	6	6	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	-	2	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Indiana												
All races, both sexes	8	8	20	25	39	51	59	56	47	16	2	-
Male	3	4	5	7	14	20	17	16	14	5	-	-
Female	5	4	15	18	25	31	42	40	33	11	2	-
White, both sexes	7	5	19	24	35	47	55	55	44	16	2	-
Male	2	3	4	6	14	20	14	16	13	5	-	-
Female	5	2	15	18	21	27	41	39	31	11	2	-
Black, both sexes	1	3	1	1	4	4	4	1	3	-	-	-
Male	1	1	1	1	-	-	3	-	1	-	-	-
Female	-	2	-	-	4	4	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Iowa												
All races, both sexes	2	5	8	11	18	28	26	28	28	10	3	-
Male	1	2	2	4	6	13	9	6	8	2	-	-
Female	1	3	6	7	12	15	17	22	20	8	3	-
White, both sexes	2	4	8	11	17	27	26	28	28	10	3	-
Male	1	2	2	4	6	13	9	6	8	2	-	-
Female	1	2	6	7	11	14	17	22	20	8	3	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	3	2	8	17	22	30	30	32	11	2	-
Male	5	1	-	5	7	12	10	8	8	2	1	-
Female	4	2	2	3	10	10	20	22	24	9	1	-
White, both sexes	8	3	2	6	17	22	26	30	32	10	2	-
Male	4	1	-	4	7	12	9	8	8	1	1	-
Female	4	2	2	2	10	10	17	22	24	9	1	-
Black, both sexes	-	-	-	1	-	-	3	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Kentucky												
All races, both sexes	4	12	7	18	26	31	41	34	25	8	-	-
Male	1	6	3	8	14	13	12	12	5	1	-	-
Female	3	6	4	10	12	18	29	22	20	7	-	-
White, both sexes	4	11	7	17	24	28	39	31	24	8	-	-
Male	1	5	3	8	12	12	11	11	5	1	-	-
Female	3	6	4	9	12	16	28	20	19	7	-	-
Black, both sexes	-	1	-	1	2	3	1	2	1	-	-	-
Male	-	1	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	1	-	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	10	7	15	26	30	36	39	19	10	1	-
Male	2	7	2	4	11	10	15	12	4	2	1	-
Female	3	3	5	11	15	20	21	27	15	8	-	-
White, both sexes	4	6	5	8	15	20	27	29	14	7	1	-
Male	2	4	1	-	4	8	12	8	3	1	1	-
Female	2	2	4	8	11	12	15	21	11	6	-	-
Black, both sexes	1	4	2	7	11	10	8	9	5	3	-	-
Male	-	3	1	4	7	2	3	4	1	1	-	-
Female	1	1	1	3	4	8	5	5	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Maine												
All races, both sexes	2	1	2	3	5	7	19	17	21	8	3	-
Male	1	-	-	2	1	2	7	8	6	3	2	-
Female	1	1	2	1	4	5	12	9	15	5	1	-
White, both sexes	2	1	2	3	5	7	19	17	21	8	3	-
Male	1	-	-	2	1	2	7	8	6	3	2	-
Female	1	1	2	1	4	5	12	9	15	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	12	7	20	28	43	52	47	33	15	-	-
Male	4	9	5	8	6	19	18	16	4	3	-	-
Female	3	3	2	12	22	24	34	31	29	12	-	-
White, both sexes	4	6	3	12	24	30	46	41	27	14	-	-
Male	2	6	2	3	5	15	16	13	3	2	-	-
Female	2	-	1	9	19	15	30	28	24	12	-	-
Black, both sexes	3	6	4	8	4	13	6	6	6	1	-	-
Male	2	3	3	5	1	4	2	3	1	1	-	-
Female	1	3	1	3	3	9	4	3	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Massachusetts												
All races, both sexes	9	8	9	11	40	49	78	78	55	22	5	-
Male	3	2	3	6	22	20	26	32	18	4	-	-
Female	6	6	6	5	18	29	52	46	37	18	5	-
White, both sexes	9	7	7	11	34	45	76	74	54	21	5	-
Male	3	2	2	6	19	17	25	31	17	3	-	-
Female	6	5	5	5	15	28	51	43	37	18	5	-
Black, both sexes	-	1	2	-	6	3	1	4	-	1	-	-
Male	-	-	1	-	3	2	-	1	-	1	-	-
Female	-	1	1	-	3	1	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	14	15	25	29	53	71	102	84	67	27	4	-
Male	7	5	11	15	21	29	36	24	17	5	1	-
Female	7	10	14	14	32	42	66	60	50	22	3	-
White, both sexes	10	12	20	18	41	57	88	76	57	24	4	-
Male	6	5	7	9	15	25	30	21	14	4	1	-
Female	4	7	13	9	26	32	58	55	43	20	3	-
Black, both sexes	4	2	5	10	11	14	11	8	10	3	-	-
Male	1	-	4	6	5	4	6	3	3	1	-	-
Female	3	2	1	4	6	10	5	5	7	2	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Minnesota												
All races, both sexes	4	7	9	15	25	28	38	49	43	16	3	-
Male	2	4	3	5	13	14	19	13	11	3	2	-
Female	2	3	6	10	12	14	19	36	32	13	1	-
White, both sexes	3	6	8	15	24	27	37	48	42	16	3	-
Male	2	3	2	5	12	13	18	13	10	3	2	-
Female	1	3	6	10	12	14	19	35	32	13	1	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	10	5	12	19	23	20	28	4	3	1	-
Male	3	6	2	6	12	9	6	7	2	-	-	-
Female	-	4	3	6	7	14	14	21	2	3	1	-
White, both sexes	1	7	4	6	12	16	13	20	3	1	-	-
Male	1	4	2	3	8	5	4	4	1	-	-	-
Female	-	3	2	3	4	11	9	16	2	1	-	-
Black, both sexes	2	3	1	6	7	7	7	8	1	2	1	-
Male	2	2	-	3	4	4	2	3	1	-	-	-
Female	-	1	1	3	3	3	5	5	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Missouri												
All races, both sexes	7	14	17	21	35	51	68	58	41	10	3	-
Male	5	9	6	5	15	23	24	18	8	2	-	-
Female	2	5	11	16	20	28	44	40	33	8	3	-
White, both sexes	4	10	13	17	32	48	66	56	37	10	3	-
Male	3	6	4	4	14	22	24	18	8	2	-	-
Female	1	4	9	13	18	26	42	38	29	8	3	-
Black, both sexes	3	4	3	3	3	3	2	2	4	-	-	-
Male	2	3	2	1	1	1	-	-	-	-	-	-
Female	1	1	1	2	2	2	2	2	4	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	3	4	14	9	8	7	1	-	-
Male	1	-	1	2	1	9	4	5	3	1	-	-
Female	-	-	-	1	3	5	5	3	4	-	-	-
White, both sexes	1	-	1	2	4	13	9	8	7	1	-	-
Male	1	-	1	2	1	8	4	5	3	1	-	-
Female	-	-	-	-	3	5	5	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Nebraska												
All races, both sexes	1	3	3	9	11	13	24	20	17	9	1	-
Male	1	1	1	5	7	6	10	5	4	3	-	-
Female	-	2	2	4	4	7	14	15	13	6	1	-
White, both sexes	1	2	3	9	10	12	24	20	16	8	1	-
Male	1	1	1	5	6	5	10	5	3	3	-	-
Female	-	1	2	4	4	7	14	15	13	5	1	-
Black, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	3	2	4	8	11	16	11	8	2	-	-
Male	2	-	-	2	2	5	6	7	1	-	-	-
Female	3	3	2	2	6	6	10	4	7	2	-	-
White, both sexes	5	2	2	4	6	9	14	11	8	2	-	-
Male	2	-	-	2	2	4	4	7	1	-	-	-
Female	3	2	2	2	4	5	10	4	7	2	-	-
Black, both sexes	-	1	-	-	2	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
New Hampshire												
All races, both sexes	2	1	2	5	6	14	22	14	5	6	-	-
Male	2	1	1	2	-	7	9	6	2	3	-	-
Female	-	-	1	3	6	7	13	8	3	3	-	-
White, both sexes	2	1	2	5	6	14	22	14	5	6	-	-
Male	2	1	1	2	-	7	9	6	2	3	-	-
Female	-	-	1	3	6	7	13	8	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8	9	10	17	36	37	67	66	45	13	1	-
Male	1	2	7	10	15	17	30	12	15	2	-	-
Female	7	7	3	7	21	20	37	54	30	11	1	-
White, both sexes	8	9	9	14	30	37	60	60	42	13	-	-
Male	1	2	6	8	13	17	26	10	14	2	-	-
Female	7	7	3	6	17	20	34	50	28	11	-	-
Black, both sexes	-	-	1	2	5	-	7	6	3	-	1	-
Male	-	-	1	1	2	-	4	2	1	-	-	-
Female	-	-	-	1	3	-	3	4	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
New Mexico												
All races, both sexes	-	2	1	6	6	11	17	10	10	8	-	-
Male	-	1	1	5	2	3	9	4	2	3	-	-
Female	-	1	-	1	4	8	8	6	8	5	-	-
White, both sexes	-	2	1	5	5	11	15	10	9	6	-	-
Male	-	1	1	5	1	3	7	4	1	2	-	-
Female	-	1	-	-	4	8	8	6	8	4	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	1	1	-	-
Male	-	-	-	-	1	-	1	-	1	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
New York												
All races, both sexes	14	23	37	40	81	112	150	171	96	51	8	-
Male	7	14	27	22	21	35	44	40	31	8	-	-
Female	7	9	10	18	60	77	106	131	65	43	8	-
White, both sexes	8	19	29	34	71	90	137	161	88	48	7	-
Male	2	12	21	17	16	26	41	38	30	7	-	-
Female	6	7	8	17	55	64	96	123	58	41	7	-
Black, both sexes	6	4	8	6	10	19	13	8	6	3	1	-
Male	5	2	6	5	5	6	3	1	1	1	-	-
Female	1	2	2	1	5	13	10	7	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	2	2	-	-	-
Male	-	-	-	-	-	3	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
North Carolina												
All races, both sexes	12	20	27	28	73	64	103	84	50	13	2	-
Male	8	8	11	12	26	31	29	18	10	-	-	-
Female	4	12	16	16	47	33	74	66	40	13	2	-
White, both sexes	10	15	17	18	60	54	82	71	39	9	2	-
Male	8	6	7	6	21	24	24	15	8	-	-	-
Female	2	9	10	12	39	30	58	56	31	9	2	-
Black, both sexes	2	5	10	10	12	10	21	13	10	4	-	-
Male	-	2	4	6	5	7	5	3	1	-	-	-
Female	2	3	6	4	7	3	16	10	9	4	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	1	-	4	7	8	6	13	3	-	-
Male	1	-	1	-	2	3	3	-	5	1	-	-
Female	-	1	-	-	2	4	5	6	8	2	-	-
White, both sexes	1	-	1	-	4	7	7	6	13	3	-	-
Male	1	-	1	-	2	3	2	-	5	1	-	-
Female	-	-	-	-	2	4	5	6	8	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Ohio												
All races, both sexes	11	15	29	46	71	106	129	126	85	31	2	-
Male	5	6	15	16	36	39	46	42	26	10	1	-
Female	6	9	14	30	35	67	83	84	59	21	1	-
White, both sexes	10	12	25	39	65	90	120	119	76	25	2	-
Male	5	5	12	15	31	34	42	40	24	9	1	-
Female	5	7	13	24	34	56	78	79	52	16	1	-
Black, both sexes	1	3	4	7	6	16	9	7	9	6	-	-
Male	-	1	3	1	5	5	4	2	2	1	-	-
Female	1	2	1	6	1	11	5	5	7	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	7	9	18	24	34	47	33	43	14	1	-
Male	1	2	5	8	10	11	10	11	11	2	-	-
Female	3	5	4	10	14	23	37	22	32	12	1	-
White, both sexes	3	7	8	12	24	33	42	29	39	11	1	-
Male	1	2	5	5	10	11	10	9	8	1	-	-
Female	2	5	3	7	14	22	32	20	31	10	1	-
Black, both sexes	-	-	1	1	-	-	3	3	3	2	-	-
Male	-	-	-	-	-	-	-	1	2	1	-	-
Female	-	-	1	1	-	-	3	2	1	1	-	-
American Indian 1/, both sexes	1	-	-	5	-	1	2	1	1	1	-	-
Male	-	-	-	3	-	-	-	1	1	-	-	-
Female	1	-	-	2	-	1	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Oregon												
All races, both sexes	5	7	9	13	19	26	36	40	19	3	1	-
Male	2	6	7	5	6	13	11	19	5	1	-	-
Female	3	1	2	8	13	13	25	21	14	2	1	-
White, both sexes	5	7	8	13	19	23	36	39	19	3	1	-
Male	2	6	6	5	6	12	11	18	5	1	-	-
Female	3	1	2	8	13	11	25	21	14	2	1	-
Black, both sexes	-	-	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	22	22	33	74	130	181	184	109	29	7	-
Male	1	11	13	13	27	48	61	42	27	4	1	-
Female	3	11	9	20	47	82	120	142	82	25	6	-
White, both sexes	3	18	16	29	68	118	171	173	101	28	7	-
Male	1	8	10	12	26	44	58	37	25	4	1	-
Female	2	10	6	17	42	74	113	136	76	24	6	-
Black, both sexes	1	4	6	4	6	12	9	11	8	1	-	-
Male	-	3	3	1	1	4	3	5	2	-	-	-
Female	1	1	3	3	5	8	6	6	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Rhode Island												
All races, both sexes	-	2	4	-	3	7	17	11	11	4	-	-
Male	-	1	2	-	-	4	5	3	3	1	-	-
Female	-	1	2	-	3	3	12	8	8	3	-	-
White, both sexes	-	2	4	-	2	7	17	11	11	4	-	-
Male	-	1	2	-	-	4	5	3	3	1	-	-
Female	-	1	2	-	2	3	12	8	8	3	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	6	19	13	28	37	46	31	13	9	1	-
Male	2	4	8	7	12	15	19	10	3	2	-	-
Female	1	2	11	6	16	22	27	21	10	7	1	-
White, both sexes	2	5	14	6	21	30	38	25	9	6	1	-
Male	1	3	6	1	9	12	16	10	1	2	-	-
Female	1	2	8	5	12	18	22	15	8	4	1	-
Black, both sexes	1	1	5	6	7	7	8	6	4	3	-	-
Male	1	1	2	5	3	3	3	-	2	-	-	-
Female	-	-	3	1	4	4	5	6	2	3	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
South Dakota												
All races, both sexes	-	1	1	2	2	10	5	11	8	5	-	-
Male	-	-	1	1	-	6	2	5	3	-	-	-
Female	-	1	-	1	2	4	3	6	5	5	-	-
White, both sexes	-	1	1	2	2	10	4	10	8	5	-	-
Male	-	-	1	1	-	6	2	5	3	-	-	-
Female	-	1	-	1	2	4	2	5	5	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	14	10	21	36	52	65	57	37	8	2	-
Male	2	10	6	9	18	12	24	13	3	1	-	-
Female	4	4	4	12	18	40	41	44	34	7	2	-
White, both sexes	5	11	8	17	30	48	63	50	35	7	2	-
Male	1	8	5	6	14	11	22	12	3	1	-	-
Female	4	3	3	11	16	37	41	38	32	6	2	-
Black, both sexes	1	3	2	4	6	4	2	7	2	1	-	-
Male	1	2	1	3	4	1	2	1	-	-	-	-
Female	-	1	1	1	2	3	-	6	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Texas												
All races, both sexes	19	31	37	64	99	159	170	158	120	40	13	-
Male	7	10	17	29	45	56	64	45	31	6	1	-
Female	12	21	20	35	54	103	106	113	89	34	12	-
White, both sexes	15	26	34	54	90	141	154	143	113	35	12	-
Male	5	10	15	24	38	49	57	39	30	5	1	-
Female	10	16	19	30	52	92	97	104	83	30	11	-
Black, both sexes	4	5	3	10	8	18	14	15	5	5	-	-
Male	2	-	2	5	6	7	7	6	1	1	-	-
Female	2	5	1	5	2	11	7	9	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	2	-	-	-
Utah												
All races, both sexes	1	1	9	8	16	12	16	16	8	-	1	-
Male	-	1	4	4	8	5	5	5	2	-	1	-
Female	1	-	5	4	8	7	11	11	6	-	-	-
White, both sexes	1	1	9	7	16	12	16	15	8	-	1	-
Male	-	1	4	4	8	5	5	4	2	-	1	-
Female	1	-	5	3	8	7	11	11	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Vermont												
All races, both sexes	-	1	1	1	3	1	7	1	7	6	1	-
Male	-	1	1	-	2	-	2	-	2	2	1	-
Female	-	-	-	1	1	1	5	1	5	4	-	-
White, both sexes	-	1	1	1	3	1	7	1	7	6	1	-
Male	-	1	1	-	2	-	2	-	2	2	1	-
Female	-	-	-	1	1	1	5	1	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	10	17	15	39	51	73	64	48	13	3	-
Male	4	8	10	8	16	23	21	24	16	3	1	-
Female	4	2	7	7	23	28	52	40	32	10	2	-
White, both sexes	2	7	11	9	26	41	63	56	42	9	1	-
Male	2	7	6	4	11	18	19	21	15	2	-	-
Female	-	-	5	5	15	23	44	35	27	7	1	-
Black, both sexes	6	3	6	6	13	10	9	7	6	4	2	-
Male	2	1	4	4	5	5	2	3	1	1	1	-
Female	4	2	2	2	8	5	7	4	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Washington												
All races, both sexes	7	6	12	19	19	42	62	54	36	8	2	-
Male	3	2	6	3	10	18	15	15	7	1	-	-
Female	4	4	6	16	9	24	47	39	29	7	2	-
White, both sexes	4	6	10	18	18	40	59	52	34	8	2	-
Male	-	2	5	3	10	17	15	13	7	1	-	-
Female	4	4	5	15	8	23	44	39	27	7	2	-
Black, both sexes	2	-	-	1	1	-	1	1	2	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	1	-	2	-	-	-
American Indian 1/, both sexes	1	-	2	-	-	1	1	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
West Virginia												
All races, both sexes	1	2	4	6	14	20	24	16	7	1	3	-
Male	-	1	2	1	5	12	7	4	1	-	-	-
Female	1	1	2	5	9	8	17	12	6	1	3	-
White, both sexes	1	2	4	5	14	19	24	16	7	1	2	-
Male	-	1	2	1	5	11	7	4	1	-	-	-
Female	1	1	2	4	9	8	17	12	6	1	2	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Wisconsin												
All races, both sexes	2	2	11	13	30	32	65	52	54	27	4	-
Male	1	1	4	6	12	12	18	16	14	9	-	-
Female	1	1	7	7	18	20	47	36	40	18	4	-
White, both sexes	2	1	8	10	28	32	61	49	53	26	3	-
Male	1	1	4	5	12	12	17	15	14	9	-	-
Female	1	-	4	5	16	20	44	34	39	17	3	-
Black, both sexes	-	-	2	1	2	-	2	3	1	-	1	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	2	-	2	-	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	2	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	2	-	2	3	5	4	6	5	1	-	-
Male	1	-	-	-	3	1	2	1	2	-	-	-
Female	-	2	-	2	-	4	2	5	3	1	-	-
White, both sexes	1	2	-	2	3	5	4	6	5	1	-	-
Male	1	-	-	-	3	1	2	1	2	-	-	-
Female	-	2	-	2	-	4	2	5	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
United States												
All races, both sexes	3,152	1	-	-	-	-	1	2	1	2	13	39
Male	999	1	-	-	-	-	1	-	1	2	11	21
Female	2,153	-	-	-	-	-	-	2	-	-	2	18
White, both sexes	2,872	1	-	-	-	-	1	1	1	1	9	31
Male	903	1	-	-	-	-	1	-	1	1	7	17
Female	1,969	-	-	-	-	-	-	1	-	-	2	14
Black, both sexes	244	-	-	-	-	-	-	1	-	1	4	8
Male	79	-	-	-	-	-	-	-	-	1	4	4
Female	165	-	-	-	-	-	-	1	-	-	-	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	60	-	-	-	-	-	-	-	-	1	1	-
Male	25	-	-	-	-	-	-	-	-	1	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	1	-	-
Male	23	-	-	-	-	-	-	-	-	1	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Arkansas												
All races, both sexes	35	-	-	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	364	-	-	-	-	-	-	1	-	-	1	6
Male	131	-	-	-	-	-	-	-	-	-	-	3
Female	233	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	327	-	-	-	-	-	-	1	-	-	1	5
Male	115	-	-	-	-	-	-	-	-	-	-	3
Female	212	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Colorado												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Florida												
All races, both sexes	220	-	-	-	-	-	-	-	-	1	3	3
Male	81	-	-	-	-	-	-	-	-	1	3	2
Female	139	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	207	-	-	-	-	-	-	-	-	-	3	2
Male	78	-	-	-	-	-	-	-	-	-	3	2
Female	129	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Hawaii												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Illinois												
All races, both sexes	148	-	-	-	-	-	-	-	-	-	-	1
Male	46	-	-	-	-	-	-	-	-	-	-	1
Female	102	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	138	-	-	-	-	-	-	-	-	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	94	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	72	-	-	-	-	-	-	-	1	-	-	1
Male	26	-	-	-	-	-	-	-	1	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	1	-	-	1
Male	25	-	-	-	-	-	-	-	1	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Iowa												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Kentucky												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Maine												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	61	-	-	-	-	-	1	-	-	-	-	-
Male	15	-	-	-	-	-	1	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	1	-	-	-	-	-
Male	14	-	-	-	-	-	1	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Massachusetts												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	77	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	110	-	-	-	-	-	-	-	-	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	81	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	99	-	-	-	-	-	-	-	-	-	-	2
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Minnesota												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Missouri												
All races, both sexes	66	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	64	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Nebraska												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
New Hampshire												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	1
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
New Mexico												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	217	-	-	-	-	-	-	-	-	-	2	4
Male	76	-	-	-	-	-	-	-	-	-	2	4
Female	141	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	182	-	-	-	-	-	-	-	-	-	-	2
Male	61	-	-	-	-	-	-	-	-	-	-	2
Female	121	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	-	-	2	2
Male	12	-	-	-	-	-	-	-	-	-	2	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
North Carolina												
All races, both sexes	96	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	65	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Ohio												
All races, both sexes	151	-	-	-	-	-	-	-	-	-	1	-
Male	52	-	-	-	-	-	-	-	-	-	1	-
Female	99	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	141	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	95	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Oregon												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	46	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	145	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	112	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	140	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	108	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Rhode Island												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
South Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	76	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Texas												
All races, both sexes	171	1	-	-	-	-	-	-	-	-	2	4
Male	58	1	-	-	-	-	-	-	-	-	1	4
Female	113	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	147	1	-	-	-	-	-	-	-	-	2	2
Male	49	1	-	-	-	-	-	-	-	-	1	2
Female	98	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Washington												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
Wisconsin												
All races, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
United States												
All races, both sexes	54	102	114	165	264	456	621	652	476	171	18	-
Male	31	52	44	67	108	154	199	167	118	21	1	-
Female	23	50	70	98	156	302	422	485	358	150	17	-
White, both sexes	45	79	98	142	237	413	572	617	449	159	16	-
Male	26	43	39	53	95	142	181	162	113	20	1	-
Female	19	36	59	89	142	271	391	455	336	139	15	-
Black, both sexes	8	22	15	19	20	38	41	29	24	12	2	-
Male	5	8	5	12	8	10	13	4	4	1	-	-
Female	3	14	10	7	12	28	28	25	20	11	2	-
American Indian 1/, both sexes	-	-	-	1	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	6	4	7	6	2	-	-	-
Male	-	1	-	2	5	1	4	1	1	-	-	-
Female	1	-	1	1	1	3	3	5	1	-	-	-
Alabama												
All races, both sexes	-	-	1	2	4	10	11	15	9	2	-	-
Male	-	-	1	1	1	1	2	4	3	-	-	-
Female	-	-	-	1	3	9	9	11	6	2	-	-
White, both sexes	-	-	1	2	3	9	9	14	9	2	-	-
Male	-	-	1	1	1	1	2	4	3	-	-	-
Female	-	-	-	1	2	8	7	10	6	2	-	-
Black, both sexes	-	-	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	4	-	5	2	8	12	12	10	3	-	-
Male	1	2	-	2	2	1	5	4	6	-	-	-
Female	1	2	-	3	-	7	7	8	4	3	-	-
White, both sexes	2	4	-	5	2	6	12	12	10	3	-	-
Male	1	2	-	2	2	-	5	4	6	-	-	-
Female	1	2	-	3	-	6	7	8	4	3	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Arkansas												
All races, both sexes	1	1	1	2	7	2	1	8	8	3	-	-
Male	1	-	1	2	6	-	-	2	4	-	-	-
Female	-	1	-	-	1	2	1	6	4	3	-	-
White, both sexes	1	1	1	2	7	2	1	7	7	3	-	-
Male	1	-	1	2	6	-	-	2	4	-	-	-
Female	-	1	-	-	1	2	1	5	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9	14	10	12	38	53	73	80	49	15	3	-
Male	4	9	4	5	14	22	25	25	17	3	-	-
Female	5	5	6	7	24	31	48	55	32	12	3	-
White, both sexes	9	14	9	11	31	48	63	70	47	15	3	-
Male	4	9	4	4	10	20	21	21	16	3	-	-
Female	5	5	5	7	21	28	42	49	31	12	3	-
Black, both sexes	-	-	1	1	1	3	6	5	1	-	-	-
Male	-	-	-	1	-	1	2	3	-	-	-	-
Female	-	-	1	-	1	2	4	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	5	2	3	5	1	-	-	-
Male	-	-	-	-	4	1	1	1	1	-	-	-
Female	-	-	-	-	1	1	2	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Colorado												
All races, both sexes	-	2	4	2	3	4	8	5	5	3	-	-
Male	-	-	-	-	-	2	4	2	1	-	-	-
Female	-	2	4	2	3	2	4	3	4	3	-	-
White, both sexes	-	2	4	2	3	4	8	5	5	3	-	-
Male	-	-	-	-	-	2	4	2	1	-	-	-
Female	-	2	4	2	3	2	4	3	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	3	-	4	7	6	8	9	4	-	-
Male	-	1	2	-	-	3	4	4	-	1	-	-
Female	-	-	1	-	4	4	2	4	9	3	-	-
White, both sexes	-	1	3	-	3	7	6	8	9	4	-	-
Male	-	1	2	-	-	3	4	4	-	1	-	-
Female	-	-	1	-	3	4	2	4	9	3	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Delaware												
All races, both sexes	-	1	-	1	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	2	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	1	-	-	-	1	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	1	-	-	-	1	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Florida												
All races, both sexes	2	8	14	12	20	30	48	45	24	10	-	-
Male	2	7	5	7	9	9	16	13	7	-	-	-
Female	-	1	9	5	11	21	32	32	17	10	-	-
White, both sexes	2	8	12	12	16	29	46	45	23	9	-	-
Male	2	7	5	7	8	9	16	13	6	-	-	-
Female	-	1	7	5	8	20	30	32	17	9	-	-
Black, both sexes	-	-	2	-	4	1	2	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	2	-	3	1	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	4	2	2	4	10	13	11	7	4	-	-
Male	1	2	1	2	1	2	6	4	1	1	-	-
Female	1	2	1	-	3	8	7	7	6	3	-	-
White, both sexes	1	-	1	2	3	8	10	10	7	2	-	-
Male	1	-	-	2	1	2	5	4	1	-	-	-
Female	-	-	1	-	2	6	5	6	6	2	-	-
Black, both sexes	1	4	1	-	1	2	3	1	-	2	-	-
Male	-	2	1	-	-	-	1	-	-	1	-	-
Female	1	2	-	-	1	2	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Hawaii												
All races, both sexes	1	-	1	-	1	1	1	-	1	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
Idaho												
All races, both sexes	-	-	1	-	3	3	4	5	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	2	3	4	4	-	-	-	-
White, both sexes	-	-	1	-	3	3	4	5	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	2	3	4	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Illinois												
All races, both sexes	1	5	4	3	14	18	29	35	24	12	2	-
Male	1	2	2	2	9	8	6	8	6	1	-	-
Female	-	3	2	1	5	10	23	27	18	11	2	-
White, both sexes	1	4	4	2	13	17	26	35	22	11	2	-
Male	1	2	2	2	8	8	5	8	6	1	-	-
Female	-	2	2	-	5	9	21	27	16	10	2	-
Black, both sexes	-	1	-	1	1	1	3	-	2	1	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	2	3	8	8	16	21	5	3	1	-
Male	2	1	-	2	3	2	2	8	3	1	-	-
Female	-	-	2	1	5	6	14	13	2	2	1	-
White, both sexes	1	1	2	3	8	8	16	21	5	3	1	-
Male	1	1	-	2	3	2	2	8	3	1	-	-
Female	-	-	2	1	5	6	14	13	2	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Iowa												
All races, both sexes	-	-	1	3	3	5	8	13	9	2	-	-
Male	-	-	-	2	1	4	4	2	3	-	-	-
Female	-	-	1	1	2	1	4	11	6	2	-	-
White, both sexes	-	-	1	2	3	5	8	13	9	1	-	-
Male	-	-	-	1	1	4	4	2	3	-	-	-
Female	-	-	1	1	2	1	4	11	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	3	-	-	1	8	5	8	3	1	-
Male	-	1	2	-	-	-	2	1	-	-	-	-
Female	-	-	1	-	-	1	6	4	8	3	1	-
White, both sexes	-	1	2	-	-	1	8	5	8	3	1	-
Male	-	1	1	-	-	-	2	1	-	-	-	-
Female	-	-	1	-	-	1	6	4	8	3	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Kentucky												
All races, both sexes	-	1	2	5	1	10	6	8	8	2	-	-
Male	-	1	-	2	-	5	1	-	2	1	-	-
Female	-	-	2	3	1	5	5	8	6	1	-	-
White, both sexes	-	1	2	4	1	10	6	8	7	2	-	-
Male	-	1	-	1	-	5	1	-	2	1	-	-
Female	-	-	2	3	1	5	5	8	5	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	3	2	4	9	6	12	7	-	-	-
Male	-	-	-	-	1	2	1	3	2	-	-	-
Female	-	-	3	2	3	7	5	9	5	-	-	-
White, both sexes	-	-	1	1	3	5	5	10	5	-	-	-
Male	-	-	-	-	1	1	1	3	1	-	-	-
Female	-	-	1	1	2	4	4	7	4	-	-	-
Black, both sexes	-	-	2	1	1	4	1	2	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	2	1	1	3	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Maine												
All races, both sexes	-	-	-	2	2	3	5	4	5	-	-	-
Male	-	-	-	1	-	-	3	1	-	-	-	-
Female	-	-	-	1	2	3	2	3	5	-	-	-
White, both sexes	-	-	-	2	2	3	5	4	5	-	-	-
Male	-	-	-	1	-	-	3	1	-	-	-	-
Female	-	-	-	1	2	3	2	3	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	2	2	7	11	7	16	11	3	-	-
Male	1	-	1	1	1	2	3	3	2	-	-	-
Female	-	-	1	1	6	9	4	13	9	3	-	-
White, both sexes	-	-	1	2	6	10	6	15	10	3	-	-
Male	-	-	1	1	1	2	3	3	2	-	-	-
Female	-	-	-	1	5	8	3	12	8	3	-	-
Black, both sexes	1	-	1	-	1	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Massachusetts												
All races, both sexes	2	3	4	4	5	5	19	19	13	7	-	-
Male	2	1	3	2	2	1	5	3	3	1	-	-
Female	-	2	1	2	3	4	14	16	10	6	-	-
White, both sexes	1	3	4	3	5	4	19	19	13	6	-	-
Male	1	1	3	1	2	1	5	3	3	1	-	-
Female	-	2	1	2	3	3	14	16	10	5	-	-
Black, both sexes	1	-	-	1	-	1	-	-	-	1	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	3	3	4	16	19	19	20	18	6	-	-
Male	-	-	1	1	6	5	5	6	4	-	-	-
Female	-	3	2	3	10	14	14	14	14	6	-	-
White, both sexes	-	1	3	4	13	19	17	19	15	6	-	-
Male	-	-	1	1	5	5	4	6	3	-	-	-
Female	-	1	2	3	8	14	13	13	12	6	-	-
Black, both sexes	-	2	-	-	3	-	2	1	3	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	2	-	-	2	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Minnesota												
All races, both sexes	1	3	1	3	3	6	14	8	8	2	1	-
Male	-	3	1	3	3	2	6	1	4	2	-	-
Female	1	-	-	-	-	4	8	7	4	-	1	-
White, both sexes	1	3	1	3	3	6	14	8	8	2	1	-
Male	-	3	1	3	3	2	6	1	4	2	-	-
Female	1	-	-	-	-	4	8	7	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	2	2	6	6	1	-	3	-	-
Male	-	1	-	2	1	3	1	-	-	-	-	-
Female	-	-	-	-	1	3	5	1	-	3	-	-
White, both sexes	-	1	-	-	2	5	6	-	-	3	-	-
Male	-	1	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	3	5	-	-	3	-	-
Black, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Missouri												
All races, both sexes	1	-	1	5	7	10	10	16	13	3	-	-
Male	-	-	-	-	4	5	2	3	2	-	-	-
Female	1	-	1	5	3	5	8	13	11	3	-	-
White, both sexes	1	-	1	4	7	9	10	16	13	3	-	-
Male	-	-	-	-	4	4	2	3	2	-	-	-
Female	1	-	1	4	3	5	8	13	11	3	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	1	-	5	6	1	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	1	-	1	-	4	4	1	-	-
White, both sexes	-	-	-	1	-	1	-	5	6	1	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	1	-	1	-	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Nebraska												
All races, both sexes	-	-	1	2	-	1	8	2	1	2	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	-	1	-	1	6	2	-	2	-	-
White, both sexes	-	-	1	2	-	1	8	2	1	2	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	-	1	-	1	6	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	3	3	3	1	1	3	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	2	2	2	1	1	3	-	-	-
White, both sexes	-	-	1	3	3	3	1	1	3	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	2	2	2	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
New Hampshire												
All races, both sexes	-	1	1	2	1	3	4	4	4	-	-	-
Male	-	-	-	1	-	-	2	2	1	-	-	-
Female	-	1	1	1	1	3	2	2	3	-	-	-
White, both sexes	-	1	1	2	1	3	4	4	4	-	-	-
Male	-	-	-	1	-	-	2	2	1	-	-	-
Female	-	1	1	1	1	3	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	5	3	3	9	18	23	12	10	1	-	-
Male	-	1	2	1	5	8	9	4	1	-	-	-
Female	-	4	1	2	4	10	14	8	9	1	-	-
White, both sexes	-	4	1	3	9	15	19	9	8	1	-	-
Male	-	1	1	1	5	7	7	4	1	-	-	-
Female	-	3	-	2	4	8	12	5	7	1	-	-
Black, both sexes	-	1	2	-	-	3	4	3	2	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	-	1	1	-	-	2	2	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
New Mexico												
All races, both sexes	-	-	1	3	2	3	5	5	1	-	-	-
Male	-	-	1	2	2	2	3	1	-	-	-	-
Female	-	-	-	1	-	1	2	4	1	-	-	-
White, both sexes	-	-	1	3	2	2	5	4	1	-	-	-
Male	-	-	1	2	2	1	3	1	-	-	-	-
Female	-	-	-	1	-	1	2	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	3	12	14	21	32	49	34	33	5	1	-
Male	3	1	3	5	10	14	19	8	7	-	-	-
Female	4	2	9	9	11	18	30	26	26	5	1	-
White, both sexes	5	2	9	9	20	28	42	30	30	4	1	-
Male	2	-	2	3	9	13	15	8	7	-	-	-
Female	3	2	7	6	11	15	27	22	23	4	1	-
Black, both sexes	2	1	2	3	1	3	5	4	3	1	-	-
Male	1	1	1	1	1	1	2	-	-	-	-	-
Female	1	-	1	2	-	2	3	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	2	-	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
North Carolina												
All races, both sexes	2	-	2	4	9	17	18	26	12	4	1	-
Male	-	-	1	1	2	5	6	4	4	-	-	-
Female	2	-	1	3	7	12	12	22	8	4	1	-
White, both sexes	1	-	2	3	9	15	14	24	11	4	1	-
Male	-	-	1	-	2	5	4	4	4	-	-	-
Female	1	-	1	3	7	10	10	20	7	4	1	-
Black, both sexes	1	-	-	1	-	2	4	2	1	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	2	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	-	1	1	4	2	1	-	-
Male	-	-	1	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	2	1	-	-
White, both sexes	-	-	1	-	-	1	1	4	2	1	-	-
Male	-	-	1	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Ohio												
All races, both sexes	2	9	8	9	9	27	32	25	19	9	1	-
Male	2	5	2	3	3	11	12	8	4	1	-	-
Female	-	4	6	6	6	16	20	17	15	8	1	-
White, both sexes	2	8	8	8	8	24	31	25	17	8	1	-
Male	2	4	2	2	2	9	12	8	3	1	-	-
Female	-	4	6	6	6	15	19	17	14	7	1	-
Black, both sexes	-	1	-	1	1	3	1	-	2	1	-	-
Male	-	1	-	1	1	2	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	2	7	7	7	3	6	2	2	-
Male	2	-	-	-	3	2	1	-	-	-	-	-
Female	-	-	-	2	4	5	6	3	6	2	2	-
White, both sexes	2	-	-	1	7	6	7	3	6	2	1	-
Male	2	-	-	-	3	2	1	-	-	-	-	-
Female	-	-	-	1	4	4	6	3	6	2	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Oregon												
All races, both sexes	-	1	-	4	3	5	8	10	10	2	-	-
Male	-	1	-	2	-	3	2	2	2	-	-	-
Female	-	-	-	2	3	2	6	8	8	2	-	-
White, both sexes	-	1	-	4	3	5	8	10	10	2	-	-
Male	-	1	-	2	-	3	2	2	2	-	-	-
Female	-	-	-	2	3	2	6	8	8	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	3	1	8	12	17	22	40	24	15	-	-
Male	-	1	1	1	3	5	7	9	3	2	-	-
Female	2	2	-	7	9	12	15	31	21	13	-	-
White, both sexes	2	1	1	7	11	16	22	40	24	15	-	-
Male	-	1	1	-	3	5	7	9	3	2	-	-
Female	2	-	-	7	8	11	15	31	21	13	-	-
Black, both sexes	-	2	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	2	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Rhode Island												
All races, both sexes	1	-	-	1	-	4	3	3	3	1	1	-
Male	1	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	3	2	2	3	1	1	-
White, both sexes	1	-	-	1	-	4	3	3	3	1	1	-
Male	1	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	3	2	2	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	2	4	7	12	11	8	1	-	-
Male	1	-	1	1	2	3	4	-	-	-	-	-
Female	-	1	-	1	2	4	8	11	8	1	-	-
White, both sexes	1	-	1	1	3	5	10	9	8	1	-	-
Male	1	-	1	-	1	3	4	-	-	-	-	-
Female	-	-	-	1	2	2	6	9	8	1	-	-
Black, both sexes	-	1	-	1	1	2	2	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	1	5	3	2	-	-
Male	-	-	-	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	-	-	1	3	2	2	-	-
White, both sexes	-	-	-	-	-	1	1	5	3	2	-	-
Male	-	-	-	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	-	-	1	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	4	4	2	6	10	18	14	12	3	-	-
Male	3	2	1	-	3	1	5	6	2	-	-	-
Female	-	2	3	2	3	9	13	8	10	3	-	-
White, both sexes	2	3	3	2	6	10	17	12	12	2	-	-
Male	2	2	-	-	3	1	5	5	2	-	-	-
Female	-	1	3	2	3	9	12	7	10	2	-	-
Black, both sexes	1	1	1	-	-	-	1	2	-	1	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Texas												
All races, both sexes	5	16	10	15	10	27	23	26	21	8	3	-
Male	3	7	1	7	4	5	8	10	4	2	1	-
Female	2	9	9	8	6	22	15	16	17	6	2	-
White, both sexes	5	9	10	13	10	23	20	25	18	7	2	-
Male	3	3	1	6	4	4	7	10	4	2	1	-
Female	2	6	9	7	6	19	13	15	14	5	1	-
Black, both sexes	-	7	-	2	-	4	3	1	3	1	1	-
Male	-	4	-	1	-	1	1	-	-	-	-	-
Female	-	3	-	1	-	3	2	1	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	1	1	3	2	1	1	-	-
Male	-	-	1	1	1	-	1	-	1	1	-	-
Female	-	-	-	-	-	1	2	2	-	-	-	-
White, both sexes	-	-	1	1	1	1	2	2	1	1	-	-
Male	-	-	1	1	1	-	-	-	1	1	-	-
Female	-	-	-	-	-	1	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Vermont												
All races, both sexes	-	-	-	-	-	2	4	3	3	3	1	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	4	3	2	3	1	-
White, both sexes	-	-	-	-	-	2	4	3	3	3	1	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	4	3	2	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	1	4	3	10	15	11	10	5	-	-
Male	-	-	1	-	2	3	4	2	3	1	-	-
Female	-	-	-	4	1	7	11	9	7	4	-	-
White, both sexes	-	-	1	3	1	8	14	11	9	4	-	-
Male	-	-	1	-	-	3	3	2	3	1	-	-
Female	-	-	-	3	1	5	11	9	6	3	-	-
Black, both sexes	-	-	-	1	2	1	1	-	1	1	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Washington												
All races, both sexes	3	2	-	3	2	9	15	9	13	7	-	-
Male	1	1	-	1	-	3	4	3	4	2	-	-
Female	2	1	-	2	2	6	11	6	9	5	-	-
White, both sexes	3	1	-	3	2	9	15	8	13	7	-	-
Male	1	-	-	1	-	3	4	3	4	2	-	-
Female	2	1	-	2	2	6	11	5	9	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	1	1	1	1	-	2	5	7	1	-	-	-
Male	-	1	-	-	-	1	1	2	-	-	-	-
Female	1	-	1	1	-	1	4	5	1	-	-	-
White, both sexes	1	1	1	1	-	2	5	7	1	-	-	-
Male	-	1	-	-	-	1	1	2	-	-	-	-
Female	1	-	1	1	-	1	4	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Wisconsin												
All races, both sexes	-	1	2	4	4	7	12	19	15	7	-	-
Male	-	-	-	1	1	3	4	2	6	1	-	-
Female	-	1	2	3	3	4	8	17	9	6	-	-
White, both sexes	-	1	-	4	4	7	12	19	15	7	-	-
Male	-	-	-	1	1	3	4	2	6	1	-	-
Female	-	1	-	3	3	4	8	17	9	6	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	2	-	2	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	2	2	-	-	-
White, both sexes	-	1	-	-	-	2	-	2	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
United States												
All races, both sexes	5,653	268	31	12	5	9	7	16	20	30	65	101
Male	2,163	166	18	9	2	7	2	11	12	15	37	45
Female	3,490	102	13	3	3	2	5	5	8	15	28	56
White, both sexes	4,922	149	26	9	2	6	7	13	14	20	49	76
Male	1,815	92	15	6	-	4	2	8	7	7	32	38
Female	3,107	57	11	3	2	2	5	5	7	13	17	38
Black, both sexes	602	110	3	3	2	3	-	3	5	7	10	22
Male	292	71	1	3	2	3	-	3	4	5	4	7
Female	310	39	2	-	-	-	-	-	1	2	6	15
American Indian 1/, both sexes	50	4	2	-	-	-	-	-	-	-	5	2
Male	21	2	2	-	-	-	-	-	-	-	1	-
Female	29	2	-	-	-	-	-	-	-	-	4	2
Asian or Pacific Islander 2/, both sexes	79	5	-	-	1	-	-	-	1	3	1	1
Male	35	1	-	-	-	-	-	-	1	3	-	-
Female	44	4	-	-	1	-	-	-	-	-	1	1
Alabama												
All races, both sexes	113	10	-	1	-	1	1	-	-	-	-	3
Male	46	6	-	1	-	1	1	-	-	-	-	2
Female	67	4	-	-	-	-	-	-	-	-	-	1
White, both sexes	85	2	-	-	-	-	1	-	-	-	-	2
Male	35	2	-	-	-	-	1	-	-	-	-	2
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	8	-	1	-	1	-	-	-	-	-	1
Male	11	4	-	1	-	1	-	-	-	-	-	-
Female	17	4	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Alaska												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	99	5	3	-	-	-	-	1	-	-	-	2
Male	34	3	2	-	-	-	-	1	-	-	-	1
Female	65	2	1	-	-	-	-	-	-	-	-	1
White, both sexes	94	5	3	-	-	-	-	1	-	-	-	1
Male	31	3	2	-	-	-	-	1	-	-	-	1
Female	63	2	1	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Arkansas												
All races, both sexes	44	3	-	-	-	-	-	-	-	-	-	-
Male	23	3	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	1	-	-	-	-	-	-	-	-	-	-
Male	19	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	2	-	-	-	-	-	-	-	-	-	-
Male	4	2	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	449	32	2	3	1	-	2	5	1	2	8	13
Male	166	19	2	2	-	-	-	3	-	-	4	4
Female	283	13	-	1	1	-	2	2	1	2	4	9
White, both sexes	398	20	2	3	-	-	2	3	1	2	7	11
Male	139	13	2	2	-	-	-	1	-	-	3	4
Female	259	7	-	1	-	-	2	2	1	2	4	7
Black, both sexes	32	11	-	-	-	-	-	2	-	-	1	2
Male	18	6	-	-	-	-	-	2	-	-	1	-
Female	14	5	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	1	-	-	1	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Colorado												
All races, both sexes	67	1	-	-	-	-	1	-	-	1	2	3
Male	26	1	-	-	-	-	-	-	-	1	1	1
Female	41	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	63	1	-	-	-	-	1	-	-	1	2	3
Male	25	1	-	-	-	-	-	-	-	1	1	1
Female	38	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	90	3	-	-	-	-	-	-	-	-	-	2
Male	32	3	-	-	-	-	-	-	-	-	-	2
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	2
Male	26	-	-	-	-	-	-	-	-	-	-	2
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	2	-	-	-	-	-	-	-	-	-	-
Male	4	2	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Delaware												
All races, both sexes	16	-	-	-	1	-	-	-	-	1	-	-
Male	7	-	-	-	1	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	2	1	-	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	1	1	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Florida												
All races, both sexes	363	24	2	-	-	-	-	-	1	5	5	4
Male	125	13	-	-	-	-	-	-	-	1	2	2
Female	238	11	2	-	-	-	-	-	1	4	3	2
White, both sexes	324	11	2	-	-	-	-	-	1	5	3	4
Male	106	7	-	-	-	-	-	-	-	1	2	2
Female	218	4	2	-	-	-	-	-	1	4	1	2
Black, both sexes	37	13	-	-	-	-	-	-	-	-	2	-
Male	19	6	-	-	-	-	-	-	-	-	-	-
Female	18	7	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	172	19	-	-	1	-	1	1	2	-	3	2
Male	69	14	-	-	1	-	-	1	-	-	-	1
Female	103	5	-	-	-	-	1	-	2	-	3	1
White, both sexes	116	9	-	-	-	-	1	1	1	-	1	1
Male	43	7	-	-	-	-	-	1	-	-	-	-
Female	73	2	-	-	-	-	1	-	1	-	1	1
Black, both sexes	55	9	-	-	1	-	-	-	1	-	2	1
Male	25	6	-	-	1	-	-	-	-	-	-	1
Female	30	3	-	-	-	-	-	-	1	-	2	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Hawaii												
All races, both sexes	21	2	-	-	-	-	1	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	14	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	8	1	-	-	-	-	1	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	21	2	-	-	-	-	-	-	-	-	-	1
Male	8	1	-	-	-	-	-	-	-	-	-	1
Female	13	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	2	-	-	-	-	-	-	-	-	-	1
Male	8	1	-	-	-	-	-	-	-	-	-	1
Female	13	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Illinois												
All races, both sexes	262	6	1	-	-	-	-	-	-	1	1	9
Male	97	6	-	-	-	-	-	-	-	1	1	3
Female	165	-	1	-	-	-	-	-	-	-	-	6
White, both sexes	226	2	-	-	-	-	-	-	-	-	1	5
Male	77	2	-	-	-	-	-	-	-	-	1	1
Female	149	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	34	4	1	-	-	-	-	-	-	1	-	4
Male	19	4	-	-	-	-	-	-	-	1	-	2
Female	15	-	1	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	126	5	1	1	-	-	-	-	1	-	2	2
Male	42	3	-	1	-	-	-	-	-	-	1	-
Female	84	2	1	-	-	-	-	-	1	-	1	2
White, both sexes	117	2	1	1	-	-	-	-	1	-	2	1
Male	39	2	-	1	-	-	-	-	-	-	1	-
Female	78	-	1	-	-	-	-	-	1	-	1	1
Black, both sexes	8	3	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Iowa												
All races, both sexes	65	1	-	-	-	-	-	1	-	-	-	-
Male	23	1	-	-	-	-	-	1	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	1	-	-	-	-	-	1	-	-	-	-
Male	23	1	-	-	-	-	-	1	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	67	2	1	-	-	-	-	-	-	-	1	1
Male	32	2	-	-	-	-	-	-	-	-	1	1
Female	35	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	1	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	34	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	4	2	-	-	-	-	-	-	-	-	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Kentucky												
All races, both sexes	89	3	-	-	-	-	-	-	-	1	-	2
Male	31	2	-	-	-	-	-	-	-	1	-	1
Female	58	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	85	2	-	-	-	-	-	-	-	1	-	2
Male	28	1	-	-	-	-	-	-	-	1	-	1
Female	57	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	88	3	-	-	-	1	-	1	-	2	-	3
Male	37	1	-	-	-	1	-	1	-	2	-	1
Female	51	2	-	-	-	-	-	-	-	-	-	2
White, both sexes	67	1	-	-	-	-	-	1	-	-	-	-
Male	24	1	-	-	-	-	-	1	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	21	2	-	-	-	1	-	-	-	2	-	3
Male	13	-	-	-	-	1	-	-	-	2	-	1
Female	8	2	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Maine												
All races, both sexes	34	1	-	-	-	-	-	-	-	1	2	-
Male	14	-	-	-	-	-	-	-	-	1	2	-
Female	20	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	1	-	-	-	-	-	-	-	1	2	-
Male	14	-	-	-	-	-	-	-	-	1	2	-
Female	20	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	102	6	-	-	-	-	-	1	-	-	1	3
Male	32	2	-	-	-	-	-	-	-	-	1	-
Female	70	4	-	-	-	-	-	1	-	-	-	3
White, both sexes	73	3	-	-	-	-	-	1	-	-	1	1
Male	23	1	-	-	-	-	-	-	-	-	1	-
Female	50	2	-	-	-	-	-	1	-	-	-	1
Black, both sexes	29	3	-	-	-	-	-	-	-	-	-	2
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	20	2	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Massachusetts												
All races, both sexes	174	4	1	-	-	-	-	-	1	1	2	4
Male	69	2	1	-	-	-	-	-	1	-	-	1
Female	105	2	-	-	-	-	-	-	-	1	2	3
White, both sexes	164	3	1	-	-	-	-	-	1	1	1	3
Male	67	2	1	-	-	-	-	-	1	-	-	1
Female	97	1	-	-	-	-	-	-	-	1	1	2
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	197	10	1	-	-	1	-	1	-	2	-	4
Male	80	7	1	-	-	1	-	1	-	1	-	2
Female	117	3	-	-	-	-	-	-	-	1	-	2
White, both sexes	174	7	1	-	-	-	-	-	-	1	-	3
Male	64	5	1	-	-	-	-	-	-	-	-	2
Female	110	2	-	-	-	-	-	-	-	1	-	1
Black, both sexes	22	3	-	-	-	1	-	1	-	1	-	1
Male	15	2	-	-	-	1	-	1	-	1	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Minnesota												
All races, both sexes	91	4	-	-	-	-	-	-	1	1	1	2
Male	39	2	-	-	-	-	-	-	1	-	1	1
Female	52	2	-	-	-	-	-	-	-	1	-	1
White, both sexes	85	3	-	-	-	-	-	-	1	1	-	1
Male	36	2	-	-	-	-	-	-	1	-	-	1
Female	49	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	48	5	-	-	-	-	-	-	-	-	-	1
Male	19	4	-	-	-	-	-	-	-	-	-	-
Female	29	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	1	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	4	-	-	-	-	-	-	-	-	-	1
Male	8	4	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Missouri												
All races, both sexes	132	2	3	1	-	-	-	-	-	-	1	2
Male	51	1	2	-	-	-	-	-	-	-	1	-
Female	81	1	1	1	-	-	-	-	-	-	-	2
White, both sexes	122	-	3	1	-	-	-	-	-	-	1	1
Male	47	-	2	-	-	-	-	-	-	-	1	-
Female	75	-	1	1	-	-	-	-	-	-	-	1
Black, both sexes	9	2	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Nebraska												
All races, both sexes	39	2	1	-	-	-	-	-	-	-	1	-
Male	15	2	1	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	2	1	-	-	-	-	-	-	-	1	-
Male	14	2	1	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	47	1	-	-	-	-	-	-	-	-	1	2
Male	27	1	-	-	-	-	-	-	-	-	1	2
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	2
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	212	7	1	-	-	1	-	1	2	1	-	1
Male	79	3	1	-	-	-	-	-	2	1	-	1
Female	133	4	-	-	-	1	-	1	-	-	-	-
White, both sexes	184	5	1	-	-	1	-	1	-	-	-	1
Male	66	1	1	-	-	-	-	-	-	-	-	1
Female	118	4	-	-	-	1	-	1	-	-	-	-
Black, both sexes	22	2	-	-	-	-	-	-	2	1	-	-
Male	11	2	-	-	-	-	-	-	2	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
New Mexico												
All races, both sexes	38	1	2	-	-	-	-	1	1	-	3	1
Male	15	-	2	-	-	-	-	1	-	-	2	1
Female	23	1	-	-	-	-	-	-	1	-	1	-
White, both sexes	30	-	1	-	-	-	-	1	1	-	2	1
Male	12	-	1	-	-	-	-	1	-	-	2	1
Female	18	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	1	1	-	-	-	-	-	-	-	1	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	342	13	3	1	-	3	-	-	2	2	4	3
Male	127	11	1	1	-	3	-	-	2	1	2	2
Female	215	2	2	-	-	-	-	-	-	1	2	1
White, both sexes	294	6	3	-	-	3	-	-	1	1	3	3
Male	103	5	1	-	-	3	-	-	1	-	1	2
Female	191	1	2	-	-	-	-	-	-	1	2	1
Black, both sexes	38	7	-	1	-	-	-	-	1	-	1	-
Male	19	6	-	1	-	-	-	-	1	-	1	-
Female	19	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
North Carolina												
All races, both sexes	175	5	-	-	-	-	-	-	1	1	6	2
Male	54	3	-	-	-	-	-	-	-	1	1	2
Female	121	2	-	-	-	-	-	-	1	-	5	-
White, both sexes	141	2	-	-	-	-	-	-	1	1	3	2
Male	40	-	-	-	-	-	-	-	-	1	1	2
Female	101	2	-	-	-	-	-	-	1	-	2	-
Black, both sexes	31	3	-	-	-	-	-	-	-	-	2	-
Male	14	3	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Ohio												
All races, both sexes	294	9	1	-	-	-	1	2	1	-	4	2
Male	129	7	-	-	-	-	1	1	1	-	4	-
Female	165	2	1	-	-	-	-	1	-	-	-	2
White, both sexes	270	5	1	-	-	-	1	2	1	-	4	2
Male	116	4	-	-	-	-	1	1	1	-	4	-
Female	154	1	1	-	-	-	-	1	-	-	-	2
Black, both sexes	20	4	-	-	-	-	-	-	-	-	-	-
Male	12	3	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	79	3	-	-	-	-	-	-	-	-	3	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	54	3	-	-	-	-	-	-	-	-	3	1
White, both sexes	66	3	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	46	3	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Oregon												
All races, both sexes	77	2	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	54	2	-	-	-	-	-	-	-	-	-	1
White, both sexes	75	1	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	52	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	321	6	2	-	-	2	-	-	1	-	1	6
Male	116	6	2	-	-	1	-	-	1	-	1	3
Female	205	-	-	-	-	1	-	-	-	-	-	3
White, both sexes	287	4	1	-	-	2	-	-	1	-	1	4
Male	103	4	1	-	-	1	-	-	1	-	1	2
Female	184	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	34	2	1	-	-	-	-	-	-	-	-	2
Male	13	2	1	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Rhode Island												
All races, both sexes	23	-	-	1	-	-	-	-	-	1	1	-
Male	12	-	-	1	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	1	-	-	-	-	-	-	1	-
Male	11	-	-	1	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	98	5	-	1	1	-	-	-	-	1	-	3
Male	39	3	-	-	-	-	-	-	-	-	-	2
Female	59	2	-	1	1	-	-	-	-	1	-	1
White, both sexes	73	1	-	1	1	-	-	-	-	1	-	2
Male	28	1	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	1	1	-	-	-	-	1	-	-
Black, both sexes	25	4	-	-	-	-	-	-	-	-	-	1
Male	11	2	-	-	-	-	-	-	-	-	-	-
Female	14	2	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
South Dakota												
All races, both sexes	19	1	-	-	-	-	-	-	-	1	1	1
Male	7	1	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	1	-
Male	2	1	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	120	5	1	-	-	-	-	-	1	-	1	2
Male	51	4	1	-	-	-	-	-	1	-	1	1
Female	69	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	100	1	1	-	-	-	-	-	-	-	1	2
Male	43	1	1	-	-	-	-	-	-	-	1	1
Female	57	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	20	4	-	-	-	-	-	-	1	-	-	-
Male	8	3	-	-	-	-	-	-	1	-	-	-
Female	12	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Texas												
All races, both sexes	328	35	2	2	1	-	-	1	2	3	7	7
Male	141	19	1	2	-	-	-	1	2	1	7	2
Female	187	16	1	-	1	-	-	-	-	2	-	5
White, both sexes	290	29	2	1	1	-	-	1	1	1	6	6
Male	119	16	1	1	-	-	-	1	1	-	6	1
Female	171	13	1	-	1	-	-	-	-	1	-	5
Black, both sexes	30	6	-	1	-	-	-	-	-	1	1	1
Male	17	3	-	1	-	-	-	-	-	-	1	1
Female	13	3	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Vermont												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	148	5	-	-	-	-	-	-	-	1	1	2
Male	56	2	-	-	-	-	-	-	-	-	-	1
Female	92	3	-	-	-	-	-	-	-	1	1	1
White, both sexes	125	3	-	-	-	-	-	-	-	-	1	1
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	82	3	-	-	-	-	-	-	-	-	1	1
Black, both sexes	23	2	-	-	-	-	-	-	-	1	-	1
Male	13	2	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Washington												
All races, both sexes	106	7	1	-	-	-	-	-	1	-	1	-
Male	38	4	-	-	-	-	-	-	-	-	1	-
Female	68	3	1	-	-	-	-	-	1	-	-	-
White, both sexes	95	5	1	-	-	-	-	-	1	-	1	-
Male	33	3	-	-	-	-	-	-	-	-	1	-
Female	62	2	1	-	-	-	-	-	1	-	-	-
Black, both sexes	5	2	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	48	-	-	1	-	-	-	-	-	1	-	-
Male	14	-	-	1	-	-	-	-	-	1	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	1	-	-	-	-	-	1	-	-
Male	14	-	-	1	-	-	-	-	-	1	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of intestines and peritoneum (K52,K58-K63)												
Wisconsin												
All races, both sexes	85	4	1	-	-	-	-	-	1	-	-	1
Male	41	1	1	-	-	-	-	-	1	-	-	1
Female	44	3	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	2	-	-	-	-	-	-	1	-	-	1
Male	39	1	-	-	-	-	-	-	1	-	-	1
Female	39	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	2	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
United States												
All races, both sexes	146	186	282	360	525	818	1,005	890	618	223	36	-
Male	74	91	121	166	242	322	368	251	155	42	7	-
Female	72	95	161	194	283	496	637	639	463	181	29	-
White, both sexes	117	150	231	306	464	734	913	822	576	206	32	-
Male	55	71	95	143	213	283	336	227	139	36	6	-
Female	62	79	136	163	251	451	577	595	437	170	26	-
Black, both sexes	23	26	44	45	46	64	76	57	35	14	4	-
Male	16	15	22	19	23	30	27	18	13	5	1	-
Female	7	11	22	26	23	34	49	39	22	9	3	-
American Indian 1/, both sexes	3	5	4	5	4	8	6	1	1	-	-	-
Male	1	3	3	2	1	3	2	1	-	-	-	-
Female	2	2	1	3	3	5	4	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	3	4	11	12	10	10	6	3	-	-
Male	2	2	1	2	5	6	3	5	3	1	-	-
Female	1	3	2	2	6	6	7	5	3	2	-	-
Alabama												
All races, both sexes	4	1	4	14	11	15	20	15	10	2	1	-
Male	-	1	1	6	7	5	8	4	1	1	1	-
Female	4	-	3	8	4	10	12	11	9	1	-	-
White, both sexes	3	-	4	12	9	14	15	12	9	2	-	-
Male	-	-	1	6	6	4	7	4	1	1	-	-
Female	3	-	3	6	3	10	8	8	8	1	-	-
Black, both sexes	1	1	-	2	2	1	5	3	1	-	1	-
Male	-	1	-	-	1	1	1	-	-	-	1	-
Female	1	-	-	2	1	-	4	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Alaska												
All races, both sexes	1	1	-	-	-	2	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	6	4	7	9	15	11	16	15	2	-	-
Male	2	3	2	2	1	5	5	5	2	-	-	-
Female	1	3	2	5	8	10	6	11	13	2	-	-
White, both sexes	2	4	4	7	9	14	11	16	15	2	-	-
Male	1	1	2	2	1	5	5	5	2	-	-	-
Female	1	3	2	5	8	9	6	11	13	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Arkansas												
All races, both sexes	2	-	4	4	6	8	10	4	2	1	-	-
Male	1	-	1	2	3	4	8	1	-	-	-	-
Female	1	-	3	2	3	4	2	3	2	1	-	-
White, both sexes	2	-	3	4	5	8	7	4	1	1	-	-
Male	1	-	1	2	3	4	6	1	-	-	-	-
Female	1	-	2	2	2	4	1	3	1	1	-	-
Black, both sexes	-	-	1	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	14	9	24	29	31	55	78	64	52	19	5	-
Male	7	3	11	15	13	17	26	25	11	3	1	-
Female	7	6	13	14	18	38	52	39	41	16	4	-
White, both sexes	12	9	19	25	29	48	75	59	48	18	5	-
Male	5	3	9	12	12	13	26	22	9	2	1	-
Female	7	6	10	13	17	35	49	37	39	16	4	-
Black, both sexes	1	-	2	4	1	3	1	2	2	-	-	-
Male	1	-	1	3	-	1	-	2	1	-	-	-
Female	-	-	1	1	1	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	-	1	4	2	3	2	1	-	-
Male	1	-	1	-	1	3	-	1	1	1	-	-
Female	-	-	2	-	-	1	2	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Colorado												
All races, both sexes	4	4	1	4	7	9	9	14	6	-	1	-
Male	2	2	-	2	3	5	3	4	1	-	-	-
Female	2	2	1	2	4	4	6	10	5	-	1	-
White, both sexes	3	4	1	2	7	9	8	14	6	-	1	-
Male	1	2	-	2	3	5	3	4	1	-	-	-
Female	2	2	1	-	4	4	5	10	5	-	1	-
Black, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	3	10	6	4	16	12	16	10	6	-	-
Male	-	2	6	2	2	5	3	5	2	-	-	-
Female	2	1	4	4	2	11	9	11	8	6	-	-
White, both sexes	2	3	7	5	4	16	11	16	10	6	-	-
Male	-	2	4	2	2	5	2	5	2	-	-	-
Female	2	1	3	3	2	11	9	11	8	6	-	-
Black, both sexes	-	-	2	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Delaware												
All races, both sexes	-	-	-	1	1	6	2	2	2	-	-	-
Male	-	-	-	-	1	4	-	-	-	-	-	-
Female	-	-	-	1	-	2	2	2	2	-	-	-
White, both sexes	-	-	-	1	1	6	2	1	2	-	-	-
Male	-	-	-	-	1	4	-	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	2	1	1	2	1	-	-	-	-
Male	-	1	-	1	1	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	1	-	1	1	-	1	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Florida												
All races, both sexes	10	16	16	17	16	62	71	60	43	11	-	-
Male	6	7	7	7	10	22	21	18	7	2	-	-
Female	4	9	9	10	6	40	50	42	36	9	-	-
White, both sexes	8	14	9	16	14	60	68	56	42	11	-	-
Male	4	5	4	6	10	20	19	18	6	2	-	-
Female	4	9	5	10	4	40	49	38	36	9	-	-
Black, both sexes	2	2	7	1	1	2	2	4	1	-	-	-
Male	2	2	3	1	-	2	2	-	1	-	-	-
Female	-	-	4	-	1	-	-	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	4	9	10	13	23	26	17	22	15	4	-	-
Male	2	3	5	7	7	9	5	8	5	1	-	-
Female	2	6	5	6	16	17	12	14	10	3	-	-
White, both sexes	3	3	6	9	18	19	11	18	12	3	-	-
Male	2	1	2	4	5	7	4	6	4	-	-	-
Female	1	2	4	5	13	12	7	12	8	3	-	-
Black, both sexes	1	6	4	4	5	7	6	4	3	1	-	-
Male	-	2	3	3	2	2	1	2	1	1	-	-
Female	1	4	1	1	3	5	5	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Hawaii												
All races, both sexes	-	1	1	2	3	3	2	3	3	-	-	-
Male	-	1	-	1	-	1	-	1	2	-	-	-
Female	-	-	1	1	3	2	2	2	1	-	-	-
White, both sexes	-	-	1	1	1	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	2	1	3	2	-	-	-
Male	-	1	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	1	2	1	1	2	1	-	-	-
Idaho												
All races, both sexes	2	-	1	-	-	-	9	3	2	1	-	-
Male	-	-	1	-	-	-	4	-	1	-	-	-
Female	2	-	-	-	-	-	5	3	1	1	-	-
White, both sexes	2	-	1	-	-	-	9	3	2	1	-	-
Male	-	-	1	-	-	-	4	-	1	-	-	-
Female	2	-	-	-	-	-	5	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Illinois												
All races, both sexes	4	11	13	17	21	41	45	45	37	8	2	-
Male	1	6	7	9	6	15	15	10	16	-	1	-
Female	3	5	6	8	15	26	30	35	21	8	1	-
White, both sexes	1	9	11	16	19	35	41	43	33	8	2	-
Male	-	4	5	8	5	11	15	10	14	-	1	-
Female	1	5	6	8	14	24	26	33	19	8	1	-
Black, both sexes	3	1	2	1	2	6	3	2	4	-	-	-
Male	1	1	2	1	1	4	-	-	2	-	-	-
Female	2	-	-	-	1	2	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4	1	7	11	14	17	15	22	18	5	-	-
Male	2	-	3	4	5	6	8	5	2	2	-	-
Female	2	1	4	7	9	11	7	17	16	3	-	-
White, both sexes	3	1	6	9	14	16	15	22	18	5	-	-
Male	1	-	2	4	5	6	8	5	2	2	-	-
Female	2	1	4	5	9	10	7	17	16	3	-	-
Black, both sexes	1	-	1	2	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Iowa												
All races, both sexes	2	-	2	7	10	13	9	5	12	3	-	-
Male	2	-	-	3	4	7	1	2	2	-	-	-
Female	-	-	2	4	6	6	8	3	10	3	-	-
White, both sexes	2	-	-	7	9	12	9	5	12	3	-	-
Male	2	-	-	3	4	7	1	2	2	-	-	-
Female	-	-	-	4	5	5	8	3	10	3	-	-
Black, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	4	3	-	1	13	11	17	11	1	1	-
Male	-	1	3	-	1	7	4	9	2	1	-	-
Female	-	3	-	-	-	6	7	8	9	-	1	-
White, both sexes	-	4	2	-	-	13	11	15	11	1	1	-
Male	-	1	2	-	-	7	4	8	2	1	-	-
Female	-	3	-	-	-	6	7	7	9	-	1	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Kentucky												
All races, both sexes	1	5	3	7	3	12	17	21	11	3	-	-
Male	-	-	1	3	1	5	9	4	3	1	-	-
Female	1	5	2	4	2	7	8	17	8	2	-	-
White, both sexes	1	5	3	7	3	11	16	21	10	3	-	-
Male	-	-	1	3	1	4	8	4	3	1	-	-
Female	1	5	2	4	2	7	8	17	7	2	-	-
Black, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	2	5	8	6	8	22	8	9	5	1	-
Male	3	1	3	4	3	3	10	2	1	1	-	-
Female	1	1	2	4	3	5	12	6	8	4	1	-
White, both sexes	3	1	4	7	5	7	17	7	9	4	1	-
Male	2	-	2	3	3	2	8	1	1	-	-	-
Female	1	1	2	4	2	5	9	6	8	4	1	-
Black, both sexes	1	1	1	1	1	1	5	1	-	1	-	-
Male	1	1	1	1	-	1	2	1	-	1	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Maine												
All races, both sexes	-	1	2	-	3	7	7	6	1	2	1	-
Male	-	1	1	-	2	4	1	-	-	1	1	-
Female	-	-	1	-	1	3	6	6	1	1	-	-
White, both sexes	-	1	2	-	3	7	7	6	1	2	1	-
Male	-	1	1	-	2	4	1	-	-	1	1	-
Female	-	-	1	-	1	3	6	6	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	7	7	8	11	19	15	11	7	1	-
Male	1	1	3	7	3	3	5	3	3	-	-	-
Female	2	1	4	-	5	8	14	12	8	7	1	-
White, both sexes	1	2	2	5	8	7	15	12	9	5	1	-
Male	-	1	-	5	3	2	4	3	3	-	-	-
Female	1	1	2	-	5	5	11	9	6	5	1	-
Black, both sexes	2	-	5	2	-	4	4	3	2	2	-	-
Male	1	-	3	2	-	1	1	-	-	-	-	-
Female	1	-	2	-	-	3	3	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Massachusetts												
All races, both sexes	4	-	7	13	15	20	34	31	25	12	-	-
Male	2	-	4	5	8	8	18	7	9	3	-	-
Female	2	-	3	8	7	12	16	24	16	9	-	-
White, both sexes	4	-	7	13	14	16	33	31	24	12	-	-
Male	2	-	4	5	8	7	17	7	9	3	-	-
Female	2	-	3	8	6	9	16	24	15	9	-	-
Black, both sexes	-	-	-	-	-	4	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Michigan												
All races, both sexes	3	6	12	11	23	30	32	39	17	5	-	-
Male	3	4	4	7	9	9	14	11	5	1	-	-
Female	-	2	8	4	14	21	18	28	12	4	-	-
White, both sexes	2	6	11	7	23	29	30	36	13	5	-	-
Male	2	4	3	4	9	8	13	10	2	1	-	-
Female	-	2	8	3	14	21	17	26	11	4	-	-
Black, both sexes	1	-	1	4	-	1	2	2	4	-	-	-
Male	1	-	1	3	-	1	1	-	3	-	-	-
Female	-	-	-	1	-	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Minnesota												
All races, both sexes	2	1	1	5	9	7	21	13	16	6	1	-
Male	2	-	-	2	4	2	12	4	6	2	-	-
Female	-	1	1	3	5	5	9	9	10	4	1	-
White, both sexes	2	1	1	5	9	5	20	13	16	6	1	-
Male	2	-	-	2	4	1	11	4	6	2	-	-
Female	-	1	1	3	5	4	9	9	10	4	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	3	3	10	6	6	5	2	5	-	-
Male	-	-	-	2	4	4	2	1	1	1	-	-
Female	2	-	3	1	6	2	4	4	1	4	-	-
White, both sexes	1	-	2	3	8	4	5	3	2	2	-	-
Male	-	-	-	2	2	3	2	-	1	1	-	-
Female	1	-	2	1	6	1	3	3	1	1	-	-
Black, both sexes	1	-	1	-	2	2	1	2	-	3	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	1	-	1	-	-	1	1	1	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Missouri												
All races, both sexes	3	3	8	12	12	17	25	21	15	5	2	-
Male	1	2	3	6	6	8	11	8	2	-	-	-
Female	2	1	5	6	6	9	14	13	13	5	2	-
White, both sexes	3	3	7	12	12	16	24	20	13	4	2	-
Male	1	2	3	6	6	7	10	7	2	-	-	-
Female	2	1	4	6	6	9	14	13	11	4	2	-
Black, both sexes	-	-	1	-	-	1	1	1	2	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	2	-	-	-	2	3	3	4	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	2	-	-	-	1	2	2	4	1	-	-
White, both sexes	-	2	-	-	-	2	2	3	4	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	2	-	-	-	1	1	2	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Nebraska												
All races, both sexes	-	-	1	5	4	8	7	4	5	1	-	-
Male	-	-	1	2	2	2	4	-	-	1	-	-
Female	-	-	-	3	2	6	3	4	5	-	-	-
White, both sexes	-	-	1	3	3	8	7	4	5	1	-	-
Male	-	-	1	2	1	2	4	-	-	1	-	-
Female	-	-	-	1	2	6	3	4	5	-	-	-
Black, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	2	6	6	6	7	9	5	-	1	-	-
Male	-	1	4	2	3	3	5	4	-	1	-	-
Female	1	1	2	4	3	4	4	1	-	-	-	-
White, both sexes	1	2	6	4	6	7	8	5	-	1	-	-
Male	-	1	4	1	3	3	4	4	-	1	-	-
Female	1	1	2	3	3	4	4	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
New Hampshire												
All races, both sexes	3	-	1	1	2	4	7	6	2	1	1	-
Male	3	-	1	-	-	2	2	3	1	-	-	-
Female	-	-	-	1	2	2	5	3	1	1	1	-
White, both sexes	3	-	1	1	2	4	7	6	2	1	1	-
Male	3	-	1	-	-	2	2	3	1	-	-	-
Female	-	-	-	1	2	2	5	3	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	5	11	11	33	26	31	37	30	10	2	-
Male	2	3	6	4	16	11	7	13	7	2	-	-
Female	-	2	5	7	17	15	24	24	23	8	2	-
White, both sexes	2	4	9	7	28	22	28	35	30	9	1	-
Male	2	3	6	3	13	8	7	12	7	2	-	-
Female	-	1	3	4	15	14	21	23	23	7	1	-
Black, both sexes	-	1	2	3	3	3	2	2	-	-	1	-
Male	-	-	-	-	2	3	-	1	-	-	-	-
Female	-	1	2	3	1	-	2	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	1	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
New Mexico												
All races, both sexes	4	2	2	3	1	5	4	3	4	1	-	-
Male	1	-	1	1	-	1	3	1	1	-	-	-
Female	3	2	1	2	1	4	1	2	3	1	-	-
White, both sexes	4	2	2	1	1	3	3	3	4	1	-	-
Male	1	-	1	-	-	1	2	1	1	-	-	-
Female	3	2	1	1	1	2	1	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8	15	22	21	36	49	46	57	40	15	2	-
Male	4	8	11	16	14	16	15	9	10	1	-	-
Female	4	7	11	5	22	33	31	48	30	14	2	-
White, both sexes	7	12	20	19	30	42	38	52	38	14	2	-
Male	3	7	10	14	13	14	11	8	9	1	-	-
Female	4	5	10	5	17	28	27	44	29	13	2	-
Black, both sexes	1	2	2	1	5	5	6	5	1	-	-	-
Male	1	1	1	1	1	1	3	1	-	-	-	-
Female	-	1	1	-	4	4	3	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	1	-	1	1	-	-
Male	-	-	-	1	-	1	1	-	1	-	-	-
Female	-	1	-	-	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
North Carolina												
All races, both sexes	4	5	11	12	14	31	26	24	24	9	-	-
Male	-	2	6	5	5	11	5	6	5	2	-	-
Female	4	3	5	7	9	20	21	18	19	7	-	-
White, both sexes	2	3	10	11	9	25	22	22	21	7	-	-
Male	-	1	5	4	3	9	5	5	4	-	-	-
Female	2	2	5	7	6	16	17	17	17	7	-	-
Black, both sexes	1	2	1	1	5	5	4	2	3	2	-	-
Male	-	1	1	1	2	2	-	1	1	2	-	-
Female	1	1	-	-	3	3	4	1	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	2	1	1	3	1	2	1	1	2	-	-
Male	-	2	1	1	1	-	1	1	-	-	-	-
Female	1	-	-	-	2	1	1	-	1	2	-	-
White, both sexes	1	2	1	1	2	1	1	1	1	2	-	-
Male	-	2	1	1	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	1	1	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Ohio												
All races, both sexes	8	6	8	23	28	51	65	49	29	5	2	-
Male	6	4	3	12	19	23	26	14	6	2	-	-
Female	2	2	5	11	9	28	39	35	23	3	2	-
White, both sexes	7	6	6	21	25	48	62	47	27	4	1	-
Male	5	4	2	12	16	21	25	13	6	1	-	-
Female	2	2	4	9	9	27	37	34	21	3	1	-
Black, both sexes	1	-	2	1	3	1	2	2	2	1	1	-
Male	1	-	1	-	3	1	1	1	-	1	-	-
Female	-	-	1	1	-	-	1	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Oklahoma												
All races, both sexes	2	5	5	5	10	13	11	12	6	3	-	-
Male	1	1	3	-	3	9	5	2	1	-	-	-
Female	1	4	2	5	7	4	6	10	5	3	-	-
White, both sexes	1	2	5	5	8	12	8	11	6	3	-	-
Male	1	-	3	-	2	8	4	1	1	-	-	-
Female	-	2	2	5	6	4	4	10	5	3	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	3	-	-	2	1	2	1	-	-	-	-
Male	-	1	-	-	1	1	1	1	-	-	-	-
Female	1	2	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Oregon												
All races, both sexes	1	5	1	1	8	14	17	13	8	3	1	-
Male	1	2	-	1	3	5	2	4	1	2	-	-
Female	-	3	1	-	5	9	15	9	7	1	1	-
White, both sexes	1	5	1	1	8	14	17	12	8	3	1	-
Male	1	2	-	1	3	5	2	4	1	2	-	-
Female	-	3	1	-	5	9	15	8	7	1	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	11	12	13	41	47	64	62	34	11	3	-
Male	4	3	6	2	22	20	19	14	11	-	1	-
Female	1	8	6	11	19	27	45	48	23	11	2	-
White, both sexes	3	7	11	10	38	44	57	59	33	10	2	-
Male	2	1	6	2	20	19	17	14	11	-	1	-
Female	1	6	5	8	18	25	40	45	22	10	1	-
Black, both sexes	2	4	1	3	3	3	7	3	1	1	1	-
Male	2	2	-	-	2	1	2	-	-	-	-	-
Female	-	2	1	3	1	2	5	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Rhode Island												
All races, both sexes	2	1	-	-	2	4	5	2	3	1	-	-
Male	-	1	-	-	1	2	4	1	1	-	-	-
Female	2	-	-	-	1	2	1	1	2	1	-	-
White, both sexes	1	1	-	-	2	4	5	2	3	1	-	-
Male	-	1	-	-	1	2	4	1	1	-	-	-
Female	1	-	-	-	1	2	1	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	1	6	9	11	18	18	13	4	3	-	-
Male	3	1	2	3	7	11	4	2	1	-	-	-
Female	1	-	4	6	4	7	14	11	3	3	-	-
White, both sexes	2	1	3	6	9	17	16	7	3	3	-	-
Male	1	1	-	1	6	10	4	1	1	-	-	-
Female	1	-	3	5	3	7	12	6	2	3	-	-
Black, both sexes	2	-	3	3	2	1	2	6	1	-	-	-
Male	2	-	2	2	1	1	-	1	-	-	-	-
Female	-	-	1	1	1	-	2	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
South Dakota												
All races, both sexes	-	-	1	-	1	1	8	2	-	1	1	-
Male	-	-	1	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	1	1	6	2	-	-	1	-
White, both sexes	-	-	1	-	1	1	8	2	-	1	1	-
Male	-	-	1	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	1	1	6	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	9	9	10	10	10	19	21	11	6	2	-
Male	2	7	3	7	5	3	7	6	2	-	1	-
Female	1	2	6	3	5	7	12	15	9	6	1	-
White, both sexes	3	8	7	9	8	6	18	20	10	4	2	-
Male	2	7	2	7	5	1	6	6	2	-	1	-
Female	1	1	5	2	3	5	12	14	8	4	1	-
Black, both sexes	-	1	2	1	2	4	1	1	1	2	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	1	1	1	2	2	-	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Texas												
All races, both sexes	10	14	18	13	37	38	65	40	21	10	2	-
Male	5	8	2	6	20	19	24	10	10	2	-	-
Female	5	6	16	7	17	19	41	30	11	8	2	-
White, both sexes	9	12	17	12	31	33	61	35	20	10	2	-
Male	4	7	2	6	15	17	23	7	9	2	-	-
Female	5	5	15	6	16	16	38	28	11	8	2	-
Black, both sexes	1	1	1	1	2	5	4	4	1	-	-	-
Male	1	1	-	-	2	2	1	3	1	-	-	-
Female	-	-	1	1	-	3	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	4	-	-	1	-	-	-	-
Male	-	-	-	-	3	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	1	4	7	7	3	1	-	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	-	-	-	1	-	3	5	6	2	1	-	-
White, both sexes	-	-	-	1	1	4	7	7	3	1	-	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	-	-	-	1	-	3	5	6	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Vermont												
All races, both sexes	-	-	1	-	-	5	3	2	2	2	1	-
Male	-	-	-	-	-	2	1	-	1	-	-	-
Female	-	-	1	-	-	3	2	2	1	2	1	-
White, both sexes	-	-	1	-	-	5	3	2	2	2	1	-
Male	-	-	-	-	-	2	1	-	1	-	-	-
Female	-	-	1	-	-	3	2	2	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	3	4	13	10	21	34	31	14	5	-	-
Male	1	1	2	4	6	5	14	14	6	-	-	-
Female	3	2	2	9	4	16	20	17	8	5	-	-
White, both sexes	4	2	4	10	9	19	29	27	11	5	-	-
Male	1	1	2	3	6	5	11	11	3	-	-	-
Female	3	1	2	7	3	14	18	16	8	5	-	-
Black, both sexes	-	1	-	3	1	2	5	4	3	-	-	-
Male	-	-	-	1	-	-	3	3	3	-	-	-
Female	-	1	-	2	1	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Washington												
All races, both sexes	3	6	7	5	11	16	15	9	13	9	2	-
Male	1	3	1	1	4	8	5	1	2	6	1	-
Female	2	3	6	4	7	8	10	8	11	3	1	-
White, both sexes	3	5	5	4	10	15	14	8	12	9	2	-
Male	1	3	-	1	3	7	5	-	2	6	1	-
Female	2	2	5	3	7	8	9	8	10	3	1	-
Black, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	1	7	1	2	8	13	3	6	3	-	-
Male	1	1	1	-	1	1	5	-	1	1	-	-
Female	1	-	6	1	1	7	8	3	5	2	-	-
White, both sexes	2	1	7	1	2	8	13	3	6	3	-	-
Male	1	1	1	-	1	1	5	-	1	1	-	-
Female	1	-	6	1	1	7	8	3	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Wisconsin												
All races, both sexes	1	4	-	5	6	15	19	14	7	6	1	-
Male	1	3	-	2	5	7	11	5	2	1	-	-
Female	-	1	-	3	1	8	8	9	5	5	1	-
White, both sexes	1	3	-	5	6	15	18	13	6	6	1	-
Male	1	2	-	2	5	7	11	5	2	1	-	-
Female	-	1	-	3	1	8	7	8	4	5	1	-
Black, both sexes	-	1	-	-	-	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	1	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
United States												
All races, both sexes	1,829	12	4	2	3	5	4	8	16	20	29	79
Male	773	8	2	1	2	3	-	3	6	11	14	40
Female	1,056	4	2	1	1	2	4	5	10	9	15	39
White, both sexes	1,547	9	3	-	2	4	1	8	11	12	24	59
Male	655	6	1	-	1	3	-	3	4	6	12	31
Female	892	3	2	-	1	1	1	5	7	6	12	28
Black, both sexes	247	3	1	2	1	1	3	-	5	6	5	19
Male	98	2	1	1	1	-	-	-	2	3	2	8
Female	149	1	-	1	-	1	3	-	3	3	3	11
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	2	-	1
Male	16	-	-	-	-	-	-	-	-	2	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	27	-	-	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	1	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Arkansas												
All races, both sexes	16	-	1	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	11	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	131	-	-	-	-	1	-	1	1	1	4	6
Male	58	-	-	-	-	1	-	-	1	1	3	2
Female	73	-	-	-	-	-	-	1	-	-	1	4
White, both sexes	115	-	-	-	-	1	-	1	1	1	2	6
Male	50	-	-	-	-	1	-	-	1	1	2	2
Female	65	-	-	-	-	-	-	1	-	-	-	4
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Colorado												
All races, both sexes	24	-	-	-	-	-	-	-	1	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	24	-	-	-	-	-	-	-	1	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	25	-	-	-	1	-	-	-	-	-	-	-
Male	10	-	-	-	1	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Florida												
All races, both sexes	105	1	-	1	-	1	1	1	-	-	1	4
Male	41	1	-	-	-	1	-	-	-	-	1	4
Female	64	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	90	-	-	-	-	1	-	1	-	-	-	4
Male	38	-	-	-	-	1	-	-	-	-	-	4
Female	52	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	14	1	-	1	-	-	1	-	-	-	1	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	47	-	-	-	-	1	1	-	2	2	-	3
Male	21	-	-	-	-	-	-	-	1	-	-	2
Female	26	-	-	-	-	1	1	-	1	2	-	1
White, both sexes	33	-	-	-	-	-	-	-	2	2	-	1
Male	17	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	1	2	-	-
Black, both sexes	13	-	-	-	-	1	1	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	1	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Illinois												
All races, both sexes	98	-	-	-	-	-	-	-	-	3	1	7
Male	37	-	-	-	-	-	-	-	-	2	-	3
Female	61	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	79	-	-	-	-	-	-	-	-	2	1	4
Male	28	-	-	-	-	-	-	-	-	1	-	1
Female	51	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Iowa												
All races, both sexes	14	-	-	-	-	1	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	13	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	16	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Kentucky												
All races, both sexes	30	-	-	-	-	-	-	-	1	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	1	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	32	-	1	-	1	-	-	-	1	-	1	3
Male	14	-	1	-	1	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	24	-	-	-	1	-	-	-	-	-	-	2
Male	11	-	-	-	1	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	1	-	-	-	-	-	1	-	1	1
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Maine												
All races, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	5
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	36	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Massachusetts												
All races, both sexes	48	-	-	-	-	-	-	-	1	1	1	1
Male	20	-	-	-	-	-	-	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	46	-	-	-	-	-	-	-	-	1	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	75	1	-	-	-	-	-	-	1	-	2	4
Male	26	1	-	-	-	-	-	-	1	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	56	1	-	-	-	-	-	-	-	-	2	1
Male	17	1	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	16	-	-	-	-	-	-	-	1	-	-	3
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Minnesota												
All races, both sexes	35	-	-	-	-	-	-	-	1	-	2	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	34	-	-	-	-	-	-	-	1	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Missouri												
All races, both sexes	54	-	-	-	-	1	-	1	-	-	-	2
Male	24	-	-	-	-	1	-	1	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	51	-	-	-	-	1	-	1	-	-	-	2
Male	22	-	-	-	-	1	-	1	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Nebraska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
New Hampshire												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	59	1	1	-	-	-	1	-	-	2	1	2
Male	30	1	-	-	-	-	-	-	-	1	-	1
Female	29	-	1	-	-	-	1	-	-	1	1	1
White, both sexes	48	-	1	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	10	1	-	-	-	-	1	-	-	1	1	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
New Mexico												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	120	-	1	-	1	-	-	-	1	3	1	5
Male	58	-	1	-	-	-	-	-	1	3	-	5
Female	62	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	101	-	1	-	1	-	-	-	1	1	1	4
Male	45	-	1	-	-	-	-	-	1	1	-	4
Female	56	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
North Carolina												
All races, both sexes	48	2	-	-	-	-	-	-	-	1	1	-
Male	15	1	-	-	-	-	-	-	-	-	-	-
Female	33	1	-	-	-	-	-	-	-	1	1	-
White, both sexes	33	1	-	-	-	-	-	-	-	-	1	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	1	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Ohio												
All races, both sexes	97	-	-	-	-	-	-	1	-	1	1	3
Male	42	-	-	-	-	-	-	-	-	1	1	1
Female	55	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	81	-	-	-	-	-	-	1	-	-	1	2
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	50	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	26	1	-	-	-	-	1	1	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	25	1	-	-	-	-	1	1	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Oregon												
All races, both sexes	21	1	-	-	-	-	-	-	-	-	1	2
Male	10	1	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	20	1	-	-	-	-	-	-	-	-	1	2
Male	10	1	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	108	-	-	-	-	-	-	-	1	1	3	3
Male	43	-	-	-	-	-	-	-	-	1	3	1
Female	65	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	93	-	-	-	-	-	-	-	1	1	3	2
Male	39	-	-	-	-	-	-	-	-	1	3	1
Female	54	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Rhode Island												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Texas												
All races, both sexes	101	3	-	1	-	-	-	-	-	2	2	8
Male	43	-	-	1	-	-	-	-	-	-	1	6
Female	58	3	-	-	-	-	-	-	-	2	1	2
White, both sexes	85	3	-	-	-	-	-	-	-	1	2	6
Male	38	-	-	-	-	-	-	-	-	-	1	5
Female	47	3	-	-	-	-	-	-	-	1	1	1
Black, both sexes	14	-	-	1	-	-	-	-	-	1	-	2
Male	3	-	-	1	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	40	1	-	-	-	-	-	-	-	-	1	2
Male	14	1	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	31	1	-	-	-	-	-	-	-	-	1	1
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Washington												
All races, both sexes	33	-	-	-	-	-	-	-	1	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	23	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Wisconsin												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
United States												
All races, both sexes	106	114	147	154	176	243	284	247	133	35	7	1
Male	68	55	72	76	78	112	123	64	26	7	1	1
Female	38	59	75	78	98	131	161	183	107	28	6	-
White, both sexes	77	93	112	133	139	215	255	228	124	31	6	1
Male	51	46	55	66	66	101	115	58	22	6	1	1
Female	26	47	57	67	73	114	140	170	102	25	5	-
Black, both sexes	23	20	30	20	32	23	25	16	7	4	1	-
Male	12	8	14	10	10	9	7	4	3	1	-	-
Female	11	12	16	10	22	14	18	12	4	3	1	-
American Indian 1/, both sexes	3	-	1	-	1	2	1	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	4	1	4	3	3	3	2	-	-	-
Male	2	1	2	-	2	2	1	2	1	-	-	-
Female	1	-	2	1	2	1	2	1	1	-	-	-
Alabama												
All races, both sexes	4	4	2	5	2	3	5	3	3	1	1	-
Male	2	3	2	3	-	1	3	2	-	1	-	-
Female	2	1	-	2	2	2	2	1	3	-	1	-
White, both sexes	2	3	1	4	1	2	5	3	3	1	-	-
Male	2	3	1	3	-	-	3	2	-	1	-	-
Female	-	-	-	1	1	2	2	1	3	-	-	-
Black, both sexes	2	1	1	1	1	1	-	-	-	-	1	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	2	1	-	1	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Alaska												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	5	4	6	2	3	1	1	-	-	1
Male	1	1	2	3	4	1	2	-	1	-	-	1
Female	-	1	3	1	2	1	1	1	-	-	-	-
White, both sexes	-	2	5	4	5	2	3	1	1	-	-	1
Male	-	1	2	3	3	1	2	-	1	-	-	1
Female	-	1	3	1	2	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Arkansas												
All races, both sexes	-	1	3	1	1	-	3	4	1	-	-	-
Male	-	1	1	1	-	-	-	1	-	-	-	-
Female	-	-	2	-	1	-	3	3	1	-	-	-
White, both sexes	-	1	2	1	-	-	2	3	1	-	-	-
Male	-	1	1	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	2	2	1	-	-	-
Black, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10	7	11	12	11	13	23	20	7	2	1	-
Male	4	3	7	5	7	7	9	5	2	1	-	-
Female	6	4	4	7	4	6	14	15	5	1	1	-
White, both sexes	7	6	7	11	10	13	21	19	6	2	1	-
Male	3	3	4	4	7	7	9	4	1	1	-	-
Female	4	3	3	7	3	6	12	15	5	1	1	-
Black, both sexes	2	1	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	2	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	2	1	1	-	-	-
Male	1	-	2	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Colorado												
All races, both sexes	1	5	-	3	2	2	1	3	1	-	1	-
Male	-	2	-	2	1	-	-	1	1	-	-	-
Female	1	3	-	1	1	2	1	2	-	-	1	-
White, both sexes	1	5	-	3	2	2	1	3	1	-	1	-
Male	-	2	-	2	1	-	-	1	1	-	-	-
Female	1	3	-	1	1	2	1	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	2	1	2	5	4	4	4	-	-	-
Male	-	-	2	-	1	1	1	3	1	-	-	-
Female	1	1	-	1	1	4	3	1	3	-	-	-
White, both sexes	1	1	2	1	2	4	4	4	4	-	-	-
Male	-	-	2	-	1	1	1	3	1	-	-	-
Female	1	1	-	1	1	3	3	1	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Delaware												
All races, both sexes	-	2	1	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	2	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	-	2	3	1	-	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	2	1	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	2	3	1	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Florida												
All races, both sexes	4	8	4	6	8	15	19	20	10	1	-	-
Male	1	4	1	1	5	7	10	3	1	1	-	-
Female	3	4	3	5	3	8	9	17	9	-	-	-
White, both sexes	3	7	3	4	6	13	19	18	10	1	-	-
Male	1	4	-	1	5	7	10	3	1	1	-	-
Female	2	3	3	3	1	6	9	15	9	-	-	-
Black, both sexes	-	1	1	2	2	2	-	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	2	2	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	2	1	2	7	3	9	8	1	2	-	-
Male	2	2	-	1	5	3	3	-	1	1	-	-
Female	1	-	1	1	2	-	6	8	-	1	-	-
White, both sexes	1	2	-	2	6	3	7	6	1	-	-	-
Male	1	2	-	1	4	3	3	-	1	-	-	-
Female	-	-	-	1	2	-	4	6	-	-	-	-
Black, both sexes	2	-	1	-	1	-	2	2	-	2	-	-
Male	1	-	-	-	1	-	-	-	-	1	-	-
Female	1	-	1	-	-	-	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Hawaii												
All races, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Idaho												
All races, both sexes	-	1	1	-	2	-	2	2	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	2	-	-	-	-
White, both sexes	-	1	1	-	2	-	2	2	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Illinois												
All races, both sexes	4	7	6	8	10	10	15	11	16	-	-	-
Male	4	3	2	3	2	3	9	2	4	-	-	-
Female	-	4	4	5	8	7	6	9	12	-	-	-
White, both sexes	2	6	3	6	7	8	15	10	15	-	-	-
Male	2	3	-	2	2	3	9	1	4	-	-	-
Female	-	3	3	4	5	5	6	9	11	-	-	-
Black, both sexes	2	1	3	1	3	2	-	1	1	-	-	-
Male	2	-	2	1	-	-	-	1	-	-	-	-
Female	-	1	1	-	3	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	6	6	4	12	9	3	3	-	-	-
Male	-	1	1	3	1	7	4	1	-	-	-	-
Female	2	-	5	3	3	5	5	2	3	-	-	-
White, both sexes	1	1	5	5	3	11	9	3	3	-	-	-
Male	-	1	1	3	-	6	4	1	-	-	-	-
Female	1	-	4	2	3	5	5	2	3	-	-	-
Black, both sexes	1	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Iowa												
All races, both sexes	2	-	1	-	1	1	3	2	2	-	-	-
Male	2	-	1	-	1	-	3	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
White, both sexes	2	-	1	-	1	1	3	2	2	-	-	-
Male	2	-	1	-	1	-	3	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	1	1	5	2	2	2	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	-	-	-	-	1	3	1	1	2	-	-	-
White, both sexes	-	1	-	1	1	4	2	2	2	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	-	-	-	-	1	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Kentucky												
All races, both sexes	-	3	3	2	4	8	2	5	1	-	-	-
Male	-	2	1	2	1	3	-	2	-	-	-	-
Female	-	1	2	-	3	5	2	3	1	-	-	-
White, both sexes	-	1	3	1	3	7	2	5	1	-	-	-
Male	-	-	1	1	1	3	-	2	-	-	-	-
Female	-	1	2	-	2	4	2	3	1	-	-	-
Black, both sexes	-	1	-	1	1	1	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	3	2	2	2	5	5	2	2	1	-	-
Male	-	3	2	1	-	4	1	-	-	-	-	-
Female	1	-	-	1	2	1	4	2	2	1	-	-
White, both sexes	1	2	1	2	1	5	5	2	2	-	-	-
Male	-	2	1	1	-	4	1	-	-	-	-	-
Female	1	-	-	1	1	1	4	2	2	-	-	-
Black, both sexes	-	1	1	-	1	-	-	-	-	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Maine												
All races, both sexes	2	-	1	1	1	3	2	2	-	1	-	-
Male	2	-	-	1	1	2	1	2	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	1	-	-
White, both sexes	2	-	1	1	1	3	2	2	-	1	-	-
Male	2	-	-	1	1	2	1	2	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	1	4	4	4	6	6	11	4	2	-	-
Male	2	-	3	-	-	1	1	2	1	2	-	-
Female	1	1	1	4	4	5	5	9	3	-	-	-
White, both sexes	1	-	3	3	2	5	4	10	2	2	-	-
Male	1	-	2	-	-	1	-	2	-	2	-	-
Female	-	-	1	3	2	4	4	8	2	-	-	-
Black, both sexes	1	1	1	1	1	1	2	1	2	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	1	1	-	1	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Massachusetts												
All races, both sexes	2	3	2	4	3	14	7	4	2	3	-	-
Male	1	2	1	2	-	8	3	-	1	-	-	-
Female	1	1	1	2	3	6	4	4	1	3	-	-
White, both sexes	2	2	2	4	3	14	7	4	2	3	-	-
Male	1	1	1	2	-	8	3	-	1	-	-	-
Female	1	1	1	2	3	6	4	4	1	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	5	5	4	6	11	17	12	4	-	-	-
Male	2	4	3	-	4	4	5	1	-	-	-	-
Female	1	1	2	4	2	7	12	11	4	-	-	-
White, both sexes	-	3	5	4	4	9	14	9	4	-	-	-
Male	-	3	3	-	2	4	4	-	-	-	-	-
Female	-	-	2	4	2	5	10	9	4	-	-	-
Black, both sexes	2	2	-	-	2	2	3	1	-	-	-	-
Male	1	1	-	-	2	-	1	-	-	-	-	-
Female	1	1	-	-	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Minnesota												
All races, both sexes	6	3	3	1	-	3	6	5	4	1	-	-
Male	5	2	2	1	-	2	3	2	2	-	-	-
Female	1	1	1	-	-	1	3	3	2	1	-	-
White, both sexes	5	3	3	1	-	3	6	5	4	1	-	-
Male	4	2	2	1	-	2	3	2	2	-	-	-
Female	1	1	1	-	-	1	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	1	3	1	2	5	1	2	-	-	-	-
Male	2	-	3	1	1	1	-	1	-	-	-	-
Female	-	1	-	-	1	4	1	1	-	-	-	-
White, both sexes	2	-	1	1	2	3	1	2	-	-	-	-
Male	2	-	1	1	1	1	-	1	-	-	-	-
Female	-	-	-	-	1	2	1	1	-	-	-	-
Black, both sexes	-	1	2	-	-	2	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Missouri												
All races, both sexes	5	4	6	3	8	4	7	5	6	1	1	-
Male	4	3	4	-	4	1	3	-	1	-	1	-
Female	1	1	2	3	4	3	4	5	5	1	-	-
White, both sexes	5	3	6	2	7	4	7	5	6	1	1	-
Male	4	2	4	-	3	1	3	-	1	-	1	-
Female	1	1	2	2	4	3	4	5	5	1	-	-
Black, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Nebraska												
All races, both sexes	-	1	-	-	1	2	2	-	1	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	1	-	-
White, both sexes	-	1	-	-	1	2	2	-	1	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	2	1	2	2	3	4	-	-	-	-
Male	-	-	2	-	2	1	2	2	-	-	-	-
Female	-	2	-	1	-	1	1	2	-	-	-	-
White, both sexes	-	2	1	1	2	-	3	4	-	-	-	-
Male	-	-	1	-	2	-	2	2	-	-	-	-
Female	-	2	-	1	-	-	1	2	-	-	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
New Hampshire												
All races, both sexes	-	2	2	-	-	3	2	1	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	2	1	-	-	2	1	1	-	-	-	-
White, both sexes	-	2	2	-	-	3	2	1	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	2	1	-	-	2	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	5	5	7	8	6	11	5	1	-	-
Male	1	2	3	3	6	2	4	5	1	-	-	-
Female	-	-	2	2	1	6	2	6	4	1	-	-
White, both sexes	1	2	4	5	5	7	5	11	4	1	-	-
Male	1	2	3	3	5	2	3	5	-	-	-	-
Female	-	-	1	2	-	5	2	6	4	1	-	-
Black, both sexes	-	-	1	-	2	1	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
New Mexico												
All races, both sexes	1	1	2	1	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
White, both sexes	1	1	1	1	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	7	8	11	9	17	18	20	10	1	-	-
Male	5	4	4	5	3	11	7	7	2	-	-	-
Female	2	3	4	6	6	6	11	13	8	1	-	-
White, both sexes	4	6	8	8	7	15	15	19	9	1	-	-
Male	3	3	4	2	3	9	6	6	2	-	-	-
Female	1	3	4	6	4	6	9	13	7	1	-	-
Black, both sexes	3	1	-	3	2	2	3	1	1	-	-	-
Male	2	1	-	3	-	2	1	1	-	-	-	-
Female	1	-	-	-	2	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
North Carolina												
All races, both sexes	5	5	5	3	5	7	5	3	3	3	-	-
Male	3	1	-	2	1	4	2	-	1	-	-	-
Female	2	4	5	1	4	3	3	3	2	3	-	-
White, both sexes	2	3	3	3	4	6	3	2	2	3	-	-
Male	1	-	-	2	1	3	1	-	-	-	-	-
Female	1	3	3	1	3	3	2	2	2	3	-	-
Black, both sexes	3	2	2	-	1	1	2	1	1	-	-	-
Male	2	1	-	-	-	1	1	-	1	-	-	-
Female	1	1	2	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
White, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Ohio												
All races, both sexes	11	2	9	11	6	11	19	13	7	2	-	-
Male	7	-	3	9	2	3	12	2	1	-	-	-
Female	4	2	6	2	4	8	7	11	6	2	-	-
White, both sexes	7	2	7	10	5	9	16	12	7	2	-	-
Male	4	-	2	8	1	2	10	2	1	-	-	-
Female	3	2	5	2	4	7	6	10	6	2	-	-
Black, both sexes	4	-	1	1	1	2	3	1	-	-	-	-
Male	3	-	1	1	1	1	2	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	4	1	7	1	-	5	4	1	-	-	-
Male	-	-	-	5	-	-	3	-	-	-	-	-
Female	-	4	1	2	1	-	2	4	1	-	-	-
White, both sexes	-	4	1	6	1	-	5	4	1	-	-	-
Male	-	-	-	4	-	-	3	-	-	-	-	-
Female	-	4	1	2	1	-	2	4	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Oregon												
All races, both sexes	-	1	2	2	1	1	3	3	1	3	-	-
Male	-	-	1	1	1	1	2	1	1	-	-	-
Female	-	1	1	1	-	-	1	2	-	3	-	-
White, both sexes	-	1	2	2	1	1	2	3	1	3	-	-
Male	-	-	1	1	1	1	2	1	1	-	-	-
Female	-	1	1	1	-	-	-	2	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	3	9	12	12	17	19	11	9	2	2	-
Male	2	1	2	5	5	12	8	2	1	-	-	-
Female	2	2	7	7	7	5	11	9	8	2	2	-
White, both sexes	4	2	4	10	10	15	19	10	9	1	2	-
Male	2	1	1	4	5	10	8	2	1	-	-	-
Female	2	1	3	6	5	5	11	8	8	1	2	-
Black, both sexes	-	1	4	2	2	2	-	1	-	1	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	1	3	1	2	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Rhode Island												
All races, both sexes	2	-	-	1	-	1	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	2	-	-	1	-	1	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	4	1	5	4	3	4	3	3	-	-	-
Male	2	2	1	1	1	1	1	-	-	-	-	-
Female	-	2	-	4	3	2	3	3	3	-	-	-
White, both sexes	1	2	1	2	3	3	3	3	3	-	-	-
Male	1	2	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	2	3	2	2	3	3	-	-	-
Black, both sexes	1	2	-	3	1	-	1	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	-	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
South Dakota												
All races, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	3	5	3	4	7	5	4	2	-	-	-
Male	1	2	3	1	2	2	1	2	1	-	-	-
Female	2	1	2	2	2	5	4	2	1	-	-	-
White, both sexes	3	3	4	3	4	6	4	4	2	-	-	-
Male	1	2	2	1	2	2	1	2	1	-	-	-
Female	2	1	2	2	2	4	3	2	1	-	-	-
Black, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Texas												
All races, both sexes	5	5	9	6	14	15	17	6	7	1	-	-
Male	5	2	4	3	7	5	6	2	1	-	-	-
Female	-	3	5	3	7	10	11	4	6	1	-	-
White, both sexes	5	4	7	6	8	14	15	6	7	1	-	-
Male	5	1	4	3	6	4	6	2	1	-	-	-
Female	-	3	3	3	2	10	9	4	6	1	-	-
Black, both sexes	-	1	2	-	5	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	5	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	2	2	1	1	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	2	1	1	1	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Vermont												
All races, both sexes	-	1	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	1	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	2	7	3	6	1	4	7	3	1	-	-
Male	1	1	3	1	3	-	-	2	1	-	-	-
Female	1	1	4	2	3	1	4	5	2	1	-	-
White, both sexes	2	2	5	2	4	1	2	6	3	1	-	-
Male	1	1	2	1	2	-	-	2	1	-	-	-
Female	1	1	3	1	2	1	2	4	2	1	-	-
Black, both sexes	-	-	2	1	2	-	2	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Washington												
All races, both sexes	2	1	5	2	2	3	4	4	4	1	1	-
Male	-	-	2	-	-	2	2	-	-	-	-	-
Female	2	1	3	2	2	1	2	4	4	1	1	-
White, both sexes	2	1	5	2	2	3	4	4	4	1	1	-
Male	-	-	2	-	-	2	2	-	-	-	-	-
Female	2	1	3	2	2	1	2	4	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	1	4	2	3	7	-	2	-	-
Male	2	-	-	1	1	1	2	3	-	-	-	-
Female	-	-	-	-	3	1	1	4	-	2	-	-
White, both sexes	2	-	-	1	4	2	3	7	-	2	-	-
Male	2	-	-	1	1	1	2	3	-	-	-	-
Female	-	-	-	-	3	1	1	4	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Wisconsin												
All races, both sexes	2	2	2	5	6	5	7	9	-	1	-	-
Male	2	2	2	4	2	3	2	4	-	1	-	-
Female	-	-	-	1	4	2	5	5	-	-	-	-
White, both sexes	2	2	1	5	6	5	7	9	-	1	-	-
Male	2	2	1	4	2	3	2	4	-	1	-	-
Female	-	-	-	1	4	2	5	5	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
United States												
All races, both sexes	36,783	96	27	15	15	27	56	138	346	925	2,469	4,261
Male	23,284	61	15	8	9	6	30	79	213	594	1,598	2,967
Female	13,499	35	12	7	6	21	26	59	133	331	871	1,294
White, both sexes	31,708	67	16	10	6	17	39	103	255	772	2,069	3,540
Male	20,175	40	9	5	3	5	20	62	168	510	1,361	2,486
Female	11,533	27	7	5	3	12	19	41	87	262	708	1,054
Black, both sexes	3,734	26	8	3	8	7	14	20	52	90	260	556
Male	2,318	19	4	1	5	1	8	7	21	47	151	372
Female	1,416	7	4	2	3	6	6	13	31	43	109	184
American Indian 1/, both sexes	746	-	2	-	1	2	2	12	29	46	106	115
Male	431	-	1	-	1	-	2	7	17	23	59	68
Female	315	-	1	-	-	2	-	5	12	23	47	47
Asian or Pacific Islander 2/, both sexes	595	3	1	2	-	1	1	3	10	17	34	50
Male	360	2	1	2	-	-	-	3	7	14	27	41
Female	235	1	-	-	-	1	1	-	3	3	7	9
Alabama												
All races, both sexes	674	1	-	1	-	2	-	2	10	16	43	77
Male	411	-	-	1	-	-	-	-	5	9	27	52
Female	263	1	-	-	-	2	-	2	5	7	16	25
White, both sexes	525	-	-	1	-	-	-	1	6	11	28	56
Male	320	-	-	1	-	-	-	-	4	8	16	40
Female	205	-	-	-	-	-	-	1	2	3	12	16
Black, both sexes	146	1	-	-	-	2	-	1	4	5	13	21
Male	89	-	-	-	-	-	-	-	1	1	9	12
Female	57	1	-	-	-	2	-	1	3	4	4	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Alaska												
All races, both sexes	71	-	-	-	-	-	-	-	-	3	10	7
Male	40	-	-	-	-	-	-	-	-	1	3	1
Female	31	-	-	-	-	-	-	-	-	2	7	6
White, both sexes	53	-	-	-	-	-	-	-	-	3	7	3
Male	33	-	-	-	-	-	-	-	-	1	3	1
Female	20	-	-	-	-	-	-	-	-	2	4	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	3	4
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	3	4
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	940	3	1	-	-	-	1	4	14	34	69	107
Male	606	2	-	-	-	-	1	3	11	23	44	65
Female	334	1	1	-	-	-	-	1	3	11	25	42
White, both sexes	795	3	-	-	-	-	1	-	7	25	53	79
Male	525	2	-	-	-	-	1	-	7	18	33	50
Female	270	1	-	-	-	-	-	-	-	7	20	29
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	123	-	1	-	-	-	-	4	6	9	15	23
Male	67	-	-	-	-	-	-	3	3	5	11	11
Female	56	-	1	-	-	-	-	1	3	4	4	12
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	3
Male	5	-	-	-	-	-	-	-	1	-	-	3
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Arkansas												
All races, both sexes	345	-	1	-	-	-	2	-	5	8	16	40
Male	203	-	1	-	-	-	2	-	2	3	6	27
Female	142	-	-	-	-	-	-	-	3	5	10	13
White, both sexes	308	-	1	-	-	-	-	-	4	7	13	36
Male	185	-	1	-	-	-	-	-	2	2	5	26
Female	123	-	-	-	-	-	-	-	2	5	8	10
Black, both sexes	33	-	-	-	-	-	2	-	1	1	2	4
Male	15	-	-	-	-	-	2	-	-	1	-	1
Female	18	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,587	14	2	2	2	3	6	25	48	148	330	553
Male	3,083	10	1	1	1	-	3	18	33	106	227	403
Female	1,504	4	1	1	1	3	3	7	15	42	103	150
White, both sexes	4,025	13	2	2	2	2	6	21	39	135	286	482
Male	2,728	9	1	1	1	-	3	16	26	98	199	347
Female	1,297	4	1	1	1	2	3	5	13	37	87	135
Black, both sexes	277	-	-	-	-	-	-	2	2	5	21	44
Male	177	-	-	-	-	-	-	1	2	2	12	33
Female	100	-	-	-	-	-	-	1	-	3	9	11
American Indian 1/, both sexes	70	-	-	-	-	-	-	1	3	3	9	8
Male	48	-	-	-	-	-	-	-	3	2	5	7
Female	22	-	-	-	-	-	-	1	-	1	4	1
Asian or Pacific Islander 2/, both sexes	215	1	-	-	-	1	-	1	4	5	14	19
Male	130	1	-	-	-	-	-	1	2	4	11	16
Female	85	-	-	-	-	1	-	-	2	1	3	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Colorado												
All races, both sexes	619	1	2	-	1	-	-	2	14	20	43	98
Male	405	1	2	-	1	-	-	2	10	10	33	70
Female	214	-	-	-	-	-	-	-	4	10	10	28
White, both sexes	584	1	1	-	1	-	-	2	13	19	42	92
Male	384	1	1	-	1	-	-	2	9	10	32	67
Female	200	-	-	-	-	-	-	-	4	9	10	25
Black, both sexes	17	-	1	-	-	-	-	-	-	1	1	4
Male	11	-	1	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	1	-	-	2
Male	8	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	432	1	1	-	-	-	1	2	2	7	28	35
Male	253	1	1	-	-	-	1	2	1	3	19	26
Female	179	-	-	-	-	-	-	-	1	4	9	9
White, both sexes	399	1	1	-	-	-	1	2	1	6	25	35
Male	235	1	1	-	-	-	1	2	1	3	17	26
Female	164	-	-	-	-	-	-	-	-	3	8	9
Black, both sexes	31	-	-	-	-	-	-	-	1	-	3	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	14	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Delaware												
All races, both sexes	96	-	-	-	-	-	-	1	-	-	7	9
Male	63	-	-	-	-	-	-	1	-	-	4	6
Female	33	-	-	-	-	-	-	-	-	-	3	3
White, both sexes	89	-	-	-	-	-	-	1	-	-	7	9
Male	60	-	-	-	-	-	-	1	-	-	4	6
Female	29	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	89	-	1	-	-	-	-	1	1	1	6	8
Male	62	-	1	-	-	-	-	-	1	1	3	6
Female	27	-	-	-	-	-	-	1	-	-	3	2
White, both sexes	16	-	-	-	-	-	-	1	1	-	1	1
Male	10	-	-	-	-	-	-	-	1	-	1	1
Female	6	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	69	-	-	-	-	-	-	-	-	1	5	7
Male	49	-	-	-	-	-	-	-	-	1	2	5
Female	20	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Florida												
All races, both sexes	2,680	7	-	-	-	2	4	5	12	44	148	315
Male	1,724	5	-	-	-	2	2	4	7	26	94	213
Female	956	2	-	-	-	-	2	1	5	18	54	102
White, both sexes	2,446	3	-	-	-	2	1	3	9	39	136	276
Male	1,575	1	-	-	-	2	-	3	5	23	84	187
Female	871	2	-	-	-	-	1	-	4	16	52	89
Black, both sexes	216	4	-	-	-	-	3	2	3	4	10	39
Male	133	4	-	-	-	-	2	1	2	2	8	26
Female	83	-	-	-	-	-	1	1	1	2	2	13
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	2	-
Male	11	-	-	-	-	-	-	-	-	1	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,063	7	1	1	2	-	3	4	8	31	74	124
Male	670	5	1	-	1	-	2	1	6	18	50	86
Female	393	2	-	1	1	-	1	3	2	13	24	38
White, both sexes	793	1	-	1	-	-	3	4	2	20	59	84
Male	512	1	-	-	-	-	2	1	2	11	39	59
Female	281	-	-	1	-	-	1	3	-	9	20	25
Black, both sexes	256	6	-	-	2	-	-	-	6	10	13	39
Male	149	4	-	-	1	-	-	-	4	6	9	27
Female	107	2	-	-	1	-	-	-	2	4	4	12
American Indian 1/, both sexes	3	-	1	-	-	-	-	-	-	-	1	1
Male	2	-	1	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Hawaii												
All races, both sexes	147	1	-	1	-	-	-	2	1	1	8	13
Male	89	-	-	1	-	-	-	2	1	1	4	9
Female	58	1	-	-	-	-	-	-	-	-	4	4
White, both sexes	62	-	-	-	-	-	-	1	-	-	2	6
Male	39	-	-	-	-	-	-	1	-	-	-	4
Female	23	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	82	1	-	1	-	-	-	1	1	1	6	7
Male	49	-	-	1	-	-	-	1	1	1	4	5
Female	33	1	-	-	-	-	-	-	-	-	2	2
Idaho												
All races, both sexes	144	1	-	-	-	-	-	2	1	5	10	15
Male	94	1	-	-	-	-	-	2	1	1	5	11
Female	50	-	-	-	-	-	-	-	-	4	5	4
White, both sexes	136	1	-	-	-	-	-	2	1	3	8	15
Male	92	1	-	-	-	-	-	2	1	1	5	11
Female	44	-	-	-	-	-	-	-	-	2	3	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	2	2	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Illinois												
All races, both sexes	1,367	1	-	2	1	-	4	12	16	35	113	156
Male	838	1	-	2	1	-	2	8	12	24	80	112
Female	529	-	-	-	-	-	2	4	4	11	33	44
White, both sexes	1,141	1	-	-	-	-	2	10	15	32	92	117
Male	697	1	-	-	-	-	2	7	12	22	67	83
Female	444	-	-	-	-	-	-	3	3	10	25	34
Black, both sexes	200	-	-	1	1	-	2	2	-	2	18	36
Male	128	-	-	1	1	-	-	1	-	1	11	27
Female	72	-	-	-	-	-	2	1	-	1	7	9
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	16	-	-	1	-	-	-	-	1	-	2	1
Male	8	-	-	1	-	-	-	-	-	-	2	1
Female	8	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	693	4	2	-	1	-	2	3	8	19	38	81
Male	422	2	-	-	1	-	-	2	6	14	24	52
Female	271	2	2	-	-	-	2	1	2	5	14	29
White, both sexes	633	3	2	-	-	-	2	3	7	19	33	78
Male	386	1	-	-	-	-	-	2	5	14	20	51
Female	247	2	2	-	-	-	2	1	2	5	13	27
Black, both sexes	55	1	-	-	1	-	-	-	1	-	4	2
Male	34	1	-	-	1	-	-	-	1	-	3	1
Female	21	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Iowa												
All races, both sexes	261	-	1	-	-	-	-	-	3	7	10	34
Male	163	-	1	-	-	-	-	-	2	4	8	24
Female	98	-	-	-	-	-	-	-	1	3	2	10
White, both sexes	251	-	1	-	-	-	-	-	3	7	10	32
Male	156	-	1	-	-	-	-	-	2	4	8	23
Female	95	-	-	-	-	-	-	-	1	3	2	9
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	295	-	-	-	-	-	-	-	1	2	23	45
Male	190	-	-	-	-	-	-	-	1	2	19	31
Female	105	-	-	-	-	-	-	-	-	-	4	14
White, both sexes	276	-	-	-	-	-	-	-	1	2	21	41
Male	181	-	-	-	-	-	-	-	1	2	17	29
Female	95	-	-	-	-	-	-	-	-	-	4	12
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	3
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Kentucky												
All races, both sexes	537	1	-	-	-	-	-	1	5	14	33	61
Male	347	1	-	-	-	-	-	-	3	12	26	46
Female	190	-	-	-	-	-	-	1	2	2	7	15
White, both sexes	509	1	-	-	-	-	-	-	5	14	30	60
Male	328	1	-	-	-	-	-	-	3	12	24	46
Female	181	-	-	-	-	-	-	-	2	2	6	14
Black, both sexes	26	-	-	-	-	-	-	1	-	-	3	1
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	592	1	-	1	-	1	1	3	11	11	43	74
Male	363	-	-	-	-	-	1	1	7	5	29	50
Female	229	1	-	1	-	1	-	2	4	6	14	24
White, both sexes	422	1	-	-	-	-	-	1	5	3	27	49
Male	261	-	-	-	-	-	-	1	4	1	18	31
Female	161	1	-	-	-	-	-	-	1	2	9	18
Black, both sexes	165	-	-	1	-	1	1	2	6	8	16	24
Male	99	-	-	-	-	-	1	-	3	4	11	18
Female	66	-	-	1	-	1	-	2	3	4	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Maine												
All races, both sexes	159	-	-	-	-	-	-	-	1	2	5	13
Male	109	-	-	-	-	-	-	-	1	-	3	7
Female	50	-	-	-	-	-	-	-	-	2	2	6
White, both sexes	157	-	-	-	-	-	-	-	1	2	5	13
Male	108	-	-	-	-	-	-	-	1	-	3	7
Female	49	-	-	-	-	-	-	-	-	2	2	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	605	-	-	1	-	-	1	2	3	20	35	65
Male	384	-	-	1	-	-	1	1	1	10	18	44
Female	221	-	-	-	-	-	-	1	2	10	17	21
White, both sexes	430	-	-	1	-	-	1	1	2	13	25	40
Male	268	-	-	1	-	-	1	-	1	7	14	28
Female	162	-	-	-	-	-	-	1	1	6	11	12
Black, both sexes	163	-	-	-	-	-	-	1	1	6	9	25
Male	112	-	-	-	-	-	-	1	-	3	3	16
Female	51	-	-	-	-	-	-	-	1	3	6	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Massachusetts												
All races, both sexes	797	-	-	1	-	-	1	1	8	18	57	85
Male	491	-	-	1	-	-	-	1	5	10	35	57
Female	306	-	-	-	-	-	1	-	3	8	22	28
White, both sexes	759	-	-	1	-	-	1	1	8	18	57	78
Male	468	-	-	1	-	-	-	1	5	10	35	52
Female	291	-	-	-	-	-	1	-	3	8	22	26
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	7
Male	19	-	-	-	-	-	-	-	-	-	-	5
Female	11	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,334	2	3	-	1	-	2	3	21	41	101	140
Male	841	2	2	-	1	-	-	1	10	25	64	97
Female	493	-	1	-	-	-	2	2	11	16	37	43
White, both sexes	1,109	1	1	-	-	-	2	2	14	36	83	116
Male	692	1	1	-	-	-	-	1	7	23	51	82
Female	417	-	-	-	-	-	2	1	7	13	32	34
Black, both sexes	202	1	2	-	-	-	-	1	5	4	15	23
Male	132	1	1	-	-	-	-	-	2	1	11	14
Female	70	-	1	-	-	-	-	1	3	3	4	9
American Indian 1/, both sexes	18	-	-	-	1	-	-	-	2	1	3	1
Male	14	-	-	-	1	-	-	-	1	1	2	1
Female	4	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Minnesota												
All races, both sexes	431	2	-	-	-	1	2	1	6	9	30	50
Male	256	2	-	-	-	-	1	1	3	4	17	34
Female	175	-	-	-	-	1	1	-	3	5	13	16
White, both sexes	397	2	-	-	-	1	1	1	3	8	24	45
Male	236	2	-	-	-	-	1	1	2	3	13	30
Female	161	-	-	-	-	1	-	-	1	5	11	15
Black, both sexes	13	-	-	-	-	-	-	-	1	1	2	1
Male	7	-	-	-	-	-	-	-	-	1	2	-
Female	6	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	2	-	4	3
Male	9	-	-	-	-	-	-	-	1	-	2	3
Female	6	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	422	1	-	-	-	-	3	1	8	16	20	58
Male	264	1	-	-	-	-	2	-	2	9	11	42
Female	158	-	-	-	-	-	1	1	6	7	9	16
White, both sexes	314	1	-	-	-	-	3	1	4	9	13	36
Male	196	1	-	-	-	-	2	-	2	5	8	27
Female	118	-	-	-	-	-	1	1	2	4	5	9
Black, both sexes	100	-	-	-	-	-	-	-	3	7	6	20
Male	66	-	-	-	-	-	-	-	-	4	3	14
Female	34	-	-	-	-	-	-	-	3	3	3	6
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Missouri												
All races, both sexes	611	3	1	-	-	-	1	1	6	10	38	81
Male	380	2	1	-	-	-	-	1	3	5	20	59
Female	231	1	-	-	-	-	1	-	3	5	18	22
White, both sexes	545	1	1	-	-	-	1	1	6	10	34	73
Male	343	1	1	-	-	-	-	1	3	5	18	53
Female	202	-	-	-	-	-	1	-	3	5	16	20
Black, both sexes	60	2	-	-	-	-	-	-	-	-	3	8
Male	34	1	-	-	-	-	-	-	-	-	1	6
Female	26	1	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	146	-	-	-	-	-	-	2	3	5	14	12
Male	87	-	-	-	-	-	-	-	1	2	10	8
Female	59	-	-	-	-	-	-	2	2	3	4	4
White, both sexes	108	-	-	-	-	-	-	-	-	1	7	8
Male	70	-	-	-	-	-	-	-	-	-	5	5
Female	38	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	37	-	-	-	-	-	-	2	3	4	7	3
Male	17	-	-	-	-	-	-	-	1	2	5	3
Female	20	-	-	-	-	-	-	2	2	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Nebraska												
All races, both sexes	199	-	-	-	-	-	-	-	3	2	25	27
Male	124	-	-	-	-	-	-	-	1	-	15	19
Female	75	-	-	-	-	-	-	-	2	2	10	8
White, both sexes	179	-	-	-	-	-	-	-	2	2	23	23
Male	112	-	-	-	-	-	-	-	1	-	14	17
Female	67	-	-	-	-	-	-	-	1	2	9	6
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	1	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	402	-	-	-	-	1	-	1	3	12	35	54
Male	267	-	-	-	-	-	-	1	1	8	19	39
Female	135	-	-	-	-	1	-	-	2	4	16	15
White, both sexes	368	-	-	-	-	1	-	1	1	12	33	48
Male	248	-	-	-	-	-	-	1	1	8	17	34
Female	120	-	-	-	-	1	-	-	-	4	16	14
Black, both sexes	18	-	-	-	-	-	-	-	2	-	1	5
Male	11	-	-	-	-	-	-	-	-	-	1	5
Female	7	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
New Hampshire												
All races, both sexes	147	1	-	-	-	-	-	-	1	2	14	12
Male	90	-	-	-	-	-	-	-	1	1	10	6
Female	57	1	-	-	-	-	-	-	-	1	4	6
White, both sexes	145	1	-	-	-	-	-	-	1	2	13	12
Male	89	-	-	-	-	-	-	-	1	1	9	6
Female	56	1	-	-	-	-	-	-	-	1	4	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	994	3	-	-	1	-	2	2	8	30	72	102
Male	626	1	-	-	-	-	1	-	7	22	49	74
Female	368	2	-	-	1	-	1	2	1	8	23	28
White, both sexes	839	2	-	-	1	-	1	1	7	26	60	84
Male	536	1	-	-	-	-	1	-	6	20	41	61
Female	303	1	-	-	1	-	-	1	1	6	19	23
Black, both sexes	127	1	-	-	-	-	1	1	1	4	11	15
Male	69	-	-	-	-	-	-	-	1	2	7	10
Female	58	1	-	-	-	-	1	1	-	2	4	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	3
Male	19	-	-	-	-	-	-	-	-	-	1	3
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
New Mexico												
All races, both sexes	395	1	-	-	-	1	-	5	7	18	33	52
Male	253	-	-	-	-	-	-	4	5	9	20	41
Female	142	1	-	-	-	1	-	1	2	9	13	11
White, both sexes	296	1	-	-	-	1	-	3	3	12	19	36
Male	199	-	-	-	-	-	-	2	2	8	14	29
Female	97	1	-	-	-	1	-	1	1	4	5	7
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	96	-	-	-	-	-	-	2	4	6	13	16
Male	54	-	-	-	-	-	-	2	3	1	6	12
Female	42	-	-	-	-	-	-	-	1	5	7	4
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,661	6	1	-	1	2	3	6	10	28	84	154
Male	1,072	5	-	-	1	-	2	3	7	20	60	117
Female	589	1	1	-	-	2	1	3	3	8	24	37
White, both sexes	1,381	5	1	-	-	1	3	5	9	20	75	124
Male	884	4	-	-	-	-	2	2	6	14	55	98
Female	497	1	1	-	-	1	1	3	3	6	20	26
Black, both sexes	218	1	-	-	1	1	-	-	1	5	8	21
Male	146	1	-	-	1	-	-	-	1	3	5	13
Female	72	-	-	-	-	1	-	-	-	2	3	8
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	1	1	4
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	7	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	48	-	-	-	-	-	-	1	-	2	-	5
Male	35	-	-	-	-	-	-	1	-	2	-	4
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
North Carolina												
All races, both sexes	1,081	5	1	-	-	4	2	3	10	19	63	113
Male	688	3	-	-	-	2	1	1	4	13	38	79
Female	393	2	1	-	-	2	1	2	6	6	25	34
White, both sexes	857	4	-	-	-	3	1	3	6	14	44	87
Male	536	2	-	-	-	1	-	1	3	9	26	60
Female	321	2	-	-	-	2	1	2	3	5	18	27
Black, both sexes	194	1	1	-	-	1	-	-	3	4	17	26
Male	132	1	-	-	-	1	-	-	1	3	10	19
Female	62	-	1	-	-	-	-	-	2	1	7	7
American Indian 1/, both sexes	26	-	-	-	-	-	1	-	1	1	2	-
Male	18	-	-	-	-	-	1	-	-	1	2	-
Female	8	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	81	-	-	-	-	-	-	-	1	2	9	9
Male	46	-	-	-	-	-	-	-	1	1	5	4
Female	35	-	-	-	-	-	-	-	-	1	4	5
White, both sexes	66	-	-	-	-	-	-	-	-	-	6	8
Male	37	-	-	-	-	-	-	-	-	-	4	3
Female	29	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	1	2	3	1
Male	9	-	-	-	-	-	-	-	1	1	1	1
Female	6	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Ohio												
All races, both sexes	1,447	4	1	-	1	1	-	5	8	40	88	167
Male	888	4	1	-	-	-	-	1	4	22	55	112
Female	559	-	-	-	1	1	-	4	4	18	33	55
White, both sexes	1,275	3	1	-	1	-	-	3	6	36	74	136
Male	778	3	1	-	-	-	-	1	4	20	45	91
Female	497	-	-	-	1	-	-	2	2	16	29	45
Black, both sexes	163	1	-	-	-	1	-	2	2	2	11	31
Male	104	1	-	-	-	-	-	-	-	1	7	21
Female	59	-	-	-	-	1	-	2	2	1	4	10
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	2	2	-
Male	3	-	-	-	-	-	-	-	-	1	2	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	560	1	2	1	1	3	-	3	3	10	55	64
Male	339	1	1	-	1	-	-	1	2	8	31	41
Female	221	-	1	1	-	3	-	2	1	2	24	23
White, both sexes	441	-	1	-	1	1	-	1	3	6	34	45
Male	269	-	-	-	1	-	-	-	2	6	22	31
Female	172	-	1	-	-	1	-	1	1	-	12	14
Black, both sexes	33	1	1	1	-	-	-	1	-	-	5	6
Male	19	1	1	-	-	-	-	1	-	-	2	4
Female	14	-	-	1	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	86	-	-	-	-	2	-	1	-	4	16	13
Male	51	-	-	-	-	-	-	-	-	2	7	6
Female	35	-	-	-	-	2	-	1	-	2	9	7
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Oregon												
All races, both sexes	470	-	-	-	-	-	1	1	5	17	21	68
Male	293	-	-	-	-	-	1	1	1	9	12	44
Female	177	-	-	-	-	-	-	-	4	8	9	24
White, both sexes	441	-	-	-	-	-	1	-	5	17	19	62
Male	278	-	-	-	-	-	1	-	1	9	11	40
Female	163	-	-	-	-	-	-	-	4	8	8	22
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	1	-	-	1	5
Male	10	-	-	-	-	-	-	1	-	-	1	3
Female	5	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
Pennsylvania												
All races, both sexes	1,664	2	-	-	-	1	1	5	10	35	94	184
Male	999	1	-	-	-	-	1	1	6	24	60	127
Female	665	1	-	-	-	1	-	4	4	11	34	57
White, both sexes	1,493	1	-	-	-	1	-	4	8	31	88	149
Male	902	-	-	-	-	-	-	1	6	22	58	103
Female	591	1	-	-	-	1	-	3	2	9	30	46
Black, both sexes	160	1	-	-	-	-	1	1	2	4	6	34
Male	92	1	-	-	-	-	1	-	-	2	2	24
Female	68	-	-	-	-	-	-	1	2	2	4	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Rhode Island												
All races, both sexes	141	1	-	-	-	-	-	1	-	4	8	10
Male	80	1	-	-	-	-	-	-	-	4	3	5
Female	61	-	-	-	-	-	-	1	-	-	5	5
White, both sexes	131	-	-	-	-	-	-	1	-	4	8	10
Male	74	-	-	-	-	-	-	-	-	4	3	5
Female	57	-	-	-	-	-	-	1	-	-	5	5
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	659	3	3	-	1	-	1	1	5	11	44	79
Male	397	1	1	-	-	-	-	1	2	4	24	50
Female	262	2	2	-	1	-	1	-	3	7	20	29
White, both sexes	508	1	-	-	-	-	-	-	3	8	30	61
Male	311	-	-	-	-	-	-	-	2	3	19	39
Female	197	1	-	-	-	-	-	-	1	5	11	22
Black, both sexes	146	2	3	-	1	-	1	1	2	3	14	17
Male	82	1	1	-	-	-	-	1	-	1	5	10
Female	64	1	2	-	1	-	1	-	2	2	9	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
South Dakota												
All races, both sexes	102	-	-	-	-	-	-	1	3	5	7	16
Male	61	-	-	-	-	-	-	1	3	3	4	12
Female	41	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	64	-	-	-	-	-	-	-	1	1	2	8
Male	38	-	-	-	-	-	-	-	1	1	1	6
Female	26	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	38	-	-	-	-	-	-	1	2	4	5	8
Male	23	-	-	-	-	-	-	1	2	2	3	6
Female	15	-	-	-	-	-	-	-	-	2	2	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	772	2	-	-	-	-	-	4	5	15	48	85
Male	504	1	-	-	-	-	-	4	3	11	35	59
Female	268	1	-	-	-	-	-	-	2	4	13	26
White, both sexes	685	1	-	-	-	-	-	3	4	14	42	71
Male	448	1	-	-	-	-	-	3	2	10	31	49
Female	237	-	-	-	-	-	-	-	2	4	11	22
Black, both sexes	82	1	-	-	-	-	-	1	-	1	6	14
Male	52	-	-	-	-	-	-	1	-	1	4	10
Female	30	1	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Texas												
All races, both sexes	3,153	8	1	4	2	3	7	13	32	79	230	408
Male	2,046	2	-	1	1	1	3	7	21	61	158	296
Female	1,107	6	1	3	1	2	4	6	11	18	72	112
White, both sexes	2,864	7	1	4	-	2	4	13	27	72	213	365
Male	1,869	1	-	1	-	1	1	7	17	58	148	271
Female	995	6	1	3	-	1	3	6	10	14	65	94
Black, both sexes	250	-	-	-	2	1	3	-	4	6	12	38
Male	153	-	-	-	1	-	2	-	3	2	7	21
Female	97	-	-	-	1	1	1	-	1	4	5	17
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	3	1
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	28	1	-	-	-	-	-	-	1	1	2	4
Male	18	1	-	-	-	-	-	-	1	1	1	4
Female	10	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	180	-	-	-	-	-	-	-	-	7	5	21
Male	119	-	-	-	-	-	-	-	-	7	3	11
Female	61	-	-	-	-	-	-	-	-	-	2	10
White, both sexes	173	-	-	-	-	-	-	-	-	6	5	18
Male	113	-	-	-	-	-	-	-	-	6	3	8
Female	60	-	-	-	-	-	-	-	-	-	2	10
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	3
Male	4	-	-	-	-	-	-	-	-	1	-	3
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Vermont												
All races, both sexes	60	-	-	-	-	-	-	1	-	1	5	5
Male	39	-	-	-	-	-	-	-	-	1	4	3
Female	21	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	60	-	-	-	-	-	-	1	-	1	5	5
Male	39	-	-	-	-	-	-	-	-	1	4	3
Female	21	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	803	2	1	-	-	2	-	3	6	21	57	93
Male	504	1	-	-	-	1	-	1	2	12	38	69
Female	299	1	1	-	-	1	-	2	4	9	19	24
White, both sexes	640	1	1	-	-	2	-	2	5	15	46	74
Male	401	-	-	-	-	1	-	1	2	8	34	56
Female	239	1	1	-	-	1	-	1	3	7	12	18
Black, both sexes	155	1	-	-	-	-	-	1	1	5	11	19
Male	102	1	-	-	-	-	-	-	-	4	4	13
Female	53	-	-	-	-	-	-	1	1	1	7	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Washington												
All races, both sexes	668	3	1	-	-	-	2	2	4	18	44	74
Male	412	2	1	-	-	-	1	1	3	14	24	49
Female	256	1	-	-	-	-	1	1	1	4	20	25
White, both sexes	599	3	1	-	-	-	2	2	3	13	39	64
Male	371	2	1	-	-	-	1	1	2	9	22	42
Female	228	1	-	-	-	-	1	1	1	4	17	22
Black, both sexes	17	-	-	-	-	-	-	-	-	1	1	4
Male	10	-	-	-	-	-	-	-	-	1	-	4
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	1	2	4	5
Male	18	-	-	-	-	-	-	-	1	2	2	2
Female	12	-	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	2	-	1
Male	13	-	-	-	-	-	-	-	-	2	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	284	1	-	-	-	-	-	-	-	5	25	26
Male	180	1	-	-	-	-	-	-	-	3	17	19
Female	104	-	-	-	-	-	-	-	-	2	8	7
White, both sexes	272	1	-	-	-	-	-	-	-	5	21	24
Male	177	1	-	-	-	-	-	-	-	3	16	19
Female	95	-	-	-	-	-	-	-	-	2	5	5
Black, both sexes	12	-	-	-	-	-	-	-	-	-	4	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
Wisconsin												
All races, both sexes	636	2	-	-	-	-	3	1	4	15	46	69
Male	371	1	-	-	-	-	2	-	2	8	30	48
Female	265	1	-	-	-	-	1	1	2	7	16	21
White, both sexes	571	2	-	-	-	-	2	1	2	13	38	55
Male	334	1	-	-	-	-	1	-	1	7	24	39
Female	237	1	-	-	-	-	1	1	1	6	14	16
Black, both sexes	36	-	-	-	-	-	-	-	-	-	4	11
Male	19	-	-	-	-	-	-	-	-	-	3	8
Female	17	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	23	-	-	-	-	-	1	-	1	1	4	3
Male	13	-	-	-	-	-	1	-	-	-	3	1
Female	10	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	87	-	-	-	-	-	-	-	2	2	5	13
Male	48	-	-	-	-	-	-	-	2	1	1	5
Female	39	-	-	-	-	-	-	-	-	1	4	8
White, both sexes	78	-	-	-	-	-	-	-	2	-	4	11
Male	44	-	-	-	-	-	-	-	2	-	1	4
Female	34	-	-	-	-	-	-	-	-	-	3	7
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	2	1	2
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
United States												
All races, both sexes	5,347	5,202	4,103	3,471	3,353	2,995	2,283	1,128	423	84	13	6
Male	3,896	3,700	2,790	2,093	1,897	1,543	1,087	508	152	29	3	6
Female	1,451	1,502	1,313	1,378	1,456	1,452	1,196	620	271	55	10	-
White, both sexes	4,452	4,446	3,534	3,054	3,020	2,724	2,076	1,039	385	70	9	5
Male	3,296	3,180	2,418	1,845	1,728	1,409	988	462	147	27	1	5
Female	1,156	1,266	1,116	1,209	1,292	1,315	1,088	577	238	43	8	-
Black, both sexes	696	618	425	310	234	173	142	51	26	11	3	1
Male	469	421	284	190	134	93	62	23	3	1	1	1
Female	227	197	141	120	100	80	80	28	23	10	2	-
American Indian 1/, both sexes	128	87	82	45	44	24	14	5	1	1	-	-
Male	79	61	51	25	16	9	8	4	-	-	-	-
Female	49	26	31	20	28	15	6	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	71	51	62	62	55	74	51	33	11	2	1	-
Male	52	38	37	33	19	32	29	19	2	1	1	-
Female	19	13	25	29	36	42	22	14	9	1	-	-
Alabama												
All races, both sexes	89	99	84	62	59	52	46	25	4	2	-	-
Male	67	67	57	40	27	30	22	7	-	-	-	-
Female	22	32	27	22	32	22	24	18	4	2	-	-
White, both sexes	65	73	71	53	54	45	36	22	2	1	-	-
Male	47	48	48	34	25	26	17	6	-	-	-	-
Female	18	25	23	19	29	19	19	16	2	1	-	-
Black, both sexes	24	26	13	8	5	7	10	3	2	1	-	-
Male	20	19	9	6	2	4	5	1	-	-	-	-
Female	4	7	4	2	3	3	5	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Alaska												
All races, both sexes	13	8	10	4	9	2	4	-	1	-	-	-
Male	8	7	7	2	8	1	2	-	-	-	-	-
Female	5	1	3	2	1	1	2	-	1	-	-	-
White, both sexes	10	7	7	2	8	1	4	-	1	-	-	-
Male	6	6	5	1	7	1	2	-	-	-	-	-
Female	4	1	2	1	1	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	3	2	-	1	-	-	-	-	-	-
Male	2	1	2	1	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	151	123	111	83	95	63	55	18	7	1	-	-
Male	115	83	69	56	59	34	30	7	3	1	-	-
Female	36	40	42	27	36	29	25	11	4	-	-	-
White, both sexes	126	109	96	76	84	60	50	18	7	1	-	-
Male	99	76	61	52	56	32	27	7	3	1	-	-
Female	27	33	35	24	28	28	23	11	4	-	-	-
Black, both sexes	3	3	4	2	1	-	1	-	-	-	-	-
Male	1	1	3	1	1	-	1	-	-	-	-	-
Female	2	2	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	11	11	5	10	2	4	-	-	-	-	-
Male	15	6	5	3	2	1	2	-	-	-	-	-
Female	7	5	6	2	8	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Arkansas												
All races, both sexes	56	52	38	35	32	22	23	7	5	3	-	-
Male	38	36	21	19	19	10	9	6	3	1	-	-
Female	18	16	17	16	13	12	14	1	2	2	-	-
White, both sexes	47	46	36	30	30	22	22	7	5	2	-	-
Male	32	32	21	17	18	10	9	6	3	1	-	-
Female	15	14	15	13	12	12	13	1	2	1	-	-
Black, both sexes	8	6	2	3	2	-	1	-	-	1	-	-
Male	6	4	-	-	1	-	-	-	-	-	-	-
Female	2	2	2	3	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	746	723	523	444	364	284	237	92	30	9	-	2
Male	538	504	376	274	218	171	132	48	14	3	-	2
Female	208	219	147	170	146	113	105	44	16	6	-	-
White, both sexes	646	646	469	390	317	247	208	77	27	6	-	2
Male	473	458	339	244	191	148	116	41	13	2	-	2
Female	173	188	130	146	126	99	92	36	14	4	-	-
Black, both sexes	57	47	27	28	25	8	7	2	1	1	-	-
Male	36	26	19	17	18	5	4	1	1	-	-	-
Female	21	21	8	11	7	3	3	1	-	1	-	-
American Indian 1/, both sexes	12	13	9	5	4	3	-	-	-	-	-	-
Male	8	9	7	2	3	2	-	-	-	-	-	-
Female	4	4	2	3	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	17	18	21	18	26	22	13	2	2	-	-
Male	21	11	11	11	6	16	12	6	-	1	-	-
Female	10	6	7	10	12	10	10	7	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Colorado												
All races, both sexes	96	93	59	54	42	51	26	11	4	2	-	-
Male	67	65	42	34	20	29	12	3	2	2	-	-
Female	29	28	17	20	22	22	14	8	2	-	-	-
White, both sexes	90	90	53	49	42	46	26	11	4	2	-	-
Male	62	63	38	31	20	28	12	3	2	2	-	-
Female	28	27	15	18	22	18	14	8	2	-	-	-
Black, both sexes	5	-	2	1	-	2	-	-	-	-	-	-
Male	4	-	2	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	3	2	-	2	-	-	-	-	-	-
Male	1	2	2	-	-	1	-	-	-	-	-	-
Female	-	1	1	2	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	53	55	44	41	37	44	39	23	17	2	-	-
Male	29	42	27	28	22	24	15	8	4	-	-	-
Female	24	13	17	13	15	20	24	15	13	2	-	-
White, both sexes	44	50	37	39	36	42	38	23	16	2	-	-
Male	25	39	22	26	21	23	15	8	4	-	-	-
Female	19	11	15	13	15	19	23	15	12	2	-	-
Black, both sexes	9	4	7	2	1	2	1	-	1	-	-	-
Male	4	2	5	2	1	1	-	-	-	-	-	-
Female	5	2	2	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Delaware												
All races, both sexes	11	19	9	7	11	9	7	6	-	-	-	-
Male	8	15	7	5	4	5	5	3	-	-	-	-
Female	3	4	2	2	7	4	2	3	-	-	-	-
White, both sexes	9	17	9	6	10	8	7	6	-	-	-	-
Male	6	14	7	5	4	5	5	3	-	-	-	-
Female	3	3	2	1	6	3	2	3	-	-	-	-
Black, both sexes	2	2	-	1	1	1	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	16	19	7	4	7	9	3	4	1	1	-	-
Male	13	14	4	3	6	6	2	2	-	-	-	-
Female	3	5	3	1	1	3	1	2	1	1	-	-
White, both sexes	2	3	-	-	2	2	2	-	-	1	-	-
Male	1	2	-	-	2	1	1	-	-	-	-	-
Female	1	1	-	-	-	1	1	-	-	1	-	-
Black, both sexes	14	16	6	4	5	7	1	2	1	-	-	-
Male	12	12	3	3	4	5	1	1	-	-	-	-
Female	2	4	3	1	1	2	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Florida												
All races, both sexes	359	422	304	253	261	249	168	85	34	8	-	-
Male	264	301	209	158	158	138	90	36	15	2	-	-
Female	95	121	95	95	103	111	78	49	19	6	-	-
White, both sexes	321	378	278	235	252	239	154	81	33	6	-	-
Male	238	270	195	148	153	131	83	35	15	2	-	-
Female	83	108	83	87	99	108	71	46	18	4	-	-
Black, both sexes	34	40	25	16	9	10	12	2	1	2	-	-
Male	22	27	13	8	5	7	6	-	-	-	-	-
Female	12	13	12	8	4	3	6	2	1	2	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	1	-	-	-	-	-
Male	3	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	2	-	-	1	2	-	-	-	-
Male	1	3	1	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Georgia												
All races, both sexes	161	169	127	101	86	74	52	25	10	2	1	-
Male	113	118	85	64	51	37	23	7	2	-	-	-
Female	48	51	42	37	35	37	29	18	8	2	1	-
White, both sexes	115	129	100	75	69	64	37	23	6	1	-	-
Male	86	93	67	53	40	32	17	7	2	-	-	-
Female	29	36	33	22	29	32	20	16	4	1	-	-
Black, both sexes	45	40	25	25	15	9	14	2	3	1	1	-
Male	26	25	17	11	10	4	5	-	-	-	-	-
Female	19	15	8	14	5	5	9	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	2	1	1	-	1	-	-	-
Male	1	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Hawaii												
All races, both sexes	23	17	14	7	15	17	14	8	4	1	-	-
Male	16	12	11	5	6	8	6	5	2	-	-	-
Female	7	5	3	2	9	9	8	3	2	1	-	-
White, both sexes	11	9	8	4	5	5	7	1	2	1	-	-
Male	7	7	6	3	2	4	3	1	1	-	-	-
Female	4	2	2	1	3	1	4	-	1	1	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	7	6	2	9	12	7	7	2	-	-	-
Male	9	5	5	1	4	4	3	4	1	-	-	-
Female	3	2	1	1	5	8	4	3	1	-	-	-
Idaho												
All races, both sexes	21	19	20	15	12	8	5	9	1	-	-	-
Male	13	14	12	11	7	5	3	7	1	-	-	-
Female	8	5	8	4	5	3	2	2	-	-	-	-
White, both sexes	20	18	19	15	11	8	5	9	1	-	-	-
Male	12	14	12	11	6	5	3	7	1	-	-	-
Female	8	4	7	4	5	3	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Illinois												
All races, both sexes	183	176	145	118	111	119	101	53	18	2	1	-
Male	126	128	100	70	59	42	44	23	4	-	-	-
Female	57	48	45	48	52	77	57	30	14	2	1	-
White, both sexes	150	137	119	97	101	105	92	51	17	2	1	-
Male	105	97	84	55	57	38	41	22	4	-	-	-
Female	45	40	35	42	44	67	51	29	13	2	1	-
Black, both sexes	29	36	26	19	7	11	8	2	-	-	-	-
Male	20	28	16	13	1	4	3	1	-	-	-	-
Female	9	8	10	6	6	7	5	1	-	-	-	-
American Indian 1/, both sexes	4	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	3	3	1	-	1	-	-	-
Male	-	3	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	2	3	1	-	1	-	-	-
Indiana												
All races, both sexes	109	94	81	61	62	61	34	27	4	2	-	-
Male	81	66	53	34	32	29	14	10	1	1	-	-
Female	28	28	28	27	30	32	20	17	3	1	-	-
White, both sexes	90	88	74	57	58	56	32	26	3	2	-	-
Male	67	63	50	32	30	27	13	9	1	1	-	-
Female	23	25	24	25	28	29	19	17	2	1	-	-
Black, both sexes	19	6	6	3	4	4	2	1	1	-	-	-
Male	14	3	2	2	2	2	1	1	-	-	-	-
Female	5	3	4	1	2	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Iowa												
All races, both sexes	40	25	29	28	19	29	22	9	4	1	-	-
Male	28	19	21	15	13	14	8	1	4	1	-	-
Female	12	6	8	13	6	15	14	8	-	-	-	-
White, both sexes	35	23	29	28	19	28	22	9	4	1	-	-
Male	23	18	21	15	13	14	8	1	4	1	-	-
Female	12	5	8	13	6	14	14	8	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	40	42	25	37	24	33	15	6	2	-	-	-
Male	29	28	19	26	12	13	7	3	-	-	-	-
Female	11	14	6	11	12	20	8	3	2	-	-	-
White, both sexes	40	39	23	32	23	31	15	6	2	-	-	-
Male	29	27	18	23	12	13	7	3	-	-	-	-
Female	11	12	5	9	11	18	8	3	2	-	-	-
Black, both sexes	-	3	1	2	1	-	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	2	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Kentucky												
All races, both sexes	81	67	68	55	56	37	35	14	8	1	-	-
Male	64	47	41	33	33	19	11	5	5	1	-	-
Female	17	20	27	22	23	18	24	9	3	-	-	-
White, both sexes	73	64	64	52	54	35	34	14	8	1	-	-
Male	56	45	38	32	32	17	11	5	5	1	-	-
Female	17	19	26	20	22	18	23	9	3	-	-	-
Black, both sexes	7	2	4	3	2	2	1	-	-	-	-	-
Male	7	1	3	1	1	2	-	-	-	-	-	-
Female	-	1	1	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	90	75	73	58	60	53	20	13	2	1	1	-
Male	66	52	47	32	33	30	7	3	-	-	-	-
Female	24	23	26	26	27	23	13	10	2	1	1	-
White, both sexes	60	54	60	44	45	43	17	10	1	1	1	-
Male	47	40	37	26	24	23	7	2	-	-	-	-
Female	13	14	23	18	21	20	10	8	1	1	1	-
Black, both sexes	30	20	13	14	14	9	2	3	1	-	-	-
Male	19	11	10	6	9	6	-	1	-	-	-	-
Female	11	9	3	8	5	3	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Maine												
All races, both sexes	26	21	23	13	10	23	11	7	3	1	-	-
Male	24	17	16	6	6	17	5	5	2	-	-	-
Female	2	4	7	7	4	6	6	2	1	1	-	-
White, both sexes	25	21	22	13	10	23	11	7	3	1	-	-
Male	23	17	16	6	6	17	5	5	2	-	-	-
Female	2	4	6	7	4	6	6	2	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	89	92	75	57	55	47	46	13	1	-	2	1
Male	71	65	48	35	31	26	23	6	-	-	2	1
Female	18	27	27	22	24	21	23	7	1	-	-	-
White, both sexes	57	64	47	45	45	38	39	10	1	-	1	-
Male	44	43	28	27	27	21	20	5	-	-	1	-
Female	13	21	19	18	18	17	19	5	1	-	-	-
Black, both sexes	31	27	25	10	10	7	7	2	-	-	1	1
Male	27	22	18	8	4	5	3	-	-	-	1	1
Female	4	5	7	2	6	2	4	2	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	2	-	2	-	1	-	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	1	1	2	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Massachusetts												
All races, both sexes	120	111	72	65	73	83	59	32	7	4	-	-
Male	89	72	45	39	43	44	27	18	3	2	-	-
Female	31	39	27	26	30	39	32	14	4	2	-	-
White, both sexes	116	104	67	63	68	77	57	32	7	4	-	-
Male	86	68	42	39	41	40	25	18	3	2	-	-
Female	30	36	25	24	27	37	32	14	4	2	-	-
Black, both sexes	2	7	5	2	4	1	2	-	-	-	-	-
Male	2	4	3	-	2	1	2	-	-	-	-	-
Female	-	3	2	2	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	5	-	-	-	-	-	-
Male	1	-	-	-	-	3	-	-	-	-	-	-
Female	1	-	-	-	1	2	-	-	-	-	-	-
Michigan												
All races, both sexes	219	194	133	118	126	85	77	47	20	1	-	-
Male	151	155	92	68	72	47	28	22	4	-	-	-
Female	68	39	41	50	54	38	49	25	16	1	-	-
White, both sexes	173	148	111	102	114	76	73	40	16	1	-	-
Male	118	118	77	56	66	44	25	19	3	-	-	-
Female	55	30	34	46	48	32	48	21	13	1	-	-
Black, both sexes	42	43	18	15	11	9	3	6	4	-	-	-
Male	29	34	12	11	6	3	3	3	1	-	-	-
Female	13	9	6	4	5	6	-	3	3	-	-	-
American Indian 1/, both sexes	3	2	4	-	-	-	1	-	-	-	-	-
Male	3	2	3	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	-	-	1	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Minnesota												
All races, both sexes	58	50	44	50	39	27	37	14	11	-	-	-
Male	45	34	30	22	20	14	19	7	3	-	-	-
Female	13	16	14	28	19	13	18	7	8	-	-	-
White, both sexes	52	45	42	47	38	27	37	13	11	-	-	-
Male	40	32	28	22	20	14	19	6	3	-	-	-
Female	12	13	14	25	18	13	18	7	8	-	-	-
Black, both sexes	3	4	-	1	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	3	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	1	1	-	-	-	-	-	-	-
Male	1	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	61	65	42	37	37	29	35	7	-	2	-	-
Male	48	48	27	22	22	14	12	3	-	1	-	-
Female	13	17	15	15	15	15	23	4	-	1	-	-
White, both sexes	38	48	36	28	32	27	33	5	-	-	-	-
Male	30	35	23	17	20	13	11	2	-	-	-	-
Female	8	13	13	11	12	14	22	3	-	-	-	-
Black, both sexes	21	17	6	8	4	2	2	2	-	2	-	-
Male	17	13	4	5	2	1	1	1	-	1	-	-
Female	4	4	2	3	2	1	1	1	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Missouri												
All races, both sexes	81	69	58	49	74	51	51	28	8	1	-	-
Male	61	44	36	34	46	29	23	10	6	-	-	-
Female	20	25	22	15	28	22	28	18	2	1	-	-
White, both sexes	72	60	48	45	69	46	44	26	7	1	-	-
Male	57	38	32	31	43	26	19	9	6	-	-	-
Female	15	22	16	14	26	20	25	17	1	1	-	-
Black, both sexes	8	9	9	4	5	3	6	2	1	-	-	-
Male	4	6	4	3	3	2	3	1	-	-	-	-
Female	4	3	5	1	2	1	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	27	19	10	17	12	12	4	8	1	-	-	-
Male	15	14	7	10	6	8	1	5	-	-	-	-
Female	12	5	3	7	6	4	3	3	1	-	-	-
White, both sexes	19	16	9	14	12	11	3	8	-	-	-	-
Male	13	13	6	8	6	8	1	5	-	-	-	-
Female	6	3	3	6	6	3	2	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	3	1	3	-	1	1	-	1	-	-	-
Male	2	1	1	2	-	-	-	-	-	-	-	-
Female	6	2	-	1	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Nebraska												
All races, both sexes	26	23	20	16	13	16	16	6	5	-	1	-
Male	21	18	12	11	6	12	7	1	1	-	-	-
Female	5	5	8	5	7	4	9	5	4	-	1	-
White, both sexes	22	17	18	15	13	16	16	6	5	-	1	-
Male	18	13	12	10	6	12	7	1	1	-	-	-
Female	4	4	6	5	7	4	9	5	4	-	1	-
Black, both sexes	1	3	2	-	-	-	-	-	-	-	-	-
Male	1	3	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	-	1	-	-	-	-	-	-	-	-
Male	2	2	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	49	59	58	35	33	27	21	9	5	-	-	-
Male	32	44	40	22	21	21	12	4	3	-	-	-
Female	17	15	18	13	12	6	9	5	2	-	-	-
White, both sexes	46	54	53	31	31	25	20	8	4	-	-	-
Male	31	41	38	19	19	21	11	4	3	-	-	-
Female	15	13	15	12	12	4	9	4	1	-	-	-
Black, both sexes	3	3	2	2	-	-	-	-	-	-	-	-
Male	1	1	1	2	-	-	-	-	-	-	-	-
Female	2	2	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	2	1	1	1	1	1	-	-	-
Male	-	2	1	1	1	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
New Hampshire												
All races, both sexes	18	17	17	10	17	16	12	6	3	-	1	-
Male	12	9	9	8	12	10	6	4	2	-	-	-
Female	6	8	8	2	5	6	6	2	1	-	1	-
White, both sexes	18	16	17	10	17	16	12	6	3	-	1	-
Male	12	9	9	8	12	10	6	4	2	-	-	-
Female	6	7	8	2	5	6	6	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	132	127	103	89	90	87	89	37	16	3	1	-
Male	95	90	69	52	50	44	45	18	7	2	-	-
Female	37	37	34	37	40	43	44	19	9	1	1	-
White, both sexes	103	111	84	73	78	77	83	34	12	2	-	-
Male	78	80	58	41	44	39	41	17	6	2	-	-
Female	25	31	26	32	34	38	42	17	6	-	-	-
Black, both sexes	25	15	16	10	9	7	4	2	3	1	1	-
Male	14	9	9	6	4	3	3	-	1	-	-	-
Female	11	6	7	4	5	4	1	2	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	3	6	2	2	2	1	1	-	-	-
Male	3	1	2	5	1	1	1	1	-	-	-	-
Female	1	-	1	1	1	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
New Mexico												
All races, both sexes	57	47	38	36	39	28	19	11	2	1	-	-
Male	44	28	29	22	21	17	9	4	-	-	-	-
Female	13	19	9	14	18	11	10	7	2	1	-	-
White, both sexes	38	37	30	33	30	25	17	9	2	-	-	-
Male	33	22	25	20	19	15	8	2	-	-	-	-
Female	5	15	5	13	11	10	9	7	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	10	8	3	8	3	2	2	-	1	-	-
Male	11	6	4	2	2	2	1	2	-	-	-	-
Female	7	4	4	1	6	1	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	227	216	195	175	179	167	111	68	24	3	1	-
Male	167	154	134	106	94	96	59	36	9	2	-	-
Female	60	62	61	69	85	71	52	32	15	1	1	-
White, both sexes	176	180	149	149	152	146	98	63	21	3	1	-
Male	134	126	98	87	79	86	51	31	9	2	-	-
Female	42	54	51	62	73	60	47	32	12	1	1	-
Black, both sexes	44	33	34	20	22	13	7	4	3	-	-	-
Male	27	26	27	14	14	7	3	4	-	-	-	-
Female	17	7	7	6	8	6	4	-	3	-	-	-
American Indian 1/, both sexes	1	1	2	1	1	-	2	-	-	-	-	-
Male	1	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	2	10	5	4	8	4	1	-	-	-	-
Male	5	1	8	5	1	3	4	1	-	-	-	-
Female	1	1	2	-	3	5	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
North Carolina												
All races, both sexes	161	149	127	122	103	93	61	32	13	-	-	-
Male	118	103	91	84	57	52	23	14	5	-	-	-
Female	43	46	36	38	46	41	38	18	8	-	-	-
White, both sexes	121	115	101	97	84	83	51	30	13	-	-	-
Male	94	81	67	67	44	44	19	13	5	-	-	-
Female	27	34	34	30	40	39	32	17	8	-	-	-
Black, both sexes	32	32	20	20	19	10	8	-	-	-	-	-
Male	19	20	19	15	13	8	3	-	-	-	-	-
Female	13	12	1	5	6	2	5	-	-	-	-	-
American Indian 1/, both sexes	8	1	5	5	-	-	1	1	-	-	-	-
Male	5	1	4	2	-	-	1	1	-	-	-	-
Female	3	-	1	3	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	1	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
North Dakota												
All races, both sexes	8	12	9	6	6	5	2	6	3	2	1	-
Male	6	9	6	4	3	2	-	3	2	-	-	-
Female	2	3	3	2	3	3	2	3	1	2	1	-
White, both sexes	5	12	7	5	5	4	2	6	3	2	1	-
Male	3	9	5	3	3	2	-	3	2	-	-	-
Female	2	3	2	2	2	2	2	3	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	2	1	1	1	-	-	-	-	-	-
Male	3	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Ohio												
All races, both sexes	220	200	160	127	137	126	89	49	19	5	-	-
Male	154	139	111	76	74	63	42	23	7	-	-	-
Female	66	61	49	51	63	63	47	26	12	5	-	-
White, both sexes	192	171	140	117	125	121	83	45	17	4	-	-
Male	132	120	96	69	69	61	40	19	7	-	-	-
Female	60	51	44	48	56	60	43	26	10	4	-	-
Black, both sexes	27	29	19	10	11	5	6	4	1	1	-	-
Male	21	19	14	7	5	2	2	4	-	-	-	-
Female	6	10	5	3	6	3	4	-	1	1	-	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Oklahoma												
All races, both sexes	76	97	75	47	41	34	23	16	6	2	-	-
Male	56	68	46	28	21	15	10	6	3	-	-	-
Female	20	29	29	19	20	19	13	10	3	2	-	-
White, both sexes	64	74	61	40	34	33	20	15	6	2	-	-
Male	49	50	37	24	17	15	7	5	3	-	-	-
Female	15	24	24	16	17	18	13	10	3	2	-	-
Black, both sexes	4	5	4	4	-	-	1	-	-	-	-	-
Male	3	2	2	2	-	-	1	-	-	-	-	-
Female	1	3	2	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	18	10	3	7	1	2	1	-	-	-	-
Male	4	16	7	2	4	-	2	1	-	-	-	-
Female	4	2	3	1	3	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Oregon												
All races, both sexes	75	72	50	42	43	39	23	10	1	2	-	-
Male	55	54	30	22	24	23	9	6	1	1	-	-
Female	20	18	20	20	19	16	14	4	-	1	-	-
White, both sexes	73	67	47	37	41	37	23	9	1	2	-	-
Male	55	51	28	20	24	22	9	5	1	1	-	-
Female	18	16	19	17	17	15	14	4	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	2	1	3	-	-	-	-	-	-	-	-
Male	-	2	1	2	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	2	2	1	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	-	2	1	2	2	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	212	220	139	151	167	182	151	75	29	3	1	2
Male	153	154	84	90	101	79	72	34	9	1	-	2
Female	59	66	55	61	66	103	79	41	20	2	1	-
White, both sexes	179	196	118	138	164	170	144	71	26	2	1	2
Male	130	139	75	82	99	75	68	33	8	1	-	2
Female	49	57	43	56	65	95	76	38	18	1	1	-
Black, both sexes	32	24	19	11	3	11	6	3	1	1	-	-
Male	22	15	9	7	2	4	3	-	-	-	-	-
Female	10	9	10	4	1	7	3	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	2	-	1	1	1	2	-	-	-
Male	1	-	-	1	-	-	1	1	1	-	-	-
Female	-	-	2	1	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Rhode Island												
All races, both sexes	20	17	15	10	15	17	13	8	2	-	-	-
Male	13	12	9	6	7	7	6	7	-	-	-	-
Female	7	5	6	4	8	10	7	1	2	-	-	-
White, both sexes	18	15	13	8	15	16	13	8	2	-	-	-
Male	11	10	9	5	7	7	6	7	-	-	-	-
Female	7	5	4	3	8	9	7	1	2	-	-	-
Black, both sexes	-	2	-	1	-	1	-	-	-	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	103	90	78	64	71	41	34	21	8	1	-	-
Male	76	62	57	36	41	17	10	13	2	-	-	-
Female	27	28	21	28	30	24	24	8	6	1	-	-
White, both sexes	75	65	61	53	60	39	27	17	7	1	-	-
Male	57	44	46	31	32	17	8	11	2	-	-	-
Female	18	21	15	22	28	22	19	6	5	1	-	-
Black, both sexes	27	24	17	11	11	1	6	4	1	-	-	-
Male	18	17	11	5	9	-	1	2	-	-	-	-
Female	9	7	6	6	2	1	5	2	1	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
South Dakota												
All races, both sexes	11	9	13	8	6	14	4	2	1	1	1	-
Male	8	6	9	4	3	5	2	-	-	1	-	-
Female	3	3	4	4	3	9	2	2	1	-	1	-
White, both sexes	8	5	7	7	4	12	4	2	1	1	1	-
Male	6	4	5	3	3	5	2	-	-	1	-	-
Female	2	1	2	4	1	7	2	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	4	6	1	2	2	-	-	-	-	-	-
Male	2	2	4	1	-	-	-	-	-	-	-	-
Female	1	2	2	-	2	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	107	110	99	72	70	63	58	29	5	-	-	-
Male	89	90	76	43	29	28	22	11	3	-	-	-
Female	18	20	23	29	41	35	36	18	2	-	-	-
White, both sexes	92	100	88	64	65	58	50	28	5	-	-	-
Male	77	81	67	38	28	26	21	11	3	-	-	-
Female	15	19	21	26	37	32	29	17	2	-	-	-
Black, both sexes	14	8	11	8	4	5	8	1	-	-	-	-
Male	11	7	9	5	1	2	1	-	-	-	-	-
Female	3	1	2	3	3	3	7	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	1	-	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Texas												
All races, both sexes	486	432	366	296	284	224	158	78	35	6	1	-
Male	374	303	263	165	173	92	82	30	9	3	1	-
Female	112	129	103	131	111	132	76	48	26	3	-	-
White, both sexes	435	388	329	270	265	206	147	75	35	6	-	-
Male	341	274	236	150	164	83	76	28	9	3	-	-
Female	94	114	93	120	101	123	71	47	26	3	-	-
Black, both sexes	47	42	31	22	16	17	6	3	-	-	-	-
Male	30	28	24	14	8	8	3	2	-	-	-	-
Female	17	14	7	8	8	9	3	1	-	-	-	-
American Indian 1/, both sexes	3	1	2	1	-	-	-	-	-	-	-	-
Male	2	-	1	1	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	3	3	1	5	-	-	-	1	-
Male	1	1	2	-	1	1	3	-	-	-	1	-
Female	-	-	2	3	2	-	2	-	-	-	-	-
Utah												
All races, both sexes	19	27	20	26	16	18	13	4	2	1	-	1
Male	19	19	15	14	6	10	10	2	1	1	-	1
Female	-	8	5	12	10	8	3	2	1	-	-	-
White, both sexes	19	26	19	25	16	18	13	4	2	1	-	1
Male	19	18	14	14	6	10	10	2	1	1	-	1
Female	-	8	5	11	10	8	3	2	1	-	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Vermont												
All races, both sexes	7	8	8	7	6	4	5	1	1	1	-	-
Male	4	7	5	4	4	2	3	1	-	1	-	-
Female	3	1	3	3	2	2	2	-	1	-	-	-
White, both sexes	7	8	8	7	6	4	5	1	1	1	-	-
Male	4	7	5	4	4	2	3	1	-	1	-	-
Female	3	1	3	3	2	2	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	113	103	97	90	70	70	51	13	7	4	-	-
Male	79	77	68	53	36	36	23	6	1	1	-	-
Female	34	26	29	37	34	34	28	7	6	3	-	-
White, both sexes	81	77	78	74	59	61	42	12	6	4	-	-
Male	60	56	54	41	30	31	19	6	1	1	-	-
Female	21	21	24	33	29	30	23	6	5	3	-	-
Black, both sexes	31	26	17	14	11	7	9	1	1	-	-	-
Male	19	21	13	12	6	5	4	-	-	-	-	-
Female	12	5	4	2	5	2	5	1	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	2	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Washington												
All races, both sexes	81	108	73	64	71	52	41	19	11	-	-	-
Male	56	80	50	33	36	24	24	10	4	-	-	-
Female	25	28	23	31	35	28	17	9	7	-	-	-
White, both sexes	71	99	64	60	62	48	39	18	11	-	-	-
Male	50	72	47	31	32	23	23	9	4	-	-	-
Female	21	27	17	29	30	25	16	9	7	-	-	-
Black, both sexes	3	3	2	2	-	1	-	-	-	-	-	-
Male	1	3	-	-	-	1	-	-	-	-	-	-
Female	2	-	2	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	5	6	-	4	-	-	-	-	-	-	-
Male	2	4	3	-	2	-	-	-	-	-	-	-
Female	1	1	3	-	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	1	2	5	3	2	1	-	-	-	-
Male	3	1	-	2	2	-	1	1	-	-	-	-
Female	1	-	1	-	3	3	1	-	-	-	-	-
West Virginia												
All races, both sexes	30	49	31	34	31	26	17	6	3	-	-	-
Male	16	41	19	23	17	12	8	3	1	-	-	-
Female	14	8	12	11	14	14	9	3	2	-	-	-
White, both sexes	30	46	29	34	30	26	17	6	3	-	-	-
Male	16	39	19	23	17	12	8	3	1	-	-	-
Female	14	7	10	11	13	14	9	3	2	-	-	-
Black, both sexes	-	3	2	-	1	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	1	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Wisconsin												
All races, both sexes	77	78	72	64	51	66	40	31	15	2	-	-
Male	52	57	40	38	25	30	21	12	4	1	-	-
Female	25	21	32	26	26	36	19	19	11	1	-	-
White, both sexes	61	70	67	59	50	64	39	31	15	2	-	-
Male	45	50	38	37	25	28	21	12	4	1	-	-
Female	16	20	29	22	25	36	18	19	11	1	-	-
Black, both sexes	8	5	4	2	1	-	1	-	-	-	-	-
Male	3	4	1	-	-	-	-	-	-	-	-	-
Female	5	1	3	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	7	1	1	2	-	2	-	-	-	-	-	-
Male	3	1	1	1	-	2	-	-	-	-	-	-
Female	4	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	1	-	-	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	13	14	12	7	6	7	6	-	-	-	-	-
Male	10	9	9	4	3	2	2	-	-	-	-	-
Female	3	5	3	3	3	5	4	-	-	-	-	-
White, both sexes	12	11	12	7	6	7	6	-	-	-	-	-
Male	9	8	9	4	3	2	2	-	-	-	-	-
Female	3	3	3	3	3	5	4	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
United States												
All races, both sexes	12,928	-	-	-	-	-	7	58	159	507	1,299	2,086
Male	9,425	-	-	-	-	-	6	43	104	336	853	1,466
Female	3,503	-	-	-	-	-	1	15	55	171	446	620
White, both sexes	11,205	-	-	-	-	-	4	44	120	430	1,093	1,781
Male	8,250	-	-	-	-	-	3	35	81	293	732	1,260
Female	2,955	-	-	-	-	-	1	9	39	137	361	521
Black, both sexes	1,169	-	-	-	-	-	2	3	12	37	115	215
Male	801	-	-	-	-	-	2	1	5	21	63	151
Female	368	-	-	-	-	-	-	2	7	16	52	64
American Indian 1/, both sexes	410	-	-	-	-	-	1	10	19	32	72	71
Male	258	-	-	-	-	-	1	6	13	15	43	38
Female	152	-	-	-	-	-	-	4	6	17	29	33
Asian or Pacific Islander 2/, both sexes	144	-	-	-	-	-	-	1	8	8	19	19
Male	116	-	-	-	-	-	-	1	5	7	15	17
Female	28	-	-	-	-	-	-	-	3	1	4	2
Alabama												
All races, both sexes	145	-	-	-	-	-	-	-	2	8	14	29
Male	103	-	-	-	-	-	-	-	-	4	12	17
Female	42	-	-	-	-	-	-	-	2	4	2	12
White, both sexes	103	-	-	-	-	-	-	-	2	5	7	19
Male	71	-	-	-	-	-	-	-	-	3	5	12
Female	32	-	-	-	-	-	-	-	2	2	2	7
Black, both sexes	41	-	-	-	-	-	-	-	-	3	6	10
Male	31	-	-	-	-	-	-	-	-	1	6	5
Female	10	-	-	-	-	-	-	-	-	2	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Alaska												
All races, both sexes	34	-	-	-	-	-	-	-	-	2	7	5
Male	17	-	-	-	-	-	-	-	-	1	2	1
Female	17	-	-	-	-	-	-	-	-	1	5	4
White, both sexes	25	-	-	-	-	-	-	-	-	2	5	2
Male	16	-	-	-	-	-	-	-	-	1	2	1
Female	9	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	2	3
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	429	-	-	-	-	-	1	3	8	24	45	60
Male	308	-	-	-	-	-	1	3	6	16	28	35
Female	121	-	-	-	-	-	-	-	2	8	17	25
White, both sexes	348	-	-	-	-	-	1	-	3	18	31	44
Male	256	-	-	-	-	-	1	-	3	13	18	27
Female	92	-	-	-	-	-	-	-	-	5	13	17
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	71	-	-	-	-	-	-	3	4	6	13	14
Male	45	-	-	-	-	-	-	3	2	3	10	6
Female	26	-	-	-	-	-	-	-	2	3	3	8
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Arkansas												
All races, both sexes	67	-	-	-	-	-	2	-	1	1	5	16
Male	43	-	-	-	-	-	2	-	-	1	1	9
Female	24	-	-	-	-	-	-	-	1	-	4	7
White, both sexes	60	-	-	-	-	-	-	-	1	1	5	16
Male	39	-	-	-	-	-	-	-	-	1	1	9
Female	21	-	-	-	-	-	-	-	1	-	4	7
Black, both sexes	7	-	-	-	-	-	2	-	-	-	-	-
Male	4	-	-	-	-	-	2	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,696	-	-	-	-	-	1	14	34	117	244	391
Male	2,008	-	-	-	-	-	1	11	25	86	169	284
Female	688	-	-	-	-	-	-	3	9	31	75	107
White, both sexes	2,406	-	-	-	-	-	1	10	26	107	214	350
Male	1,795	-	-	-	-	-	1	9	19	81	148	254
Female	611	-	-	-	-	-	-	1	7	26	66	96
Black, both sexes	176	-	-	-	-	-	-	2	2	4	13	31
Male	125	-	-	-	-	-	-	1	2	1	9	22
Female	51	-	-	-	-	-	-	1	-	3	4	9
American Indian 1/, both sexes	42	-	-	-	-	-	-	1	2	2	6	3
Male	31	-	-	-	-	-	-	-	2	1	3	3
Female	11	-	-	-	-	-	-	1	-	1	3	-
Asian or Pacific Islander 2/, both sexes	72	-	-	-	-	-	-	1	4	4	11	7
Male	57	-	-	-	-	-	-	1	2	3	9	5
Female	15	-	-	-	-	-	-	-	2	1	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Colorado												
All races, both sexes	254	-	-	-	-	-	-	-	6	10	23	53
Male	179	-	-	-	-	-	-	-	2	4	17	39
Female	75	-	-	-	-	-	-	-	4	6	6	14
White, both sexes	241	-	-	-	-	-	-	-	5	10	23	50
Male	171	-	-	-	-	-	-	-	1	4	17	38
Female	70	-	-	-	-	-	-	-	4	6	6	12
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	94	-	-	-	-	-	-	-	-	2	9	16
Male	65	-	-	-	-	-	-	-	-	-	6	10
Female	29	-	-	-	-	-	-	-	-	2	3	6
White, both sexes	87	-	-	-	-	-	-	-	-	2	8	16
Male	62	-	-	-	-	-	-	-	-	-	5	10
Female	25	-	-	-	-	-	-	-	-	2	3	6
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Delaware												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	4	1
Male	21	-	-	-	-	-	-	-	-	-	2	1
Female	7	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	4	1
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	7	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	43	-	-	-	-	-	-	-	1	-	5	5
Male	38	-	-	-	-	-	-	-	1	-	3	5
Female	5	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	4	5
Male	31	-	-	-	-	-	-	-	-	-	2	5
Female	5	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Florida												
All races, both sexes	925	-	-	-	-	-	-	3	3	25	82	144
Male	671	-	-	-	-	-	-	3	2	13	48	96
Female	254	-	-	-	-	-	-	-	1	12	34	48
White, both sexes	860	-	-	-	-	-	-	3	3	22	78	129
Male	626	-	-	-	-	-	-	3	2	11	45	86
Female	234	-	-	-	-	-	-	-	1	11	33	43
Black, both sexes	61	-	-	-	-	-	-	-	-	3	4	15
Male	41	-	-	-	-	-	-	-	-	2	3	10
Female	20	-	-	-	-	-	-	-	-	1	1	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	227	-	-	-	-	-	-	1	2	9	25	38
Male	170	-	-	-	-	-	-	-	1	5	20	27
Female	57	-	-	-	-	-	-	1	1	4	5	11
White, both sexes	175	-	-	-	-	-	-	1	1	5	20	29
Male	141	-	-	-	-	-	-	-	1	4	16	21
Female	34	-	-	-	-	-	-	1	-	1	4	8
Black, both sexes	50	-	-	-	-	-	-	-	1	4	4	8
Male	28	-	-	-	-	-	-	-	-	1	3	6
Female	22	-	-	-	-	-	-	-	1	3	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Hawaii												
All races, both sexes	28	-	-	-	-	-	-	1	-	-	4	5
Male	21	-	-	-	-	-	-	1	-	-	3	3
Female	7	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	18	-	-	-	-	-	-	1	-	-	1	5
Male	12	-	-	-	-	-	-	1	-	-	-	3
Female	6	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	3	-
Male	8	-	-	-	-	-	-	-	-	-	3	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	103	-	-	-	-	-	-	2	1	5	8	12
Male	69	-	-	-	-	-	-	2	1	1	3	8
Female	34	-	-	-	-	-	-	-	-	4	5	4
White, both sexes	96	-	-	-	-	-	-	2	1	3	6	12
Male	67	-	-	-	-	-	-	2	1	1	3	8
Female	29	-	-	-	-	-	-	-	-	2	3	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	2	2	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Illinois												
All races, both sexes	371	-	-	-	-	-	-	5	11	20	51	73
Male	269	-	-	-	-	-	-	3	9	14	39	52
Female	102	-	-	-	-	-	-	2	2	6	12	21
White, both sexes	317	-	-	-	-	-	-	4	10	19	44	56
Male	234	-	-	-	-	-	-	3	9	13	35	40
Female	83	-	-	-	-	-	-	1	1	6	9	16
Black, both sexes	50	-	-	-	-	-	-	1	-	1	7	16
Male	32	-	-	-	-	-	-	-	-	1	4	11
Female	18	-	-	-	-	-	-	1	-	-	3	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	167	-	-	-	-	-	-	2	6	8	17	33
Male	129	-	-	-	-	-	-	2	5	5	10	21
Female	38	-	-	-	-	-	-	-	1	3	7	12
White, both sexes	159	-	-	-	-	-	-	2	5	8	17	33
Male	123	-	-	-	-	-	-	2	4	5	10	21
Female	36	-	-	-	-	-	-	-	1	3	7	12
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Iowa												
All races, both sexes	114	-	-	-	-	-	-	-	2	5	7	23
Male	81	-	-	-	-	-	-	-	1	4	5	18
Female	33	-	-	-	-	-	-	-	1	1	2	5
White, both sexes	109	-	-	-	-	-	-	-	2	5	7	22
Male	77	-	-	-	-	-	-	-	1	4	5	17
Female	32	-	-	-	-	-	-	-	1	1	2	5
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	91	-	-	-	-	-	-	-	1	2	9	21
Male	69	-	-	-	-	-	-	-	1	2	6	11
Female	22	-	-	-	-	-	-	-	-	-	3	10
White, both sexes	85	-	-	-	-	-	-	-	1	2	7	18
Male	65	-	-	-	-	-	-	-	1	2	4	10
Female	20	-	-	-	-	-	-	-	-	-	3	8
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	3
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Kentucky												
All races, both sexes	127	-	-	-	-	-	-	-	2	6	12	15
Male	95	-	-	-	-	-	-	-	1	5	9	11
Female	32	-	-	-	-	-	-	-	1	1	3	4
White, both sexes	122	-	-	-	-	-	-	-	2	6	11	15
Male	92	-	-	-	-	-	-	-	1	5	9	11
Female	30	-	-	-	-	-	-	-	1	1	2	4
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	126	-	-	-	-	-	-	-	5	3	20	18
Male	93	-	-	-	-	-	-	-	4	2	12	13
Female	33	-	-	-	-	-	-	-	1	1	8	5
White, both sexes	95	-	-	-	-	-	-	-	4	1	11	13
Male	71	-	-	-	-	-	-	-	3	-	8	8
Female	24	-	-	-	-	-	-	-	1	1	3	5
Black, both sexes	30	-	-	-	-	-	-	-	1	2	9	4
Male	21	-	-	-	-	-	-	-	1	2	4	4
Female	9	-	-	-	-	-	-	-	-	-	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Maine												
All races, both sexes	63	-	-	-	-	-	-	-	1	-	2	9
Male	49	-	-	-	-	-	-	-	1	-	1	6
Female	14	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	61	-	-	-	-	-	-	-	1	-	2	9
Male	48	-	-	-	-	-	-	-	1	-	1	6
Female	13	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	164	-	-	-	-	-	-	-	2	10	17	26
Male	108	-	-	-	-	-	-	-	-	6	8	16
Female	56	-	-	-	-	-	-	-	2	4	9	10
White, both sexes	122	-	-	-	-	-	-	-	1	8	15	16
Male	81	-	-	-	-	-	-	-	-	5	8	11
Female	41	-	-	-	-	-	-	-	1	3	7	5
Black, both sexes	41	-	-	-	-	-	-	-	1	2	2	10
Male	26	-	-	-	-	-	-	-	-	1	-	5
Female	15	-	-	-	-	-	-	-	1	1	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Massachusetts												
All races, both sexes	145	-	-	-	-	-	-	-	2	4	16	24
Male	92	-	-	-	-	-	-	-	1	2	8	17
Female	53	-	-	-	-	-	-	-	1	2	8	7
White, both sexes	138	-	-	-	-	-	-	-	2	4	16	22
Male	86	-	-	-	-	-	-	-	1	2	8	15
Female	52	-	-	-	-	-	-	-	1	2	8	7
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	451	-	-	-	-	-	-	1	9	22	50	79
Male	330	-	-	-	-	-	-	1	6	15	36	56
Female	121	-	-	-	-	-	-	-	3	7	14	23
White, both sexes	382	-	-	-	-	-	-	1	7	22	45	67
Male	278	-	-	-	-	-	-	1	5	15	32	48
Female	104	-	-	-	-	-	-	-	2	7	13	19
Black, both sexes	59	-	-	-	-	-	-	-	1	-	4	11
Male	43	-	-	-	-	-	-	-	-	-	3	7
Female	16	-	-	-	-	-	-	-	1	-	1	4
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	1	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Minnesota												
All races, both sexes	171	-	-	-	-	-	-	-	1	7	17	25
Male	121	-	-	-	-	-	-	-	1	3	12	17
Female	50	-	-	-	-	-	-	-	-	4	5	8
White, both sexes	158	-	-	-	-	-	-	-	-	6	14	24
Male	112	-	-	-	-	-	-	-	-	2	10	16
Female	46	-	-	-	-	-	-	-	-	4	4	8
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	2	1
Male	5	-	-	-	-	-	-	-	1	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	88	-	-	-	-	-	-	-	1	5	8	13
Male	68	-	-	-	-	-	-	-	-	2	2	11
Female	20	-	-	-	-	-	-	-	1	3	6	2
White, both sexes	65	-	-	-	-	-	-	-	1	3	5	7
Male	49	-	-	-	-	-	-	-	-	1	1	5
Female	16	-	-	-	-	-	-	-	1	2	4	2
Black, both sexes	23	-	-	-	-	-	-	-	-	2	3	6
Male	19	-	-	-	-	-	-	-	-	1	1	6
Female	4	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Missouri												
All races, both sexes	159	-	-	-	-	-	-	1	4	5	17	32
Male	115	-	-	-	-	-	-	1	2	3	10	25
Female	44	-	-	-	-	-	-	-	2	2	7	7
White, both sexes	141	-	-	-	-	-	-	1	4	5	14	29
Male	106	-	-	-	-	-	-	1	2	3	9	23
Female	35	-	-	-	-	-	-	-	2	2	5	6
Black, both sexes	17	-	-	-	-	-	-	-	-	-	3	3
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	80	-	-	-	-	-	-	2	1	4	10	6
Male	48	-	-	-	-	-	-	-	-	1	7	3
Female	32	-	-	-	-	-	-	2	1	3	3	3
White, both sexes	53	-	-	-	-	-	-	-	-	1	4	4
Male	37	-	-	-	-	-	-	-	-	-	3	2
Female	16	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	26	-	-	-	-	-	-	2	1	3	6	1
Male	11	-	-	-	-	-	-	-	-	1	4	1
Female	15	-	-	-	-	-	-	2	1	2	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Nebraska												
All races, both sexes	75	-	-	-	-	-	-	-	2	-	18	11
Male	49	-	-	-	-	-	-	-	-	-	9	8
Female	26	-	-	-	-	-	-	-	2	-	9	3
White, both sexes	65	-	-	-	-	-	-	-	1	-	17	8
Male	44	-	-	-	-	-	-	-	-	-	9	6
Female	21	-	-	-	-	-	-	-	1	-	8	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	132	-	-	-	-	-	-	-	1	5	19	25
Male	94	-	-	-	-	-	-	-	1	3	10	19
Female	38	-	-	-	-	-	-	-	-	2	9	6
White, both sexes	128	-	-	-	-	-	-	-	1	5	19	23
Male	92	-	-	-	-	-	-	-	1	3	10	18
Female	36	-	-	-	-	-	-	-	-	2	9	5
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
New Hampshire												
All races, both sexes	53	-	-	-	-	-	-	-	-	2	8	8
Male	33	-	-	-	-	-	-	-	-	1	5	4
Female	20	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	52	-	-	-	-	-	-	-	-	2	7	8
Male	32	-	-	-	-	-	-	-	-	1	4	4
Female	20	-	-	-	-	-	-	-	-	1	3	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	283	-	-	-	-	-	1	-	1	15	30	41
Male	201	-	-	-	-	-	1	-	1	14	19	28
Female	82	-	-	-	-	-	-	-	-	1	11	13
White, both sexes	236	-	-	-	-	-	1	-	1	13	25	36
Male	169	-	-	-	-	-	1	-	1	12	17	24
Female	67	-	-	-	-	-	-	-	-	1	8	12
Black, both sexes	39	-	-	-	-	-	-	-	-	2	5	3
Male	25	-	-	-	-	-	-	-	-	2	2	2
Female	14	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
New Mexico												
All races, both sexes	158	-	-	-	-	-	-	4	4	10	20	24
Male	112	-	-	-	-	-	-	3	4	6	10	20
Female	46	-	-	-	-	-	-	1	-	4	10	4
White, both sexes	117	-	-	-	-	-	-	3	2	6	10	14
Male	87	-	-	-	-	-	-	2	2	5	6	12
Female	30	-	-	-	-	-	-	1	-	1	4	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	40	-	-	-	-	-	-	1	2	4	9	10
Male	25	-	-	-	-	-	-	1	2	1	4	8
Female	15	-	-	-	-	-	-	-	-	3	5	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	562	-	-	-	-	-	-	3	3	16	49	79
Male	409	-	-	-	-	-	-	2	2	12	36	55
Female	153	-	-	-	-	-	-	1	1	4	13	24
White, both sexes	451	-	-	-	-	-	-	3	3	11	44	64
Male	331	-	-	-	-	-	-	2	2	8	33	48
Female	120	-	-	-	-	-	-	1	1	3	11	16
Black, both sexes	88	-	-	-	-	-	-	-	-	3	5	10
Male	59	-	-	-	-	-	-	-	-	2	3	4
Female	29	-	-	-	-	-	-	-	-	1	2	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	2
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	17	-	-	-	-	-	-	-	-	1	-	3
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
North Carolina												
All races, both sexes	338	-	-	-	-	-	-	-	4	10	36	42
Male	264	-	-	-	-	-	-	-	3	7	25	29
Female	74	-	-	-	-	-	-	-	1	3	11	13
White, both sexes	245	-	-	-	-	-	-	-	2	7	23	27
Male	188	-	-	-	-	-	-	-	2	5	17	16
Female	57	-	-	-	-	-	-	-	-	2	6	11
Black, both sexes	83	-	-	-	-	-	-	-	1	2	11	15
Male	67	-	-	-	-	-	-	-	1	1	6	13
Female	16	-	-	-	-	-	-	-	-	1	5	2
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	1	2	-
Male	9	-	-	-	-	-	-	-	-	1	2	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	51	-	-	-	-	-	-	-	1	2	7	6
Male	35	-	-	-	-	-	-	-	1	1	3	3
Female	16	-	-	-	-	-	-	-	-	1	4	3
White, both sexes	38	-	-	-	-	-	-	-	-	-	4	5
Male	26	-	-	-	-	-	-	-	-	-	2	2
Female	12	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	2	3	1
Male	9	-	-	-	-	-	-	-	1	1	1	1
Female	4	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Ohio												
All races, both sexes	500	-	-	-	-	-	-	1	2	20	43	92
Male	363	-	-	-	-	-	-	1	1	12	30	67
Female	137	-	-	-	-	-	-	-	1	8	13	25
White, both sexes	444	-	-	-	-	-	-	1	1	20	39	75
Male	323	-	-	-	-	-	-	1	1	12	28	54
Female	121	-	-	-	-	-	-	-	-	8	11	21
Black, both sexes	54	-	-	-	-	-	-	-	1	-	2	17
Male	38	-	-	-	-	-	-	-	-	-	-	13
Female	16	-	-	-	-	-	-	-	1	-	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	240	-	-	-	-	-	-	2	2	5	29	40
Male	176	-	-	-	-	-	-	-	1	4	18	27
Female	64	-	-	-	-	-	-	2	1	1	11	13
White, both sexes	185	-	-	-	-	-	-	1	2	4	19	29
Male	139	-	-	-	-	-	-	-	1	4	12	20
Female	46	-	-	-	-	-	-	1	1	-	7	9
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	4
Male	8	-	-	-	-	-	-	-	-	-	-	4
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	41	-	-	-	-	-	-	1	-	1	9	7
Male	29	-	-	-	-	-	-	-	-	-	6	3
Female	12	-	-	-	-	-	-	1	-	1	3	4
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Oregon												
All races, both sexes	325	-	-	-	-	-	-	1	4	16	19	57
Male	226	-	-	-	-	-	-	1	1	8	10	39
Female	99	-	-	-	-	-	-	-	3	8	9	18
White, both sexes	309	-	-	-	-	-	-	-	4	16	17	51
Male	215	-	-	-	-	-	-	-	1	8	9	35
Female	94	-	-	-	-	-	-	-	3	8	8	16
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	1	-	-	1	5
Male	9	-	-	-	-	-	-	1	-	-	1	3
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Pennsylvania												
All races, both sexes	289	-	-	-	-	-	-	1	6	9	29	58
Male	204	-	-	-	-	-	-	1	3	6	20	38
Female	85	-	-	-	-	-	-	-	3	3	9	20
White, both sexes	268	-	-	-	-	-	-	1	4	9	29	53
Male	190	-	-	-	-	-	-	1	3	6	20	34
Female	78	-	-	-	-	-	-	-	1	3	9	19
Black, both sexes	21	-	-	-	-	-	-	-	2	-	-	5
Male	14	-	-	-	-	-	-	-	-	-	-	4
Female	7	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Rhode Island												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	3	5
Male	19	-	-	-	-	-	-	-	-	-	1	2
Female	20	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	35	-	-	-	-	-	-	-	-	-	3	5
Male	18	-	-	-	-	-	-	-	-	-	1	2
Female	17	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	249	-	-	-	-	-	-	-	2	8	23	42
Male	181	-	-	-	-	-	-	-	1	3	13	32
Female	68	-	-	-	-	-	-	-	1	5	10	10
White, both sexes	184	-	-	-	-	-	-	-	2	6	16	32
Male	140	-	-	-	-	-	-	-	1	3	10	24
Female	44	-	-	-	-	-	-	-	1	3	6	8
Black, both sexes	61	-	-	-	-	-	-	-	-	2	7	9
Male	38	-	-	-	-	-	-	-	-	-	3	7
Female	23	-	-	-	-	-	-	-	-	2	4	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
South Dakota												
All races, both sexes	59	-	-	-	-	-	-	1	3	4	7	11
Male	40	-	-	-	-	-	-	1	3	2	4	8
Female	19	-	-	-	-	-	-	-	-	2	3	3
White, both sexes	31	-	-	-	-	-	-	-	1	-	2	5
Male	22	-	-	-	-	-	-	-	1	-	1	4
Female	9	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	1	2	4	5	6
Male	18	-	-	-	-	-	-	1	2	2	3	4
Female	10	-	-	-	-	-	-	-	-	2	2	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	247	-	-	-	-	-	-	2	2	8	21	38
Male	197	-	-	-	-	-	-	2	1	5	16	27
Female	50	-	-	-	-	-	-	-	1	3	5	11
White, both sexes	220	-	-	-	-	-	-	2	1	7	19	31
Male	178	-	-	-	-	-	-	2	-	4	15	24
Female	42	-	-	-	-	-	-	-	1	3	4	7
Black, both sexes	26	-	-	-	-	-	-	-	-	1	2	7
Male	18	-	-	-	-	-	-	-	-	1	1	3
Female	8	-	-	-	-	-	-	-	-	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Texas												
All races, both sexes	843	-	-	-	-	-	-	6	7	30	100	160
Male	669	-	-	-	-	-	-	4	4	27	76	122
Female	174	-	-	-	-	-	-	2	3	3	24	38
White, both sexes	773	-	-	-	-	-	-	6	7	27	93	155
Male	623	-	-	-	-	-	-	4	4	25	71	120
Female	150	-	-	-	-	-	-	2	3	2	22	35
Black, both sexes	59	-	-	-	-	-	-	-	-	2	4	3
Male	39	-	-	-	-	-	-	-	-	1	3	1
Female	20	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	2	1
Male	3	-	-	-	-	-	-	-	-	1	1	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	64	-	-	-	-	-	-	-	-	4	3	9
Male	49	-	-	-	-	-	-	-	-	4	2	5
Female	15	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	60	-	-	-	-	-	-	-	-	3	3	7
Male	46	-	-	-	-	-	-	-	-	3	2	3
Female	14	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	2
Male	3	-	-	-	-	-	-	-	-	1	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Vermont												
All races, both sexes	35	-	-	-	-	-	-	-	-	1	3	4
Male	25	-	-	-	-	-	-	-	-	1	2	3
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	35	-	-	-	-	-	-	-	-	1	3	4
Male	25	-	-	-	-	-	-	-	-	1	2	3
Female	10	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	195	-	-	-	-	-	-	-	3	9	22	35
Male	157	-	-	-	-	-	-	-	2	6	15	31
Female	38	-	-	-	-	-	-	-	1	3	7	4
White, both sexes	153	-	-	-	-	-	-	-	2	7	17	30
Male	128	-	-	-	-	-	-	-	2	4	14	27
Female	25	-	-	-	-	-	-	-	-	3	3	3
Black, both sexes	41	-	-	-	-	-	-	-	1	2	5	5
Male	29	-	-	-	-	-	-	-	-	2	1	4
Female	12	-	-	-	-	-	-	-	1	-	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Washington												
All races, both sexes	444	-	-	-	-	-	1	2	3	15	37	63
Male	292	-	-	-	-	-	-	1	3	11	21	40
Female	152	-	-	-	-	-	1	1	-	4	16	23
White, both sexes	401	-	-	-	-	-	1	2	2	11	35	55
Male	265	-	-	-	-	-	-	1	2	7	20	35
Female	136	-	-	-	-	-	1	1	-	4	15	20
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	1	2	2	4
Male	14	-	-	-	-	-	-	-	1	2	1	1
Female	11	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	76	-	-	-	-	-	-	-	-	1	11	13
Male	62	-	-	-	-	-	-	-	-	1	7	12
Female	14	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	72	-	-	-	-	-	-	-	-	1	8	13
Male	61	-	-	-	-	-	-	-	-	1	6	12
Female	11	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	3	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Wisconsin												
All races, both sexes	244	-	-	-	-	-	1	-	3	11	31	42
Male	164	-	-	-	-	-	1	-	2	6	21	31
Female	80	-	-	-	-	-	-	-	1	5	10	11
White, both sexes	213	-	-	-	-	-	-	-	1	9	24	36
Male	145	-	-	-	-	-	-	-	1	5	16	27
Female	68	-	-	-	-	-	-	-	-	4	8	9
Black, both sexes	16	-	-	-	-	-	-	-	-	-	4	4
Male	9	-	-	-	-	-	-	-	-	-	3	3
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	12	-	-	-	-	-	1	-	1	1	3	2
Male	8	-	-	-	-	-	1	-	-	-	2	1
Female	4	-	-	-	-	-	-	-	1	1	1	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	1	-	-
Male	2	-	-	-	-	-	-	-	1	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	37	-	-	-	-	-	-	-	-	2	3	9
Male	24	-	-	-	-	-	-	-	-	1	1	4
Female	13	-	-	-	-	-	-	-	-	1	2	5
White, both sexes	30	-	-	-	-	-	-	-	-	-	2	7
Male	21	-	-	-	-	-	-	-	-	-	1	3
Female	9	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	2	1	2
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
United States												
All races, both sexes	2,404	2,156	1,565	1,047	797	478	269	76	16	2	-	2
Male	1,808	1,646	1,204	780	572	350	187	57	10	1	-	2
Female	596	510	361	267	225	128	82	19	6	1	-	-
White, both sexes	2,078	1,894	1,349	927	714	434	248	70	16	1	-	2
Male	1,577	1,460	1,047	694	513	316	174	52	10	1	-	2
Female	501	434	302	233	201	118	74	18	6	-	-	-
Black, both sexes	232	198	149	89	65	32	14	5	-	1	-	-
Male	158	137	113	65	48	27	6	4	-	-	-	-
Female	74	61	36	24	17	5	8	1	-	1	-	-
American Indian 1/, both sexes	66	52	46	15	15	6	4	1	-	-	-	-
Male	49	39	29	9	9	2	4	1	-	-	-	-
Female	17	13	17	6	6	4	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	12	21	16	3	6	3	-	-	-	-	-
Male	24	10	15	12	2	5	3	-	-	-	-	-
Female	4	2	6	4	1	1	-	-	-	-	-	-
Alabama												
All races, both sexes	35	22	20	5	4	2	3	1	-	-	-	-
Male	27	17	17	4	2	1	2	-	-	-	-	-
Female	8	5	3	1	2	1	1	1	-	-	-	-
White, both sexes	24	18	15	5	3	2	3	-	-	-	-	-
Male	17	14	12	4	1	1	2	-	-	-	-	-
Female	7	4	3	1	2	1	1	-	-	-	-	-
Black, both sexes	11	4	5	-	1	-	-	1	-	-	-	-
Male	10	3	5	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Alaska												
All races, both sexes	7	5	2	1	3	1	1	-	-	-	-	-
Male	3	5	1	1	2	-	1	-	-	-	-	-
Female	4	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	6	5	-	1	3	-	1	-	-	-	-	-
Male	3	5	-	1	2	-	1	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	80	69	48	33	34	14	7	2	1	-	-	-
Male	65	50	36	24	25	10	6	2	1	-	-	-
Female	15	19	12	9	9	4	1	-	-	-	-	-
White, both sexes	64	61	43	31	30	13	6	2	1	-	-	-
Male	53	44	34	23	23	9	5	2	1	-	-	-
Female	11	17	9	8	7	4	1	-	-	-	-	-
Black, both sexes	1	2	2	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	6	3	2	4	-	1	-	-	-	-	-
Male	11	5	1	1	2	-	1	-	-	-	-	-
Female	4	1	2	1	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Arkansas												
All races, both sexes	11	18	6	2	2	1	2	-	-	-	-	-
Male	6	14	4	1	2	1	2	-	-	-	-	-
Female	5	4	2	1	-	-	-	-	-	-	-	-
White, both sexes	9	17	5	1	2	1	2	-	-	-	-	-
Male	5	13	4	1	2	1	2	-	-	-	-	-
Female	4	4	1	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	502	450	325	247	185	100	61	18	4	1	-	2
Male	379	339	254	180	138	75	49	13	2	1	-	2
Female	123	111	71	67	47	25	12	5	2	-	-	-
White, both sexes	448	404	290	222	162	90	58	17	4	1	-	2
Male	342	306	225	161	120	66	46	12	2	1	-	2
Female	106	98	65	61	42	24	12	5	2	-	-	-
Black, both sexes	32	30	21	16	17	5	2	1	-	-	-	-
Male	20	19	17	13	14	4	2	1	-	-	-	-
Female	12	11	4	3	3	1	-	-	-	-	-	-
American Indian 1/, both sexes	6	8	6	2	4	2	-	-	-	-	-	-
Male	4	7	6	-	3	2	-	-	-	-	-	-
Female	2	1	-	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	8	8	7	2	3	1	-	-	-	-	-
Male	13	7	6	6	1	3	1	-	-	-	-	-
Female	3	1	2	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Colorado												
All races, both sexes	62	33	21	21	12	10	2	1	-	-	-	-
Male	41	22	18	19	8	8	1	-	-	-	-	-
Female	21	11	3	2	4	2	1	1	-	-	-	-
White, both sexes	60	31	18	20	12	9	2	1	-	-	-	-
Male	40	21	15	18	8	8	1	-	-	-	-	-
Female	20	10	3	2	4	1	1	1	-	-	-	-
Black, both sexes	1	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	1	-	-	1	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	21	15	10	2	1	3	1	-	-	-	-
Male	8	18	10	9	2	1	1	-	-	-	-	-
Female	6	3	5	1	-	-	2	1	-	-	-	-
White, both sexes	13	20	11	10	2	1	3	1	-	-	-	-
Male	8	18	8	9	2	1	1	-	-	-	-	-
Female	5	2	3	1	-	-	2	1	-	-	-	-
Black, both sexes	1	1	4	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	1	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Delaware												
All races, both sexes	6	8	2	5	2	-	-	-	-	-	-	-
Male	5	6	2	4	1	-	-	-	-	-	-	-
Female	1	2	-	1	1	-	-	-	-	-	-	-
White, both sexes	6	7	2	5	2	-	-	-	-	-	-	-
Male	5	5	2	4	1	-	-	-	-	-	-	-
Female	1	2	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	8	4	2	4	4	1	1	-	-	-	-
Male	7	7	3	2	4	4	1	1	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	2	-	-	1	1	-	-	-	-	-	-
Male	1	2	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	6	4	2	3	3	1	1	-	-	-	-
Male	6	5	3	2	3	3	1	1	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Florida												
All races, both sexes	139	174	131	82	67	45	22	5	3	-	-	-
Male	107	134	97	61	54	33	17	4	2	-	-	-
Female	32	40	34	21	13	12	5	1	1	-	-	-
White, both sexes	129	158	121	79	66	44	20	5	3	-	-	-
Male	102	122	90	58	53	32	16	4	2	-	-	-
Female	27	36	31	21	13	12	4	1	1	-	-	-
Black, both sexes	9	14	10	3	1	1	1	-	-	-	-	-
Male	4	10	7	3	1	1	-	-	-	-	-	-
Female	5	4	3	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	43	42	33	11	12	4	6	-	-	1	-	-
Male	35	30	27	9	9	3	4	-	-	-	-	-
Female	8	12	6	2	3	1	2	-	-	1	-	-
White, both sexes	34	34	25	7	10	4	5	-	-	-	-	-
Male	29	27	21	7	8	3	4	-	-	-	-	-
Female	5	7	4	-	2	1	1	-	-	-	-	-
Black, both sexes	9	8	8	4	2	-	1	-	-	1	-	-
Male	6	3	6	2	1	-	-	-	-	-	-	-
Female	3	5	2	2	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Hawaii												
All races, both sexes	6	3	3	3	2	-	1	-	-	-	-	-
Male	6	2	2	2	1	-	1	-	-	-	-	-
Female	-	1	1	1	1	-	-	-	-	-	-	-
White, both sexes	2	3	2	2	1	-	1	-	-	-	-	-
Male	2	2	2	1	-	-	1	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	1	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	18	17	16	10	8	4	1	1	-	-	-	-
Male	12	12	12	9	4	3	1	1	-	-	-	-
Female	6	5	4	1	4	1	-	-	-	-	-	-
White, both sexes	17	16	16	10	7	4	1	1	-	-	-	-
Male	11	12	12	9	3	3	1	1	-	-	-	-
Female	6	4	4	1	4	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Illinois												
All races, both sexes	74	50	34	21	12	12	7	1	-	-	-	-
Male	53	39	26	15	8	7	3	1	-	-	-	-
Female	21	11	8	6	4	5	4	-	-	-	-	-
White, both sexes	65	42	29	16	12	12	7	1	-	-	-	-
Male	47	34	23	11	8	7	3	1	-	-	-	-
Female	18	8	6	5	4	5	4	-	-	-	-	-
Black, both sexes	8	8	5	4	-	-	-	-	-	-	-	-
Male	5	5	3	3	-	-	-	-	-	-	-	-
Female	3	3	2	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	34	26	20	9	4	5	2	1	-	-	-	-
Male	30	20	18	6	4	5	2	1	-	-	-	-
Female	4	6	2	3	-	-	-	-	-	-	-	-
White, both sexes	29	25	19	9	4	5	2	1	-	-	-	-
Male	26	20	17	6	4	5	2	1	-	-	-	-
Female	3	5	2	3	-	-	-	-	-	-	-	-
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Iowa												
All races, both sexes	27	10	16	7	6	8	2	1	-	-	-	-
Male	17	8	12	6	5	5	-	-	-	-	-	-
Female	10	2	4	1	1	3	2	1	-	-	-	-
White, both sexes	25	8	16	7	6	8	2	1	-	-	-	-
Male	15	7	12	6	5	5	-	-	-	-	-	-
Female	10	1	4	1	1	3	2	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	14	16	11	8	5	2	2	-	-	-	-	-
Male	12	11	10	8	5	1	2	-	-	-	-	-
Female	2	5	1	-	-	1	-	-	-	-	-	-
White, both sexes	14	16	11	7	5	2	2	-	-	-	-	-
Male	12	11	10	7	5	1	2	-	-	-	-	-
Female	2	5	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Kentucky												
All races, both sexes	32	17	14	12	9	6	2	-	-	-	-	-
Male	27	13	8	9	6	4	2	-	-	-	-	-
Female	5	4	6	3	3	2	-	-	-	-	-	-
White, both sexes	32	17	13	12	8	4	2	-	-	-	-	-
Male	27	13	7	9	6	2	2	-	-	-	-	-
Female	5	4	6	3	2	2	-	-	-	-	-	-
Black, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	21	18	18	10	7	4	1	1	-	-	-	-
Male	15	14	15	7	6	4	-	1	-	-	-	-
Female	6	4	3	3	1	-	1	-	-	-	-	-
White, both sexes	16	15	17	8	6	2	1	1	-	-	-	-
Male	12	12	14	6	5	2	-	1	-	-	-	-
Female	4	3	3	2	1	-	1	-	-	-	-	-
Black, both sexes	5	3	1	2	1	2	-	-	-	-	-	-
Male	3	2	1	1	1	2	-	-	-	-	-	-
Female	2	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Maine												
All races, both sexes	16	12	11	2	4	4	2	-	-	-	-	-
Male	14	9	7	1	4	4	2	-	-	-	-	-
Female	2	3	4	1	-	-	-	-	-	-	-	-
White, both sexes	15	12	10	2	4	4	2	-	-	-	-	-
Male	13	9	7	1	4	4	2	-	-	-	-	-
Female	2	3	3	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	27	29	22	10	13	4	3	1	-	-	-	-
Male	21	23	15	8	9	2	-	-	-	-	-	-
Female	6	6	7	2	4	2	3	1	-	-	-	-
White, both sexes	20	23	15	8	11	2	2	1	-	-	-	-
Male	14	18	10	7	7	1	-	-	-	-	-	-
Female	6	5	5	1	4	1	2	1	-	-	-	-
Black, both sexes	7	6	6	2	2	2	1	-	-	-	-	-
Male	7	5	4	1	2	1	-	-	-	-	-	-
Female	-	1	2	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Massachusetts												
All races, both sexes	32	18	17	12	9	4	5	2	-	-	-	-
Male	20	12	10	7	6	3	4	2	-	-	-	-
Female	12	6	7	5	3	1	1	-	-	-	-	-
White, both sexes	30	16	16	12	9	4	5	2	-	-	-	-
Male	18	11	9	7	6	3	4	2	-	-	-	-
Female	12	5	7	5	3	1	1	-	-	-	-	-
Black, both sexes	1	2	1	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	97	75	35	35	19	15	12	1	1	-	-	-
Male	64	64	28	25	16	13	5	1	-	-	-	-
Female	33	11	7	10	3	2	7	-	1	-	-	-
White, both sexes	82	58	27	28	17	14	12	1	1	-	-	-
Male	55	48	22	20	14	12	5	1	-	-	-	-
Female	27	10	5	8	3	2	7	-	1	-	-	-
Black, both sexes	14	15	5	6	2	1	-	-	-	-	-	-
Male	8	14	4	4	2	1	-	-	-	-	-	-
Female	6	1	1	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	3	-	-	-	-	-	-	-	-	-
Male	1	2	2	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Minnesota												
All races, both sexes	26	32	25	17	10	6	5	-	-	-	-	-
Male	21	23	20	11	8	3	2	-	-	-	-	-
Female	5	9	5	6	2	3	3	-	-	-	-	-
White, both sexes	23	30	24	16	10	6	5	-	-	-	-	-
Male	18	23	19	11	8	3	2	-	-	-	-	-
Female	5	7	5	5	2	3	3	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	18	12	6	4	3	1	-	-	-	-	-
Male	14	15	11	6	3	3	1	-	-	-	-	-
Female	3	3	1	-	1	-	-	-	-	-	-	-
White, both sexes	13	15	10	4	3	3	1	-	-	-	-	-
Male	10	12	9	4	3	3	1	-	-	-	-	-
Female	3	3	1	-	-	-	-	-	-	-	-	-
Black, both sexes	4	3	2	2	1	-	-	-	-	-	-	-
Male	4	3	2	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Missouri												
All races, both sexes	28	28	14	13	11	4	1	1	-	-	-	-
Male	21	19	9	12	8	4	1	-	-	-	-	-
Female	7	9	5	1	3	-	-	1	-	-	-	-
White, both sexes	25	26	10	11	10	4	1	1	-	-	-	-
Male	20	18	7	10	8	4	1	-	-	-	-	-
Female	5	8	3	1	2	-	-	1	-	-	-	-
Black, both sexes	2	2	4	2	1	-	-	-	-	-	-	-
Male	1	1	2	2	-	-	-	-	-	-	-	-
Female	1	1	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	20	13	8	6	3	5	1	1	-	-	-	-
Male	10	11	6	4	1	4	-	1	-	-	-	-
Female	10	2	2	2	2	1	1	-	-	-	-	-
White, both sexes	13	10	7	5	3	4	1	1	-	-	-	-
Male	8	10	5	3	1	4	-	1	-	-	-	-
Female	5	-	2	2	2	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	3	1	1	-	1	-	-	-	-	-	-
Male	2	1	1	1	-	-	-	-	-	-	-	-
Female	5	2	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Nebraska												
All races, both sexes	11	5	7	8	3	2	8	-	-	-	-	-
Male	7	4	5	7	2	2	5	-	-	-	-	-
Female	4	1	2	1	1	-	3	-	-	-	-	-
White, both sexes	8	5	6	7	3	2	8	-	-	-	-	-
Male	5	4	5	6	2	2	5	-	-	-	-	-
Female	3	1	1	1	1	-	3	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	20	18	9	8	7	1	-	1	-	-	-
Male	13	14	14	6	5	7	1	-	1	-	-	-
Female	5	6	4	3	3	-	-	-	-	-	-	-
White, both sexes	18	20	17	9	7	7	1	-	1	-	-	-
Male	13	14	14	6	4	7	1	-	1	-	-	-
Female	5	6	3	3	3	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
New Hampshire												
All races, both sexes	9	6	7	3	4	4	2	-	-	-	-	-
Male	8	3	3	3	2	3	1	-	-	-	-	-
Female	1	3	4	-	2	1	1	-	-	-	-	-
White, both sexes	9	6	7	3	4	4	2	-	-	-	-	-
Male	8	3	3	3	2	3	1	-	-	-	-	-
Female	1	3	4	-	2	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	54	48	36	17	21	10	8	1	-	-	-	-
Male	32	38	25	14	14	8	6	1	-	-	-	-
Female	22	10	11	3	7	2	2	-	-	-	-	-
White, both sexes	43	41	31	14	17	7	6	1	-	-	-	-
Male	26	32	22	12	11	5	5	1	-	-	-	-
Female	17	9	9	2	6	2	1	-	-	-	-	-
Black, both sexes	10	7	4	-	4	2	2	-	-	-	-	-
Male	5	6	2	-	3	2	1	-	-	-	-	-
Female	5	1	2	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	-	1	-	-	-	-	-	-
Male	1	-	1	2	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
New Mexico												
All races, both sexes	28	23	18	12	8	3	1	3	-	-	-	-
Male	23	14	13	9	6	2	1	1	-	-	-	-
Female	5	9	5	3	2	1	-	2	-	-	-	-
White, both sexes	21	21	14	12	8	3	1	2	-	-	-	-
Male	18	14	10	9	6	2	1	-	-	-	-	-
Female	3	7	4	3	2	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	2	4	-	-	-	-	1	-	-	-	-
Male	5	-	3	-	-	-	-	1	-	-	-	-
Female	2	2	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	99	97	76	47	48	30	10	5	-	-	-	-
Male	70	71	61	36	34	19	6	5	-	-	-	-
Female	29	26	15	11	14	11	4	-	-	-	-	-
White, both sexes	78	77	54	39	40	26	9	3	-	-	-	-
Male	58	57	42	28	28	17	5	3	-	-	-	-
Female	20	20	12	11	12	9	4	-	-	-	-	-
Black, both sexes	18	17	15	6	8	4	-	2	-	-	-	-
Male	9	12	13	6	6	2	-	2	-	-	-	-
Female	9	5	2	-	2	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	6	2	-	-	1	-	-	-	-	-
Male	3	1	6	2	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
North Carolina												
All races, both sexes	77	56	39	35	21	9	4	5	-	-	-	-
Male	64	44	33	30	16	8	2	3	-	-	-	-
Female	13	12	6	5	5	1	2	2	-	-	-	-
White, both sexes	58	42	29	28	12	8	4	5	-	-	-	-
Male	48	35	23	23	7	7	2	3	-	-	-	-
Female	10	7	6	5	5	1	2	2	-	-	-	-
Black, both sexes	15	13	9	7	9	1	-	-	-	-	-	-
Male	12	8	9	7	9	1	-	-	-	-	-	-
Female	3	5	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	1	1	-	-	-	-	-	-	-	-	-
Male	4	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	9	8	5	2	2	-	1	2	-	-	-
Male	5	7	5	4	2	1	-	1	2	-	-	-
Female	1	2	3	1	-	1	-	-	-	-	-	-
White, both sexes	3	9	6	4	2	2	-	1	2	-	-	-
Male	2	7	4	3	2	1	-	1	2	-	-	-
Female	1	2	2	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	2	1	-	-	-	-	-	-	-	-
Male	3	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Ohio												
All races, both sexes	101	90	51	36	29	18	14	3	-	-	-	-
Male	74	70	40	26	17	15	7	3	-	-	-	-
Female	27	20	11	10	12	3	7	-	-	-	-	-
White, both sexes	92	82	43	32	27	17	12	3	-	-	-	-
Male	68	62	33	23	17	15	6	3	-	-	-	-
Female	24	20	10	9	10	2	6	-	-	-	-	-
Black, both sexes	9	8	8	4	2	1	2	-	-	-	-	-
Male	6	8	7	3	-	-	1	-	-	-	-	-
Female	3	-	1	1	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	40	57	31	13	6	7	7	1	-	-	-	-
Male	32	43	23	10	5	7	6	-	-	-	-	-
Female	8	14	8	3	1	-	1	1	-	-	-	-
White, both sexes	38	41	24	10	5	7	4	1	-	-	-	-
Male	30	31	19	8	4	7	3	-	-	-	-	-
Female	8	10	5	2	1	-	1	1	-	-	-	-
Black, both sexes	1	5	1	1	-	-	1	-	-	-	-	-
Male	1	2	-	-	-	-	1	-	-	-	-	-
Female	-	3	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	11	6	2	1	-	2	-	-	-	-	-
Male	1	10	4	2	1	-	2	-	-	-	-	-
Female	-	1	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Oregon												
All races, both sexes	63	55	34	29	25	17	4	1	-	-	-	-
Male	48	43	22	17	19	14	3	1	-	-	-	-
Female	15	12	12	12	6	3	1	-	-	-	-	-
White, both sexes	63	53	33	26	25	16	4	1	-	-	-	-
Male	48	41	21	16	19	13	3	1	-	-	-	-
Female	15	12	12	10	6	3	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	2	-	-	-	-	-	-	-	-
Male	-	2	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	49	47	31	25	14	8	8	4	-	-	-	-
Male	37	34	21	19	9	6	6	4	-	-	-	-
Female	12	13	10	6	5	2	2	-	-	-	-	-
White, both sexes	45	42	28	25	13	7	8	4	-	-	-	-
Male	33	31	20	19	8	5	6	4	-	-	-	-
Female	12	11	8	6	5	2	2	-	-	-	-	-
Black, both sexes	4	5	3	-	1	1	-	-	-	-	-	-
Male	4	3	1	-	1	1	-	-	-	-	-	-
Female	-	2	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Rhode Island												
All races, both sexes	10	4	7	3	2	4	1	-	-	-	-	-
Male	6	2	4	2	1	-	1	-	-	-	-	-
Female	4	2	3	1	1	4	-	-	-	-	-	-
White, both sexes	10	4	5	1	2	4	1	-	-	-	-	-
Male	6	2	4	1	1	-	1	-	-	-	-	-
Female	4	2	1	-	1	4	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	54	34	32	19	14	13	4	2	2	-	-	-
Male	43	24	28	15	8	8	3	2	1	-	-	-
Female	11	10	4	4	6	5	1	-	1	-	-	-
White, both sexes	37	23	24	14	12	12	2	2	2	-	-	-
Male	32	17	22	11	7	8	2	2	1	-	-	-
Female	5	6	2	3	5	4	-	-	1	-	-	-
Black, both sexes	17	10	8	5	2	-	1	-	-	-	-	-
Male	11	6	6	4	1	-	-	-	-	-	-	-
Female	6	4	2	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
South Dakota												
All races, both sexes	8	5	8	3	4	5	-	-	-	-	-	-
Male	6	3	5	3	2	3	-	-	-	-	-	-
Female	2	2	3	-	2	2	-	-	-	-	-	-
White, both sexes	6	2	5	3	3	4	-	-	-	-	-	-
Male	4	1	3	3	2	3	-	-	-	-	-	-
Female	2	1	2	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	3	-	1	1	-	-	-	-	-	-
Male	2	2	2	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	43	43	41	22	14	7	4	2	-	-	-	-
Male	39	40	35	15	9	5	2	1	-	-	-	-
Female	4	3	6	7	5	2	2	1	-	-	-	-
White, both sexes	38	42	36	19	14	6	3	2	-	-	-	-
Male	35	39	31	12	9	4	2	1	-	-	-	-
Female	3	3	5	7	5	2	1	1	-	-	-	-
Black, both sexes	5	1	5	3	-	1	1	-	-	-	-	-
Male	4	1	4	3	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Texas												
All races, both sexes	168	132	107	61	44	18	8	1	1	-	-	-
Male	133	111	89	48	33	12	8	1	1	-	-	-
Female	35	21	18	13	11	6	-	-	-	-	-	-
White, both sexes	149	121	97	54	39	15	8	1	1	-	-	-
Male	118	103	82	46	31	9	8	1	1	-	-	-
Female	31	18	15	8	8	6	-	-	-	-	-	-
Black, both sexes	17	11	7	7	5	3	-	-	-	-	-	-
Male	13	8	6	2	2	3	-	-	-	-	-	-
Female	4	3	1	5	3	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	15	11	8	10	3	-	1	-	-	-	-	-
Male	15	9	7	4	2	-	1	-	-	-	-	-
Female	-	2	1	6	1	-	-	-	-	-	-	-
White, both sexes	15	11	8	9	3	-	1	-	-	-	-	-
Male	15	9	7	4	2	-	1	-	-	-	-	-
Female	-	2	1	5	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Vermont												
All races, both sexes	6	7	5	3	3	1	1	1	-	-	-	-
Male	4	6	4	2	1	1	-	1	-	-	-	-
Female	2	1	1	1	2	-	1	-	-	-	-	-
White, both sexes	6	7	5	3	3	1	1	1	-	-	-	-
Male	4	6	4	2	1	1	-	1	-	-	-	-
Female	2	1	1	1	2	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	29	33	27	13	15	5	3	1	-	-	-	-
Male	23	28	22	12	10	5	2	1	-	-	-	-
Female	6	5	5	1	5	-	1	-	-	-	-	-
White, both sexes	19	27	21	8	14	4	3	1	-	-	-	-
Male	17	23	18	7	9	4	2	1	-	-	-	-
Female	2	4	3	1	5	-	1	-	-	-	-	-
Black, both sexes	10	6	5	5	1	1	-	-	-	-	-	-
Male	6	5	4	5	1	1	-	-	-	-	-	-
Female	4	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Washington												
All races, both sexes	62	83	55	42	34	29	14	3	1	-	-	-
Male	44	59	40	23	22	16	9	3	-	-	-	-
Female	18	24	15	19	12	13	5	-	1	-	-	-
White, both sexes	55	75	48	40	31	28	14	3	1	-	-	-
Male	40	52	37	23	21	15	9	3	-	-	-	-
Female	15	23	11	17	10	13	5	-	1	-	-	-
Black, both sexes	2	2	1	2	-	1	-	-	-	-	-	-
Male	1	2	-	-	-	1	-	-	-	-	-	-
Female	1	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	5	6	-	3	-	-	-	-	-	-	-
Male	1	4	3	-	1	-	-	-	-	-	-	-
Female	1	1	3	-	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	12	17	7	10	5	-	-	-	-	-	-	-
Male	7	13	7	10	5	-	-	-	-	-	-	-
Female	5	4	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	16	7	10	5	-	-	-	-	-	-	-
Male	7	13	7	10	5	-	-	-	-	-	-	-
Female	5	3	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Wisconsin												
All races, both sexes	38	37	24	20	15	11	9	2	-	-	-	-
Male	28	26	16	13	5	7	7	1	-	-	-	-
Female	10	11	8	7	10	4	2	1	-	-	-	-
White, both sexes	33	34	24	17	14	11	8	2	-	-	-	-
Male	23	24	16	13	5	7	7	1	-	-	-	-
Female	10	10	8	4	9	4	1	1	-	-	-	-
Black, both sexes	2	2	-	2	1	-	1	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	5	5	2	2	-	1	-	-	-	-	-
Male	7	3	4	2	2	-	-	-	-	-	-	-
Female	1	2	1	-	-	-	1	-	-	-	-	-
White, both sexes	7	4	5	2	2	-	1	-	-	-	-	-
Male	6	3	4	2	2	-	-	-	-	-	-	-
Female	1	1	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
United States												
All races, both sexes	99	-	-	-	1	-	-	-	1	1	9	12
Male	49	-	-	-	-	-	-	-	-	-	4	10
Female	50	-	-	-	1	-	-	-	1	1	5	2
White, both sexes	79	-	-	-	1	-	-	-	-	1	7	10
Male	42	-	-	-	-	-	-	-	-	-	4	9
Female	37	-	-	-	1	-	-	-	-	1	3	1
Black, both sexes	12	-	-	-	-	-	-	-	1	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Colorado												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Florida												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Iowa												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Kentucky												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Missouri												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	-	-	1	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	1	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
North Carolina												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Ohio												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Texas												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Washington												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
United States												
All races, both sexes	12	6	11	10	13	11	8	4	-	-	-	-
Male	9	5	5	3	5	4	2	2	-	-	-	-
Female	3	1	6	7	8	7	6	2	-	-	-	-
White, both sexes	8	5	9	9	13	8	4	4	-	-	-	-
Male	6	4	5	3	5	3	1	2	-	-	-	-
Female	2	1	4	6	8	5	3	2	-	-	-	-
Black, both sexes	2	1	2	-	-	1	2	-	-	-	-	-
Male	2	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	2	2	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	1	2	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	2	-	2	1	1	2	-	-	-	-	-
Male	2	1	-	1	-	-	-	-	-	-	-	-
Female	3	1	-	1	1	1	2	-	-	-	-	-
White, both sexes	4	2	-	1	1	1	1	-	-	-	-	-
Male	2	1	-	1	-	-	-	-	-	-	-	-
Female	2	1	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Colorado												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Florida												
All races, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Hawaii												
All races, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Illinois												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Iowa												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Kentucky												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Massachusetts												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Missouri												
All races, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
North Carolina												
All races, both sexes	-	-	1	1	1	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
White, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Ohio												
All races, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Oregon												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
South Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Texas												
All races, both sexes	2	-	1	-	4	2	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	3	2	-	-	-	-	-	-
White, both sexes	2	-	1	-	4	2	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	3	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Vermont												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Washington												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
United States												
All races, both sexes	14,503	10	1	1	-	4	12	17	76	211	661	1,278
Male	8,463	6	1	1	-	-	7	7	42	140	453	885
Female	6,040	4	-	-	-	4	5	10	34	71	208	393
White, both sexes	12,719	8	1	1	-	3	8	15	52	178	575	1,059
Male	7,387	5	1	1	-	-	4	7	29	119	391	740
Female	5,332	3	-	-	-	3	4	8	23	59	184	319
Black, both sexes	1,349	2	-	-	-	-	2	1	16	20	65	179
Male	850	1	-	-	-	-	2	-	8	12	48	117
Female	499	1	-	-	-	-	-	1	8	8	17	62
American Indian 1/, both sexes	186	-	-	-	-	1	1	1	6	9	12	26
Male	90	-	-	-	-	-	1	-	3	6	6	15
Female	96	-	-	-	-	1	-	1	3	3	6	11
Asian or Pacific Islander 2/, both sexes	249	-	-	-	-	-	1	-	2	4	9	14
Male	136	-	-	-	-	-	-	-	2	3	8	13
Female	113	-	-	-	-	-	1	-	-	1	1	1
Alabama												
All races, both sexes	331	-	-	-	-	-	-	1	4	5	13	31
Male	201	-	-	-	-	-	-	-	2	4	5	21
Female	130	-	-	-	-	-	-	1	2	1	8	10
White, both sexes	276	-	-	-	-	-	-	1	1	5	11	26
Male	164	-	-	-	-	-	-	-	1	4	4	18
Female	112	-	-	-	-	-	-	1	-	1	7	8
Black, both sexes	55	-	-	-	-	-	-	-	3	-	2	5
Male	37	-	-	-	-	-	-	-	1	-	1	3
Female	18	-	-	-	-	-	-	-	2	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Alaska												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	325	3	-	-	-	-	-	1	3	6	13	28
Male	190	2	-	-	-	-	-	-	2	4	10	16
Female	135	1	-	-	-	-	-	1	1	2	3	12
White, both sexes	283	3	-	-	-	-	-	-	1	3	12	20
Male	171	2	-	-	-	-	-	-	1	2	10	12
Female	112	1	-	-	-	-	-	-	-	1	2	8
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	32	-	-	-	-	-	-	1	2	3	1	6
Male	13	-	-	-	-	-	-	-	1	2	-	3
Female	19	-	-	-	-	-	-	1	1	1	1	3
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Arkansas												
All races, both sexes	161	-	-	-	-	-	-	-	4	3	3	16
Male	93	-	-	-	-	-	-	-	2	-	1	12
Female	68	-	-	-	-	-	-	-	2	3	2	4
White, both sexes	147	-	-	-	-	-	-	-	3	3	3	14
Male	86	-	-	-	-	-	-	-	2	-	1	11
Female	61	-	-	-	-	-	-	-	1	3	2	3
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,112	-	-	1	-	-	1	4	7	16	43	82
Male	611	-	-	1	-	-	-	3	6	10	30	56
Female	501	-	-	-	-	-	1	1	1	6	13	26
White, both sexes	963	-	-	1	-	-	1	4	6	16	38	70
Male	535	-	-	1	-	-	-	3	5	10	26	46
Female	428	-	-	-	-	-	1	1	1	6	12	24
Black, both sexes	52	-	-	-	-	-	-	-	-	-	2	4
Male	27	-	-	-	-	-	-	-	-	-	2	3
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	1	-	2	3
Male	8	-	-	-	-	-	-	-	1	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	82	-	-	-	-	-	-	-	-	-	1	5
Male	41	-	-	-	-	-	-	-	-	-	1	5
Female	41	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Colorado												
All races, both sexes	180	-	-	-	-	-	-	-	3	2	8	20
Male	109	-	-	-	-	-	-	-	3	2	7	14
Female	71	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	168	-	-	-	-	-	-	-	3	2	7	19
Male	102	-	-	-	-	-	-	-	3	2	6	14
Female	66	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	188	-	-	-	-	-	-	1	-	3	9	12
Male	107	-	-	-	-	-	-	1	-	2	7	9
Female	81	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	176	-	-	-	-	-	-	1	-	3	7	12
Male	99	-	-	-	-	-	-	1	-	2	6	9
Female	77	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Delaware												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	2	5
Male	23	-	-	-	-	-	-	-	-	-	1	3
Female	16	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	37	-	-	-	-	-	-	-	-	-	2	5
Male	22	-	-	-	-	-	-	-	-	-	1	3
Female	15	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Florida												
All races, both sexes	1,207	-	-	-	-	-	1	-	4	15	43	110
Male	702	-	-	-	-	-	1	-	2	9	30	74
Female	505	-	-	-	-	-	-	-	2	6	13	36
White, both sexes	1,120	-	-	-	-	-	-	-	2	14	39	96
Male	651	-	-	-	-	-	-	-	-	9	27	65
Female	469	-	-	-	-	-	-	-	2	5	12	31
Black, both sexes	79	-	-	-	-	-	1	-	2	1	2	14
Male	44	-	-	-	-	-	1	-	2	-	1	9
Female	35	-	-	-	-	-	-	-	-	1	1	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	461	-	-	-	-	-	-	-	-	13	27	50
Male	272	-	-	-	-	-	-	-	-	7	18	33
Female	189	-	-	-	-	-	-	-	-	6	9	17
White, both sexes	354	-	-	-	-	-	-	-	-	10	21	35
Male	210	-	-	-	-	-	-	-	-	4	13	23
Female	144	-	-	-	-	-	-	-	-	6	8	12
Black, both sexes	100	-	-	-	-	-	-	-	-	2	5	15
Male	58	-	-	-	-	-	-	-	-	2	4	10
Female	42	-	-	-	-	-	-	-	-	-	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Hawaii												
All races, both sexes	65	-	-	-	-	-	-	-	1	1	2	4
Male	40	-	-	-	-	-	-	-	1	1	1	3
Female	25	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	1	1	2	4
Male	25	-	-	-	-	-	-	-	1	1	1	3
Female	18	-	-	-	-	-	-	-	-	-	1	1
Idaho												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Illinois												
All races, both sexes	630	1	-	-	-	-	1	2	4	7	40	52
Male	357	1	-	-	-	-	1	1	2	4	25	38
Female	273	-	-	-	-	-	-	1	2	3	15	14
White, both sexes	528	1	-	-	-	-	1	2	4	5	31	39
Male	290	1	-	-	-	-	1	1	2	3	20	28
Female	238	-	-	-	-	-	-	1	2	2	11	11
Black, both sexes	89	-	-	-	-	-	-	-	-	1	7	11
Male	62	-	-	-	-	-	-	-	-	-	4	9
Female	27	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	2
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	326	1	-	-	-	-	-	-	-	6	13	31
Male	188	-	-	-	-	-	-	-	-	5	9	23
Female	138	1	-	-	-	-	-	-	-	1	4	8
White, both sexes	301	1	-	-	-	-	-	-	-	6	11	31
Male	171	-	-	-	-	-	-	-	-	5	7	23
Female	130	1	-	-	-	-	-	-	-	1	4	8
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Iowa												
All races, both sexes	100	-	1	-	-	-	-	-	-	1	2	8
Male	49	-	1	-	-	-	-	-	-	-	2	4
Female	51	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	97	-	1	-	-	-	-	-	-	1	2	8
Male	47	-	1	-	-	-	-	-	-	-	2	4
Female	50	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	108	-	-	-	-	-	-	-	-	-	3	15
Male	66	-	-	-	-	-	-	-	-	-	3	14
Female	42	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	104	-	-	-	-	-	-	-	-	-	3	14
Male	63	-	-	-	-	-	-	-	-	-	3	13
Female	41	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Kentucky												
All races, both sexes	251	1	-	-	-	-	-	-	1	5	13	22
Male	153	1	-	-	-	-	-	-	-	4	11	16
Female	98	-	-	-	-	-	-	-	1	1	2	6
White, both sexes	241	1	-	-	-	-	-	-	1	5	11	22
Male	144	1	-	-	-	-	-	-	-	4	9	16
Female	97	-	-	-	-	-	-	-	1	1	2	6
Black, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	251	-	-	-	-	-	-	-	3	4	11	31
Male	151	-	-	-	-	-	-	-	2	2	9	21
Female	100	-	-	-	-	-	-	-	1	2	2	10
White, both sexes	179	-	-	-	-	-	-	-	-	2	9	19
Male	107	-	-	-	-	-	-	-	-	1	7	13
Female	72	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	71	-	-	-	-	-	-	-	3	2	2	12
Male	44	-	-	-	-	-	-	-	2	1	2	8
Female	27	-	-	-	-	-	-	-	1	1	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Maine												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	1	3
Male	30	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	53	-	-	-	-	-	-	-	-	-	1	3
Male	30	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	293	-	-	-	-	-	-	1	1	6	9	24
Male	184	-	-	-	-	-	-	-	1	2	6	19
Female	109	-	-	-	-	-	-	1	-	4	3	5
White, both sexes	216	-	-	-	-	-	-	1	1	3	6	17
Male	128	-	-	-	-	-	-	-	1	1	4	13
Female	88	-	-	-	-	-	-	1	-	2	2	4
Black, both sexes	70	-	-	-	-	-	-	-	-	2	2	7
Male	54	-	-	-	-	-	-	-	-	1	1	6
Female	16	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Massachusetts												
All races, both sexes	358	-	-	-	-	-	-	-	3	4	17	26
Male	227	-	-	-	-	-	-	-	2	3	12	19
Female	131	-	-	-	-	-	-	-	1	1	5	7
White, both sexes	345	-	-	-	-	-	-	-	3	4	17	24
Male	219	-	-	-	-	-	-	-	2	3	12	17
Female	126	-	-	-	-	-	-	-	1	1	5	7
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	556	1	-	-	-	-	-	1	5	13	33	37
Male	326	1	-	-	-	-	-	-	1	7	19	22
Female	230	-	-	-	-	-	-	1	4	6	14	15
White, both sexes	468	-	-	-	-	-	-	1	3	8	25	27
Male	268	-	-	-	-	-	-	-	-	5	13	17
Female	200	-	-	-	-	-	-	1	3	3	12	10
Black, both sexes	79	1	-	-	-	-	-	-	2	4	6	10
Male	51	1	-	-	-	-	-	-	1	1	5	5
Female	28	-	-	-	-	-	-	-	1	3	1	5
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	2	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Minnesota												
All races, both sexes	146	-	-	-	-	-	1	-	4	-	8	15
Male	74	-	-	-	-	-	-	-	1	-	4	11
Female	72	-	-	-	-	-	1	-	3	-	4	4
White, both sexes	133	-	-	-	-	-	-	-	2	-	6	12
Male	68	-	-	-	-	-	-	-	1	-	2	9
Female	65	-	-	-	-	-	-	-	1	-	4	3
Black, both sexes	5	-	-	-	-	-	-	-	1	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	183	-	-	-	-	-	1	-	2	6	7	21
Male	114	-	-	-	-	-	1	-	1	5	6	15
Female	69	-	-	-	-	-	-	-	1	1	1	6
White, both sexes	142	-	-	-	-	-	1	-	1	4	6	12
Male	86	-	-	-	-	-	1	-	1	3	5	10
Female	56	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	38	-	-	-	-	-	-	-	1	2	1	7
Male	27	-	-	-	-	-	-	-	-	2	1	4
Female	11	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Missouri												
All races, both sexes	256	-	-	-	-	-	-	-	-	-	9	28
Male	157	-	-	-	-	-	-	-	-	-	6	19
Female	99	-	-	-	-	-	-	-	-	-	3	9
White, both sexes	230	-	-	-	-	-	-	-	-	-	8	24
Male	140	-	-	-	-	-	-	-	-	-	5	16
Female	90	-	-	-	-	-	-	-	-	-	3	8
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	4
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	40	-	-	-	-	-	-	-	1	1	2	2
Male	23	-	-	-	-	-	-	-	-	1	1	2
Female	17	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	2	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	1	-	2
Male	3	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Nebraska												
All races, both sexes	58	-	-	-	-	-	-	-	1	-	1	9
Male	34	-	-	-	-	-	-	-	1	-	1	7
Female	24	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	55	-	-	-	-	-	-	-	1	-	1	9
Male	32	-	-	-	-	-	-	-	1	-	1	7
Female	23	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	142	-	-	-	-	1	-	-	-	3	13	14
Male	95	-	-	-	-	-	-	-	-	2	9	11
Female	47	-	-	-	-	1	-	-	-	1	4	3
White, both sexes	124	-	-	-	-	1	-	-	-	3	11	11
Male	83	-	-	-	-	-	-	-	-	2	7	8
Female	41	-	-	-	-	1	-	-	-	1	4	3
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	3
Male	8	-	-	-	-	-	-	-	-	-	1	3
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
New Hampshire												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	5	4
Male	42	-	-	-	-	-	-	-	-	-	5	2
Female	18	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	60	-	-	-	-	-	-	-	-	-	5	4
Male	42	-	-	-	-	-	-	-	-	-	5	2
Female	18	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	445	2	-	-	-	-	-	-	1	10	31	39
Male	268	1	-	-	-	-	-	-	-	7	23	29
Female	177	1	-	-	-	-	-	-	1	3	8	10
White, both sexes	389	1	-	-	-	-	-	-	1	10	25	34
Male	240	1	-	-	-	-	-	-	-	7	18	26
Female	149	-	-	-	-	-	-	-	1	3	7	8
Black, both sexes	45	1	-	-	-	-	-	-	-	-	6	5
Male	21	-	-	-	-	-	-	-	-	-	5	3
Female	24	1	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
New Mexico												
All races, both sexes	126	-	-	-	-	-	-	-	2	2	8	11
Male	76	-	-	-	-	-	-	-	1	-	7	6
Female	50	-	-	-	-	-	-	-	1	2	1	5
White, both sexes	101	-	-	-	-	-	-	-	1	1	7	9
Male	63	-	-	-	-	-	-	-	-	-	6	6
Female	38	-	-	-	-	-	-	-	1	1	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	1	1	1	2
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	659	-	-	-	-	1	2	1	-	3	20	43
Male	397	-	-	-	-	-	1	-	-	2	18	35
Female	262	-	-	-	-	1	1	1	-	1	2	8
White, both sexes	570	-	-	-	-	1	2	1	-	3	20	34
Male	340	-	-	-	-	-	1	-	-	2	18	28
Female	230	-	-	-	-	1	1	1	-	1	2	6
Black, both sexes	67	-	-	-	-	-	-	-	-	-	-	6
Male	44	-	-	-	-	-	-	-	-	-	-	4
Female	23	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
North Carolina												
All races, both sexes	439	-	-	-	-	1	1	-	2	4	8	39
Male	250	-	-	-	-	-	1	-	-	2	5	27
Female	189	-	-	-	-	1	-	-	2	2	3	12
White, both sexes	367	-	-	-	-	1	-	-	2	3	6	33
Male	205	-	-	-	-	-	-	-	-	1	4	24
Female	162	-	-	-	-	1	-	-	2	2	2	9
Black, both sexes	58	-	-	-	-	-	-	-	-	1	2	6
Male	37	-	-	-	-	-	-	-	-	1	1	3
Female	21	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	12	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Ohio												
All races, both sexes	600	-	-	-	-	-	-	1	1	12	23	48
Male	336	-	-	-	-	-	-	-	-	6	14	32
Female	264	-	-	-	-	-	-	1	1	6	9	16
White, both sexes	533	-	-	-	-	-	-	-	1	10	18	41
Male	293	-	-	-	-	-	-	-	-	4	9	27
Female	240	-	-	-	-	-	-	-	1	6	9	14
Black, both sexes	64	-	-	-	-	-	-	1	-	1	5	7
Male	41	-	-	-	-	-	-	-	-	1	5	5
Female	23	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	188	-	-	-	-	1	-	-	-	2	15	16
Male	93	-	-	-	-	-	-	-	-	2	9	10
Female	95	-	-	-	-	1	-	-	-	-	6	6
White, both sexes	157	-	-	-	-	-	-	-	-	1	12	13
Male	78	-	-	-	-	-	-	-	-	1	8	9
Female	79	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	25	-	-	-	-	1	-	-	-	1	3	2
Male	12	-	-	-	-	-	-	-	-	1	1	1
Female	13	-	-	-	-	1	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Oregon												
All races, both sexes	68	-	-	-	-	-	-	-	-	1	2	4
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	41	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	62	-	-	-	-	-	-	-	-	1	2	4
Male	25	-	-	-	-	-	-	-	-	1	2	1
Female	37	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	769	-	-	-	-	-	-	-	1	7	31	70
Male	457	-	-	-	-	-	-	-	1	7	22	51
Female	312	-	-	-	-	-	-	-	-	-	9	19
White, both sexes	705	-	-	-	-	-	-	-	1	6	28	57
Male	420	-	-	-	-	-	-	-	1	6	21	41
Female	285	-	-	-	-	-	-	-	-	-	7	16
Black, both sexes	60	-	-	-	-	-	-	-	-	1	3	13
Male	36	-	-	-	-	-	-	-	-	1	1	10
Female	24	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Rhode Island												
All races, both sexes	67	-	-	-	-	-	-	-	-	2	3	3
Male	38	-	-	-	-	-	-	-	-	2	1	1
Female	29	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	64	-	-	-	-	-	-	-	-	2	3	3
Male	35	-	-	-	-	-	-	-	-	2	1	1
Female	29	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	229	-	-	-	-	-	-	-	1	2	8	19
Male	125	-	-	-	-	-	-	-	-	-	7	10
Female	104	-	-	-	-	-	-	-	1	2	1	9
White, both sexes	195	-	-	-	-	-	-	-	-	2	7	15
Male	101	-	-	-	-	-	-	-	-	-	6	8
Female	94	-	-	-	-	-	-	-	-	2	1	7
Black, both sexes	33	-	-	-	-	-	-	-	1	-	1	4
Male	23	-	-	-	-	-	-	-	-	-	1	2
Female	10	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
South Dakota												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	383	-	-	-	-	-	-	1	1	5	18	34
Male	227	-	-	-	-	-	-	1	1	5	12	21
Female	156	-	-	-	-	-	-	-	-	-	6	13
White, both sexes	341	-	-	-	-	-	-	1	1	5	15	29
Male	201	-	-	-	-	-	-	1	1	5	10	16
Female	140	-	-	-	-	-	-	-	-	-	5	13
Black, both sexes	39	-	-	-	-	-	-	-	-	-	3	5
Male	24	-	-	-	-	-	-	-	-	-	2	5
Female	15	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Texas												
All races, both sexes	1,603	1	-	-	-	-	3	2	12	28	99	174
Male	985	-	-	-	-	-	2	-	9	24	65	126
Female	618	1	-	-	-	-	1	2	3	4	34	48
White, both sexes	1,470	1	-	-	-	-	2	2	9	27	94	151
Male	904	-	-	-	-	-	1	-	6	23	62	114
Female	566	1	-	-	-	-	1	2	3	4	32	37
Black, both sexes	116	-	-	-	-	-	1	-	2	1	5	21
Male	70	-	-	-	-	-	1	-	2	1	3	10
Female	46	-	-	-	-	-	-	-	-	-	2	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	2
Male	11	-	-	-	-	-	-	-	1	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	63	-	-	-	-	-	-	-	-	1	-	5
Male	39	-	-	-	-	-	-	-	-	1	-	3
Female	24	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	61	-	-	-	-	-	-	-	-	1	-	4
Male	37	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Vermont												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	373	-	-	-	-	-	-	1	2	7	24	35
Male	210	-	-	-	-	-	-	1	-	4	17	24
Female	163	-	-	-	-	-	-	-	2	3	7	11
White, both sexes	303	-	-	-	-	-	-	1	2	4	23	26
Male	165	-	-	-	-	-	-	1	-	3	16	17
Female	138	-	-	-	-	-	-	-	2	1	7	9
Black, both sexes	65	-	-	-	-	-	-	-	-	2	1	9
Male	44	-	-	-	-	-	-	-	-	1	1	7
Female	21	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Washington												
All races, both sexes	112	-	-	-	-	-	-	-	-	1	1	4
Male	57	-	-	-	-	-	-	-	-	1	-	4
Female	55	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	105	-	-	-	-	-	-	-	-	1	1	4
Male	54	-	-	-	-	-	-	-	-	1	-	4
Female	51	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	137	-	-	-	-	-	-	-	-	2	8	8
Male	77	-	-	-	-	-	-	-	-	1	6	6
Female	60	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	132	-	-	-	-	-	-	-	-	2	7	8
Male	76	-	-	-	-	-	-	-	-	1	6	6
Female	56	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Wisconsin												
All races, both sexes	222	-	-	-	-	-	1	-	1	2	7	16
Male	113	-	-	-	-	-	-	-	-	1	3	10
Female	109	-	-	-	-	-	1	-	1	1	4	6
White, both sexes	200	-	-	-	-	-	1	-	1	2	7	13
Male	103	-	-	-	-	-	-	-	-	1	3	8
Female	97	-	-	-	-	-	1	-	1	1	4	5
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	23	-	-	-	-	-	-	-	1	-	1	2
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	23	-	-	-	-	-	-	-	1	-	1	2
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
United States												
All races, both sexes	1,725	1,833	1,555	1,512	1,687	1,719	1,296	667	205	28	4	1
Male	1,245	1,254	980	817	875	807	570	282	74	15	1	1
Female	480	579	575	695	812	912	726	385	131	13	3	-
White, both sexes	1,406	1,547	1,334	1,366	1,552	1,576	1,202	621	190	23	2	-
Male	1,041	1,053	838	734	812	741	524	260	72	15	-	-
Female	365	494	496	632	740	835	678	361	118	8	2	-
Black, both sexes	263	245	168	109	94	86	58	25	9	4	2	1
Male	174	175	110	65	52	46	26	10	2	-	1	1
Female	89	70	58	44	42	40	32	15	7	4	1	-
American Indian 1/, both sexes	33	20	25	15	16	11	6	3	1	-	-	-
Male	14	10	17	7	4	3	2	2	-	-	-	-
Female	19	10	8	8	12	8	4	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	23	21	28	22	25	46	30	18	5	1	-	-
Male	16	16	15	11	7	17	18	10	-	-	-	-
Female	7	5	13	11	18	29	12	8	5	1	-	-
Alabama												
All races, both sexes	30	45	43	41	42	29	27	18	2	-	-	-
Male	24	32	31	27	20	15	13	7	-	-	-	-
Female	6	13	12	14	22	14	14	11	2	-	-	-
White, both sexes	22	32	37	37	40	23	24	16	1	-	-	-
Male	19	21	27	23	19	11	11	6	-	-	-	-
Female	3	11	10	14	21	12	13	10	1	-	-	-
Black, both sexes	8	13	6	4	2	6	3	2	1	-	-	-
Male	5	11	4	4	1	4	2	1	-	-	-	-
Female	3	2	2	-	1	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Alaska												
All races, both sexes	3	1	4	1	5	-	2	-	-	-	-	-
Male	3	1	2	-	5	-	1	-	-	-	-	-
Female	-	-	2	1	-	-	1	-	-	-	-	-
White, both sexes	2	1	3	-	4	-	2	-	-	-	-	-
Male	2	1	1	-	4	-	1	-	-	-	-	-
Female	-	-	2	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	41	37	43	36	42	29	28	12	2	1	-	-
Male	29	22	25	24	24	16	12	3	-	1	-	-
Female	12	15	18	12	18	13	16	9	2	-	-	-
White, both sexes	36	32	37	33	38	28	25	12	2	1	-	-
Male	27	21	20	22	23	16	11	3	-	1	-	-
Female	9	11	17	11	15	12	14	9	2	-	-	-
Black, both sexes	1	1	2	2	1	-	1	-	-	-	-	-
Male	-	-	2	1	1	-	1	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	4	4	1	3	1	2	-	-	-	-	-
Male	2	1	3	1	-	-	-	-	-	-	-	-
Female	2	3	1	-	3	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Arkansas												
All races, both sexes	22	20	17	21	21	13	14	3	2	2	-	-
Male	14	14	10	13	12	6	5	2	1	1	-	-
Female	8	6	7	8	9	7	9	1	1	1	-	-
White, both sexes	19	18	16	18	20	13	13	3	2	2	-	-
Male	12	13	10	11	11	6	5	2	1	1	-	-
Female	7	5	6	7	9	7	8	1	1	1	-	-
Black, both sexes	2	2	1	1	1	-	1	-	-	-	-	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	136	153	110	130	119	129	110	47	19	5	-	-
Male	87	91	64	58	52	65	53	24	10	1	-	-
Female	49	62	46	72	67	64	57	23	9	4	-	-
White, both sexes	116	134	96	113	104	110	97	37	17	3	-	-
Male	76	81	58	50	47	56	46	20	9	1	-	-
Female	40	53	38	63	57	54	51	17	8	2	-	-
Black, both sexes	13	8	6	8	5	2	1	1	1	1	-	-
Male	8	4	2	3	3	1	-	-	1	-	-	-
Female	5	4	4	5	2	1	1	1	-	1	-	-
American Indian 1/, both sexes	1	5	2	-	-	1	-	-	-	-	-	-
Male	1	2	1	-	-	-	-	-	-	-	-	-
Female	-	3	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	6	6	9	10	16	12	9	1	1	-	-
Male	2	4	3	5	2	8	7	4	-	-	-	-
Female	4	2	3	4	8	8	5	5	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Colorado												
All races, both sexes	20	35	19	17	15	17	18	4	2	-	-	-
Male	15	23	14	6	7	7	8	2	1	-	-	-
Female	5	12	5	11	8	10	10	2	1	-	-	-
White, both sexes	17	34	17	14	15	16	18	4	2	-	-	-
Male	12	22	13	5	7	7	8	2	1	-	-	-
Female	5	12	4	9	8	9	10	2	1	-	-	-
Black, both sexes	3	-	-	1	-	-	-	-	-	-	-	-
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	2	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	17	16	18	15	22	29	25	12	9	-	-	-
Male	9	13	12	10	12	16	8	5	3	-	-	-
Female	8	3	6	5	10	13	17	7	6	-	-	-
White, both sexes	15	14	15	15	21	29	24	12	8	-	-	-
Male	8	11	9	10	11	16	8	5	3	-	-	-
Female	7	3	6	5	10	13	16	7	5	-	-	-
Black, both sexes	2	2	3	-	1	-	1	-	1	-	-	-
Male	1	2	3	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Delaware												
All races, both sexes	2	5	5	-	7	3	5	5	-	-	-	-
Male	2	4	4	-	1	2	4	2	-	-	-	-
Female	-	1	1	-	6	1	1	3	-	-	-	-
White, both sexes	1	5	5	-	6	3	5	5	-	-	-	-
Male	1	4	4	-	1	2	4	2	-	-	-	-
Female	-	1	1	-	5	1	1	3	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	6	2	1	1	3	1	2	1	1	-	-
Male	3	3	1	1	1	1	-	1	-	-	-	-
Female	1	3	1	-	-	2	1	1	1	1	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
Black, both sexes	4	6	1	1	1	2	-	-	1	-	-	-
Male	3	3	-	1	1	1	-	-	-	-	-	-
Female	1	3	1	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Florida												
All races, both sexes	151	167	118	118	137	160	107	57	16	3	-	-
Male	109	110	77	65	68	77	50	23	6	1	-	-
Female	42	57	41	53	69	83	57	34	10	2	-	-
White, both sexes	135	153	107	111	133	154	104	54	16	2	-	-
Male	98	99	73	62	66	73	50	22	6	1	-	-
Female	37	54	34	49	67	81	54	32	10	1	-	-
Black, both sexes	14	13	11	6	4	6	2	2	-	1	-	-
Male	9	10	4	2	2	4	-	-	-	-	-	-
Female	5	3	7	4	2	2	2	2	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	1	1	-	-	-	-
Male	1	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	60	69	53	52	39	52	27	13	5	-	1	-
Male	38	46	33	35	20	25	10	5	2	-	-	-
Female	22	23	20	17	19	27	17	8	3	-	1	-
White, both sexes	42	52	42	42	31	45	19	12	3	-	-	-
Male	27	35	26	31	15	22	7	5	2	-	-	-
Female	15	17	16	11	16	23	12	7	1	-	-	-
Black, both sexes	17	17	11	9	7	7	7	1	1	-	1	-
Male	10	11	7	4	5	3	2	-	-	-	-	-
Female	7	6	4	5	2	4	5	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Hawaii												
All races, both sexes	11	10	6	2	5	11	6	4	2	-	-	-
Male	7	8	6	2	2	5	2	2	-	-	-	-
Female	4	2	-	-	3	6	4	2	2	-	-	-
White, both sexes	7	5	2	1	3	1	1	1	1	-	-	-
Male	4	4	2	1	2	1	-	1	-	-	-	-
Female	3	1	-	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	4	1	2	10	5	3	1	-	-	-
Male	3	4	4	1	-	4	2	1	-	-	-	-
Female	1	1	-	-	2	6	3	2	1	-	-	-
Idaho												
All races, both sexes	1	1	3	3	2	3	3	4	1	-	-	-
Male	1	1	-	1	2	1	1	3	1	-	-	-
Female	-	-	3	2	-	2	2	1	-	-	-	-
White, both sexes	1	1	2	3	2	3	3	4	1	-	-	-
Male	1	1	-	1	2	1	1	3	1	-	-	-
Female	-	-	2	2	-	2	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Illinois												
All races, both sexes	73	82	67	59	61	76	61	31	12	-	1	-
Male	50	59	42	38	33	22	25	14	2	-	-	-
Female	23	23	25	21	28	54	36	17	10	-	1	-
White, both sexes	54	61	54	50	57	70	58	29	11	-	1	-
Male	39	42	33	32	31	19	23	13	2	-	-	-
Female	15	19	21	18	26	51	35	16	9	-	1	-
Black, both sexes	17	20	13	9	2	5	2	2	-	-	-	-
Male	11	16	9	6	1	3	2	1	-	-	-	-
Female	6	4	4	3	1	2	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	1	1	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Indiana												
All races, both sexes	43	40	39	27	43	41	21	17	2	2	-	-
Male	29	29	22	14	20	18	10	7	1	1	-	-
Female	14	11	17	13	23	23	11	10	1	1	-	-
White, both sexes	35	37	35	26	41	37	20	17	2	2	-	-
Male	24	26	21	13	18	16	9	7	1	1	-	-
Female	11	11	14	13	23	21	11	10	1	1	-	-
Black, both sexes	8	3	4	1	2	3	1	-	-	-	-	-
Male	5	3	1	1	2	2	1	-	-	-	-	-
Female	3	-	3	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Iowa												
All races, both sexes	7	9	12	14	9	16	14	6	1	-	-	-
Male	5	7	8	5	4	7	5	-	1	-	-	-
Female	2	2	4	9	5	9	9	6	-	-	-	-
White, both sexes	5	9	12	14	9	15	14	6	1	-	-	-
Male	3	7	8	5	4	7	5	-	1	-	-	-
Female	2	2	4	9	5	8	9	6	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	13	15	11	8	10	21	7	5	-	-	-	-
Male	11	9	6	4	4	10	2	3	-	-	-	-
Female	2	6	5	4	6	11	5	2	-	-	-	-
White, both sexes	13	14	9	8	10	21	7	5	-	-	-	-
Male	11	8	5	4	4	10	2	3	-	-	-	-
Female	2	6	4	4	6	11	5	2	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Kentucky												
All races, both sexes	26	33	34	30	29	21	21	11	4	-	-	-
Male	21	24	18	17	18	11	6	3	3	-	-	-
Female	5	9	16	13	11	10	15	8	1	-	-	-
White, both sexes	22	32	32	29	29	21	21	11	4	-	-	-
Male	17	23	17	16	18	11	6	3	3	-	-	-
Female	5	9	15	13	11	10	15	8	1	-	-	-
Black, both sexes	3	1	2	1	-	-	-	-	-	-	-	-
Male	3	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	44	35	28	26	26	26	11	4	1	-	1	-
Male	36	21	16	12	14	12	4	2	-	-	-	-
Female	8	14	12	14	12	14	7	2	1	-	1	-
White, both sexes	27	21	22	20	21	23	10	3	1	-	1	-
Male	24	15	10	10	11	11	4	1	-	-	-	-
Female	3	6	12	10	10	12	6	2	1	-	1	-
Black, both sexes	17	14	6	6	4	3	1	1	-	-	-	-
Male	12	6	6	2	3	1	-	1	-	-	-	-
Female	5	8	-	4	1	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Maine												
All races, both sexes	4	4	6	6	5	13	5	5	1	-	-	-
Male	4	4	4	3	1	8	2	3	-	-	-	-
Female	-	-	2	3	4	5	3	2	1	-	-	-
White, both sexes	4	4	6	6	5	13	5	5	1	-	-	-
Male	4	4	4	3	1	8	2	3	-	-	-	-
Female	-	-	2	3	4	5	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	42	41	37	33	28	36	27	5	1	-	1	1
Male	31	28	22	21	15	19	15	3	-	-	1	1
Female	11	13	15	12	13	17	12	2	1	-	-	-
White, both sexes	26	27	21	29	24	31	25	4	1	-	-	-
Male	19	16	11	17	13	17	14	2	-	-	-	-
Female	7	11	10	12	11	14	11	2	1	-	-	-
Black, both sexes	15	14	15	4	4	3	2	-	-	-	1	1
Male	12	12	11	4	2	2	1	-	-	-	1	1
Female	3	2	4	-	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Massachusetts												
All races, both sexes	49	41	33	30	37	56	38	19	2	3	-	-
Male	40	29	23	19	22	29	17	9	1	2	-	-
Female	9	12	10	11	15	27	21	10	1	1	-	-
White, both sexes	49	40	30	29	36	52	37	19	2	3	-	-
Male	40	29	21	19	21	27	16	9	1	2	-	-
Female	9	11	9	10	15	25	21	10	1	1	-	-
Black, both sexes	-	1	3	1	1	1	1	-	-	-	-	-
Male	-	-	2	-	1	1	1	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Michigan												
All races, both sexes	72	82	63	55	80	45	35	24	10	-	-	-
Male	55	65	38	29	42	23	12	9	3	-	-	-
Female	17	17	25	26	38	22	23	15	7	-	-	-
White, both sexes	54	63	55	53	73	42	32	22	10	-	-	-
Male	40	49	34	27	39	22	10	9	3	-	-	-
Female	14	14	21	26	34	20	22	13	7	-	-	-
Black, both sexes	15	18	7	2	7	3	2	2	-	-	-	-
Male	12	15	3	2	3	1	2	-	-	-	-	-
Female	3	3	4	-	4	2	-	2	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	1	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Minnesota												
All races, both sexes	16	8	9	26	19	11	16	8	5	-	-	-
Male	11	4	5	10	9	6	8	3	2	-	-	-
Female	5	4	4	16	10	5	8	5	3	-	-	-
White, both sexes	13	8	9	24	19	11	16	8	5	-	-	-
Male	9	4	5	10	9	6	8	3	2	-	-	-
Female	4	4	4	14	10	5	8	5	3	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	27	25	13	15	19	20	25	2	-	-	-	-
Male	23	20	10	6	10	8	8	1	-	-	-	-
Female	4	5	3	9	9	12	17	1	-	-	-	-
White, both sexes	17	17	12	13	17	18	23	1	-	-	-	-
Male	15	13	9	6	9	7	7	-	-	-	-	-
Female	2	4	3	7	8	11	16	1	-	-	-	-
Black, both sexes	10	8	1	1	2	2	2	1	-	-	-	-
Male	8	7	1	-	1	1	1	1	-	-	-	-
Female	2	1	-	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Missouri												
All races, both sexes	27	24	30	23	37	30	30	15	3	-	-	-
Male	21	14	21	14	23	15	14	7	3	-	-	-
Female	6	10	9	9	14	15	16	8	-	-	-	-
White, both sexes	26	19	27	23	35	27	25	13	3	-	-	-
Male	21	10	20	14	21	13	11	6	3	-	-	-
Female	5	9	7	9	14	14	14	7	-	-	-	-
Black, both sexes	1	5	3	-	2	3	4	2	-	-	-	-
Male	-	4	1	-	2	2	2	1	-	-	-	-
Female	1	1	2	-	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	1	2	7	8	4	2	5	1	-	-	-
Male	3	-	1	5	4	3	1	2	-	-	-	-
Female	1	1	1	2	4	1	1	3	1	-	-	-
White, both sexes	3	1	2	7	8	4	2	5	-	-	-	-
Male	3	-	1	5	4	3	1	2	-	-	-	-
Female	-	1	1	2	4	1	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Nebraska												
All races, both sexes	6	7	7	4	4	6	6	2	5	-	-	-
Male	6	6	3	2	1	4	1	1	1	-	-	-
Female	-	1	4	2	3	2	5	1	4	-	-	-
White, both sexes	6	5	6	4	4	6	6	2	5	-	-	-
Male	6	4	3	2	1	4	1	1	1	-	-	-
Female	-	1	3	2	3	2	5	1	4	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	24	18	14	13	10	11	4	1	-	-	-
Male	11	21	11	8	8	5	7	2	-	-	-	-
Female	5	3	7	6	5	5	4	2	1	-	-	-
White, both sexes	14	23	15	12	12	8	10	3	1	-	-	-
Male	10	20	9	7	7	5	6	2	-	-	-	-
Female	4	3	6	5	5	3	4	1	1	-	-	-
Black, both sexes	2	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	1	1	1	1	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
New Hampshire												
All races, both sexes	3	5	7	4	12	10	6	2	2	-	-	-
Male	2	4	5	4	9	6	3	-	2	-	-	-
Female	1	1	2	-	3	4	3	2	-	-	-	-
White, both sexes	3	5	7	4	12	10	6	2	2	-	-	-
Male	2	4	5	4	9	6	3	-	2	-	-	-
Female	1	1	2	-	3	4	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	35	45	46	40	51	54	51	29	9	2	-	-
Male	30	33	30	20	26	27	22	14	5	1	-	-
Female	5	12	16	20	25	27	29	15	4	1	-	-
White, both sexes	29	39	37	35	43	49	50	27	8	1	-	-
Male	27	30	25	16	23	26	22	14	4	1	-	-
Female	2	9	12	19	20	23	28	13	4	-	-	-
Black, both sexes	4	6	8	3	5	3	-	2	1	1	-	-
Male	1	3	5	2	1	-	-	-	1	-	-	-
Female	3	3	3	1	4	3	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	2	1	1	-	-	-	-	-
Male	2	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
New Mexico												
All races, both sexes	12	13	13	15	16	21	8	4	1	-	-	-
Male	11	7	10	5	11	12	5	1	-	-	-	-
Female	1	6	3	10	5	9	3	3	1	-	-	-
White, both sexes	8	9	10	13	13	19	7	3	1	-	-	-
Male	8	5	9	4	10	11	4	-	-	-	-	-
Female	-	4	1	9	3	8	3	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	4	3	2	3	2	1	1	-	-	-	-
Male	3	2	1	1	1	1	1	1	-	-	-	-
Female	1	2	2	1	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	68	72	68	79	88	90	72	41	9	2	-	-
Male	51	48	43	45	42	50	38	19	3	2	-	-
Female	17	24	25	34	46	40	34	22	6	-	-	-
White, both sexes	52	63	56	68	78	79	64	39	8	2	-	-
Male	42	40	34	37	38	44	34	17	3	2	-	-
Female	10	23	22	31	40	35	30	22	5	-	-	-
Black, both sexes	15	9	7	11	6	6	5	1	1	-	-	-
Male	9	8	6	8	3	3	2	1	-	-	-	-
Female	6	1	1	3	3	3	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	4	-	3	5	1	1	-	-	-	-
Male	-	-	2	-	1	3	1	1	-	-	-	-
Female	1	-	2	-	2	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
North Carolina												
All races, both sexes	50	56	51	56	55	59	34	15	8	-	-	-
Male	35	36	31	34	26	34	12	4	3	-	-	-
Female	15	20	20	22	29	25	22	11	5	-	-	-
White, both sexes	38	44	43	44	49	53	28	15	8	-	-	-
Male	28	27	25	28	23	29	9	4	3	-	-	-
Female	10	17	18	16	26	24	19	11	5	-	-	-
Black, both sexes	10	11	3	7	6	6	6	-	-	-	-	-
Male	7	8	2	4	3	5	3	-	-	-	-	-
Female	3	3	1	3	3	1	3	-	-	-	-	-
American Indian 1/, both sexes	2	-	4	5	-	-	-	-	-	-	-	-
Male	-	-	3	2	-	-	-	-	-	-	-	-
Female	2	-	1	3	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	-	-	2	3	1	4	1	-	-	-
Male	1	2	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	2	2	1	2	1	-	-	-
White, both sexes	2	2	-	-	1	2	1	4	1	-	-	-
Male	1	2	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	1	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Ohio												
All races, both sexes	76	71	66	61	68	76	54	32	11	-	-	-
Male	51	46	41	35	40	33	22	13	3	-	-	-
Female	25	25	25	26	28	43	32	19	8	-	-	-
White, both sexes	66	57	59	56	62	73	51	30	9	-	-	-
Male	43	40	36	32	36	31	21	11	3	-	-	-
Female	23	17	23	24	26	42	30	19	6	-	-	-
Black, both sexes	9	14	7	5	6	3	3	2	1	-	-	-
Male	7	6	5	3	4	2	1	2	-	-	-	-
Female	2	8	2	2	2	1	2	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Oklahoma												
All races, both sexes	18	23	31	22	25	13	11	7	4	-	-	-
Male	10	14	12	10	12	5	3	3	3	-	-	-
Female	8	9	19	12	13	8	8	4	1	-	-	-
White, both sexes	11	19	26	21	20	13	11	6	4	-	-	-
Male	7	11	9	10	10	5	3	2	3	-	-	-
Female	4	8	17	11	10	8	8	4	1	-	-	-
Black, both sexes	3	-	2	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	4	3	1	5	-	-	1	-	-	-	-
Male	1	3	2	-	2	-	-	1	-	-	-	-
Female	3	1	1	1	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Oregon												
All races, both sexes	4	5	7	4	10	13	12	5	-	1	-	-
Male	2	2	4	-	1	7	4	3	-	-	-	-
Female	2	3	3	4	9	6	8	2	-	1	-	-
White, both sexes	3	3	5	4	10	13	12	4	-	1	-	-
Male	2	2	3	-	1	7	4	2	-	-	-	-
Female	1	1	2	4	9	6	8	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	2	1	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	91	95	48	73	98	114	89	39	11	2	-	-
Male	67	68	28	40	60	52	39	17	4	1	-	-
Female	24	27	20	33	38	62	50	22	7	1	-	-
White, both sexes	79	85	41	67	96	110	85	38	11	1	-	-
Male	60	63	25	37	59	49	36	17	4	1	-	-
Female	19	22	16	30	37	61	49	21	7	-	-	-
Black, both sexes	12	10	5	5	2	4	3	1	-	1	-	-
Male	7	5	3	3	1	3	2	-	-	-	-	-
Female	5	5	2	2	1	1	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Rhode Island												
All races, both sexes	9	7	3	6	9	7	10	7	1	-	-	-
Male	6	6	1	4	4	3	4	6	-	-	-	-
Female	3	1	2	2	5	4	6	1	1	-	-	-
White, both sexes	7	6	3	6	9	7	10	7	1	-	-	-
Male	4	5	1	4	4	3	4	6	-	-	-	-
Female	3	1	2	2	5	4	6	1	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	25	24	31	22	43	18	18	15	3	-	-	-
Male	18	16	21	8	23	6	6	9	1	-	-	-
Female	7	8	10	14	20	12	12	6	2	-	-	-
White, both sexes	19	19	26	21	38	18	15	13	2	-	-	-
Male	13	11	17	7	19	6	5	8	1	-	-	-
Female	6	8	9	14	19	12	10	5	1	-	-	-
Black, both sexes	5	5	5	1	5	-	3	2	1	-	-	-
Male	4	5	4	1	4	-	1	1	-	-	-	-
Female	1	-	1	-	1	-	2	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
South Dakota												
All races, both sexes	1	1	2	3	2	6	4	1	1	-	-	-
Male	1	1	2	1	1	1	2	-	-	-	-	-
Female	-	-	-	2	1	5	2	1	1	-	-	-
White, both sexes	1	1	1	2	1	5	4	1	1	-	-	-
Male	1	1	1	-	1	1	2	-	-	-	-	-
Female	-	-	-	2	-	4	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	51	46	41	40	42	45	36	21	2	-	-	-
Male	41	37	30	23	13	18	16	8	1	-	-	-
Female	10	9	11	17	29	27	20	13	1	-	-	-
White, both sexes	44	41	37	35	37	41	32	21	2	-	-	-
Male	35	33	26	21	12	17	15	8	1	-	-	-
Female	9	8	11	14	25	24	17	13	1	-	-	-
Black, both sexes	6	4	4	5	4	4	4	-	-	-	-	-
Male	5	3	4	2	1	1	1	-	-	-	-	-
Female	1	1	-	3	3	3	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Texas												
All races, both sexes	221	215	189	163	167	153	105	55	15	1	-	-
Male	168	137	134	87	99	58	54	18	3	1	-	-
Female	53	78	55	76	68	95	51	37	12	-	-	-
White, both sexes	198	192	167	155	162	141	100	53	15	1	-	-
Male	155	123	117	82	97	53	50	17	3	1	-	-
Female	43	69	50	73	65	88	50	36	12	-	-	-
Black, both sexes	21	21	19	7	4	11	1	2	-	-	-	-
Male	12	13	15	5	2	4	1	1	-	-	-	-
Female	9	8	4	2	2	7	-	1	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	1	1	1	4	-	-	-	-	-
Male	1	1	2	-	-	1	3	-	-	-	-	-
Female	-	-	1	1	1	-	1	-	-	-	-	-
Utah												
All races, both sexes	1	7	6	9	8	13	10	2	1	-	-	-
Male	1	6	5	4	2	8	7	1	1	-	-	-
Female	-	1	1	5	6	5	3	1	-	-	-	-
White, both sexes	1	7	5	9	8	13	10	2	1	-	-	-
Male	1	6	4	4	2	8	7	1	1	-	-	-
Female	-	1	1	5	6	5	3	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Vermont												
All races, both sexes	-	-	1	1	1	2	3	-	1	1	-	-
Male	-	-	-	1	1	-	2	-	-	1	-	-
Female	-	-	1	-	-	2	1	-	1	-	-	-
White, both sexes	-	-	1	1	1	2	3	-	1	1	-	-
Male	-	-	-	1	1	-	2	-	-	1	-	-
Female	-	-	1	-	-	2	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	44	48	40	49	41	43	26	10	2	1	-	-
Male	28	32	27	22	20	20	9	5	-	1	-	-
Female	16	16	13	27	21	23	17	5	2	-	-	-
White, both sexes	28	36	34	42	33	38	24	9	2	1	-	-
Male	18	21	21	18	17	18	9	5	-	1	-	-
Female	10	15	13	24	16	20	15	4	2	-	-	-
Black, both sexes	16	12	5	6	8	3	2	1	-	-	-	-
Male	10	11	5	4	3	2	-	-	-	-	-	-
Female	6	1	-	2	5	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Washington												
All races, both sexes	9	14	6	10	19	15	16	13	4	-	-	-
Male	4	11	4	4	8	7	8	5	1	-	-	-
Female	5	3	2	6	11	8	8	8	3	-	-	-
White, both sexes	8	14	6	9	18	12	15	13	4	-	-	-
Male	3	11	4	3	8	7	7	5	1	-	-	-
Female	5	3	2	6	10	5	8	8	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	3	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	3	-	-	-	-	-	-
West Virginia												
All races, both sexes	11	22	15	16	20	18	10	5	2	-	-	-
Male	7	18	6	8	10	8	5	2	-	-	-	-
Female	4	4	9	8	10	10	5	3	2	-	-	-
White, both sexes	11	21	13	16	19	18	10	5	2	-	-	-
Male	7	17	6	8	10	8	5	2	-	-	-	-
Female	4	4	7	8	9	10	5	3	2	-	-	-
Black, both sexes	-	1	2	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Wisconsin												
All races, both sexes	25	24	30	22	23	32	15	16	7	1	-	-
Male	12	20	15	12	13	14	5	4	3	1	-	-
Female	13	4	15	10	10	18	10	12	4	-	-	-
White, both sexes	15	20	27	21	23	31	15	16	7	1	-	-
Male	11	16	14	11	13	13	5	4	3	1	-	-
Female	4	4	13	10	10	18	10	12	4	-	-	-
Black, both sexes	6	2	2	-	-	-	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	5	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	2	4	3	2	4	2	-	-	-	-	-
Male	1	2	2	1	-	1	-	-	-	-	-	-
Female	1	-	2	2	2	3	2	-	-	-	-	-
White, both sexes	2	2	4	3	2	4	2	-	-	-	-	-
Male	1	2	2	1	-	1	-	-	-	-	-	-
Female	1	-	2	2	2	3	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
United States												
All races, both sexes	9,253	86	26	14	14	23	37	63	110	206	500	885
Male	5,347	55	14	7	9	6	17	29	67	118	288	606
Female	3,906	31	12	7	5	17	20	34	43	88	212	279
White, both sexes	7,705	59	15	9	5	14	27	44	83	163	394	690
Male	4,496	35	8	4	3	5	13	20	58	98	234	477
Female	3,209	24	7	5	2	9	14	24	25	65	160	213
Black, both sexes	1,204	24	8	3	8	7	10	16	23	33	78	161
Male	662	18	4	1	5	1	4	6	8	14	40	103
Female	542	6	4	2	3	6	6	10	15	19	38	58
American Indian 1/, both sexes	150	-	2	-	1	1	-	1	4	5	22	18
Male	83	-	1	-	1	-	-	1	1	2	10	15
Female	67	-	1	-	-	1	-	-	3	3	12	3
Asian or Pacific Islander 2/, both sexes	194	3	1	2	-	1	-	2	-	5	6	16
Male	106	2	1	2	-	-	-	2	-	4	4	11
Female	88	1	-	-	-	1	-	-	-	1	2	5
Alabama												
All races, both sexes	196	1	-	1	-	2	-	1	4	3	16	16
Male	105	-	-	1	-	-	-	-	3	1	10	13
Female	91	1	-	-	-	2	-	1	1	2	6	3
White, both sexes	145	-	-	1	-	-	-	-	3	1	10	11
Male	84	-	-	1	-	-	-	-	3	1	7	10
Female	61	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	49	1	-	-	-	2	-	1	1	2	5	5
Male	20	-	-	-	-	-	-	-	-	-	2	3
Female	29	1	-	-	-	2	-	1	1	2	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Alaska												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	2	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	183	-	1	-	-	-	-	-	3	4	11	18
Male	105	-	-	-	-	-	-	-	3	3	6	13
Female	78	-	1	-	-	-	-	-	-	1	5	5
White, both sexes	161	-	-	-	-	-	-	-	3	4	10	14
Male	95	-	-	-	-	-	-	-	3	3	5	10
Female	66	-	-	-	-	-	-	-	-	1	5	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	1	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	1	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Arkansas												
All races, both sexes	117	-	1	-	-	-	-	-	-	4	8	8
Male	67	-	1	-	-	-	-	-	-	2	4	6
Female	50	-	-	-	-	-	-	-	-	2	4	2
White, both sexes	101	-	1	-	-	-	-	-	-	3	5	6
Male	60	-	1	-	-	-	-	-	-	1	3	6
Female	41	-	-	-	-	-	-	-	-	2	2	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	2	2
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	765	14	2	1	2	3	4	7	7	15	43	79
Male	460	10	1	-	1	-	2	4	2	10	28	63
Female	305	4	1	1	1	3	2	3	5	5	15	16
White, both sexes	646	13	2	1	2	2	4	7	7	12	34	62
Male	394	9	1	-	1	-	2	4	2	7	25	47
Female	252	4	1	1	1	2	2	3	5	5	9	15
Black, both sexes	49	-	-	-	-	-	-	-	-	1	6	9
Male	25	-	-	-	-	-	-	-	-	1	1	8
Female	24	-	-	-	-	-	-	-	-	-	5	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	1	1	2
Male	9	-	-	-	-	-	-	-	-	1	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	1	-	-	-	1	-	-	-	1	2	6
Male	32	1	-	-	-	-	-	-	-	1	1	6
Female	25	-	-	-	-	1	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Colorado												
All races, both sexes	183	1	2	-	1	-	-	2	5	8	11	25
Male	115	1	2	-	1	-	-	2	5	4	8	17
Female	68	-	-	-	-	-	-	-	-	4	3	8
White, both sexes	173	1	1	-	1	-	-	2	5	7	11	23
Male	109	1	1	-	1	-	-	2	5	4	8	15
Female	64	-	-	-	-	-	-	-	-	3	3	8
Black, both sexes	6	-	1	-	-	-	-	-	-	1	-	1
Male	3	-	1	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	149	1	1	-	-	-	1	1	2	2	10	7
Male	80	1	1	-	-	-	1	1	1	1	6	7
Female	69	-	-	-	-	-	-	-	1	1	4	-
White, both sexes	135	1	1	-	-	-	1	1	1	1	10	7
Male	73	1	1	-	-	-	1	1	1	1	6	7
Female	62	-	-	-	-	-	-	-	-	-	4	-
Black, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Delaware												
All races, both sexes	29	-	-	-	-	-	-	1	-	-	1	3
Male	19	-	-	-	-	-	-	1	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	25	-	-	-	-	-	-	1	-	-	1	3
Male	18	-	-	-	-	-	-	1	-	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	-	1	-	-	-	-	1	-	1	1	1
Male	12	-	1	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Florida												
All races, both sexes	541	7	-	-	-	2	3	2	5	4	22	60
Male	347	5	-	-	-	2	1	1	3	4	16	42
Female	194	2	-	-	-	-	2	1	2	-	6	18
White, both sexes	461	3	-	-	-	2	1	-	4	3	18	50
Male	295	1	-	-	-	2	-	-	3	3	12	35
Female	166	2	-	-	-	-	1	-	1	-	6	15
Black, both sexes	74	4	-	-	-	-	2	2	1	-	4	10
Male	47	4	-	-	-	-	1	1	-	-	4	7
Female	27	-	-	-	-	-	1	1	1	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	374	7	1	1	2	-	3	3	6	9	21	36
Male	228	5	1	-	1	-	2	1	5	6	12	26
Female	146	2	-	1	1	-	1	2	1	3	9	10
White, both sexes	264	1	-	1	-	-	3	3	1	5	18	20
Male	161	1	-	-	-	-	2	1	1	3	10	15
Female	103	-	-	1	-	-	1	2	-	2	8	5
Black, both sexes	105	6	-	-	2	-	-	-	5	4	3	16
Male	63	4	-	-	1	-	-	-	4	3	2	11
Female	42	2	-	-	1	-	-	-	1	1	1	5
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Hawaii												
All races, both sexes	51	1	-	1	-	-	-	1	-	-	1	4
Male	27	-	-	1	-	-	-	1	-	-	-	3
Female	24	1	-	-	-	-	-	-	-	-	1	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	1	-	1	-	-	-	1	-	-	1	3
Male	16	-	-	1	-	-	-	1	-	-	-	2
Female	14	1	-	-	-	-	-	-	-	-	1	1
Idaho												
All races, both sexes	18	1	-	-	-	-	-	-	-	-	1	2
Male	12	1	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	1	-	-	-	-	-	-	-	-	1	2
Male	12	1	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Illinois												
All races, both sexes	365	-	-	2	1	-	3	5	1	8	22	31
Male	211	-	-	2	1	-	1	4	1	6	16	22
Female	154	-	-	-	-	-	2	1	-	2	6	9
White, both sexes	295	-	-	-	-	-	1	4	1	8	17	22
Male	172	-	-	-	-	-	1	3	1	6	12	15
Female	123	-	-	-	-	-	-	1	-	2	5	7
Black, both sexes	61	-	-	1	1	-	2	1	-	-	4	9
Male	34	-	-	1	1	-	-	1	-	-	3	7
Female	27	-	-	-	-	-	2	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	1	-	-	-	-	-	-	1	-
Male	4	-	-	1	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	197	3	2	-	1	-	2	1	2	5	8	16
Male	103	2	-	-	1	-	-	-	1	4	5	7
Female	94	1	2	-	-	-	2	1	1	1	3	9
White, both sexes	170	2	2	-	-	-	2	1	2	5	5	13
Male	90	1	-	-	-	-	-	-	1	4	3	6
Female	80	1	2	-	-	-	2	1	1	1	2	7
Black, both sexes	24	1	-	-	1	-	-	-	-	-	2	2
Male	12	1	-	-	1	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Iowa												
All races, both sexes	46	-	-	-	-	-	-	-	1	1	1	3
Male	33	-	-	-	-	-	-	-	1	-	1	2
Female	13	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	44	-	-	-	-	-	-	-	1	1	1	2
Male	32	-	-	-	-	-	-	-	1	-	1	2
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	95	-	-	-	-	-	-	-	-	-	11	9
Male	55	-	-	-	-	-	-	-	-	-	10	6
Female	40	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	86	-	-	-	-	-	-	-	-	-	11	9
Male	53	-	-	-	-	-	-	-	-	-	10	6
Female	33	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Kentucky												
All races, both sexes	157	-	-	-	-	-	-	1	2	3	8	23
Male	98	-	-	-	-	-	-	-	2	3	6	18
Female	59	-	-	-	-	-	-	1	-	-	2	5
White, both sexes	144	-	-	-	-	-	-	-	2	3	8	22
Male	91	-	-	-	-	-	-	-	2	3	6	18
Female	53	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	12	-	-	-	-	-	-	1	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	212	1	-	1	-	1	1	3	3	4	12	25
Male	118	-	-	-	-	-	1	1	1	1	8	16
Female	94	1	-	1	-	1	-	2	2	3	4	9
White, both sexes	146	1	-	-	-	-	-	1	1	-	7	17
Male	82	-	-	-	-	-	-	1	1	-	3	10
Female	64	1	-	-	-	-	-	-	-	-	4	7
Black, both sexes	63	-	-	1	-	1	1	2	2	4	5	8
Male	34	-	-	-	-	-	1	-	-	1	5	6
Female	29	-	-	1	-	1	-	2	2	3	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Maine												
All races, both sexes	43	-	-	-	-	-	-	-	-	2	2	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	2	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	2	2	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	146	-	-	1	-	-	1	1	-	4	9	15
Male	92	-	-	1	-	-	1	1	-	2	4	9
Female	54	-	-	-	-	-	-	-	-	2	5	6
White, both sexes	90	-	-	1	-	-	1	-	-	2	4	7
Male	59	-	-	1	-	-	1	-	-	1	2	4
Female	31	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	52	-	-	-	-	-	-	1	-	2	5	8
Male	32	-	-	-	-	-	-	1	-	1	2	5
Female	20	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Massachusetts												
All races, both sexes	292	-	-	1	-	-	1	1	3	10	24	35
Male	170	-	-	1	-	-	-	1	2	5	15	21
Female	122	-	-	-	-	-	1	-	1	5	9	14
White, both sexes	275	-	-	1	-	-	1	1	3	10	24	32
Male	162	-	-	1	-	-	-	1	2	5	15	20
Female	113	-	-	-	-	-	1	-	1	5	9	12
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	324	1	3	-	1	-	2	1	7	6	18	24
Male	184	1	2	-	1	-	-	-	3	3	9	19
Female	140	-	1	-	-	-	2	1	4	3	9	5
White, both sexes	257	1	1	-	-	-	2	-	4	6	13	22
Male	146	1	1	-	-	-	-	-	2	3	6	17
Female	111	-	-	-	-	-	2	-	2	3	7	5
Black, both sexes	63	-	2	-	-	-	-	1	2	-	5	2
Male	37	-	1	-	-	-	-	-	1	-	3	2
Female	26	-	1	-	-	-	-	1	1	-	2	-
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Minnesota												
All races, both sexes	114	2	-	-	-	1	1	1	1	2	5	10
Male	61	2	-	-	-	-	1	1	1	1	1	6
Female	53	-	-	-	-	1	-	-	-	1	4	4
White, both sexes	106	2	-	-	-	1	1	1	1	2	4	9
Male	56	2	-	-	-	-	1	1	1	1	1	5
Female	50	-	-	-	-	1	-	-	-	1	3	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	150	1	-	-	-	-	2	1	5	5	4	24
Male	81	1	-	-	-	-	1	-	1	2	2	16
Female	69	-	-	-	-	-	1	1	4	3	2	8
White, both sexes	106	1	-	-	-	-	2	1	2	2	1	17
Male	60	1	-	-	-	-	1	-	1	1	1	12
Female	46	-	-	-	-	-	1	1	1	1	-	5
Black, both sexes	39	-	-	-	-	-	-	-	2	3	2	7
Male	20	-	-	-	-	-	-	-	-	1	1	4
Female	19	-	-	-	-	-	-	-	2	2	1	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Missouri												
All races, both sexes	192	3	1	-	-	-	1	-	2	5	12	20
Male	105	2	1	-	-	-	-	-	1	2	4	14
Female	87	1	-	-	-	-	1	-	1	3	8	6
White, both sexes	172	1	1	-	-	-	1	-	2	5	12	19
Male	95	1	1	-	-	-	-	-	1	2	4	13
Female	77	-	-	-	-	-	1	-	1	3	8	6
Black, both sexes	18	2	-	-	-	-	-	-	-	-	-	1
Male	9	1	-	-	-	-	-	-	-	-	-	1
Female	9	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	26	-	-	-	-	-	-	-	1	-	2	4
Male	16	-	-	-	-	-	-	-	1	-	2	3
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	4
Male	13	-	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Nebraska												
All races, both sexes	66	-	-	-	-	-	-	-	-	2	6	7
Male	41	-	-	-	-	-	-	-	-	-	5	4
Female	25	-	-	-	-	-	-	-	-	2	1	3
White, both sexes	59	-	-	-	-	-	-	-	-	2	5	6
Male	36	-	-	-	-	-	-	-	-	-	4	4
Female	23	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	127	-	-	-	-	-	-	1	1	4	3	15
Male	78	-	-	-	-	-	-	1	-	3	-	9
Female	49	-	-	-	-	-	-	-	1	1	3	6
White, both sexes	116	-	-	-	-	-	-	1	-	4	3	14
Male	73	-	-	-	-	-	-	1	-	3	-	8
Female	43	-	-	-	-	-	-	-	-	1	3	6
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
New Hampshire												
All races, both sexes	33	1	-	-	-	-	-	-	1	-	1	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	19	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	32	1	-	-	-	-	-	-	1	-	1	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	18	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	264	1	-	-	-	-	1	2	6	5	10	22
Male	156	-	-	-	-	-	-	-	6	1	6	17
Female	108	1	-	-	-	-	1	2	-	4	4	5
White, both sexes	212	1	-	-	-	-	-	1	5	3	9	14
Male	126	-	-	-	-	-	-	-	5	1	5	11
Female	86	1	-	-	-	-	-	1	-	2	4	3
Black, both sexes	43	-	-	-	-	-	1	1	1	2	-	7
Male	23	-	-	-	-	-	-	-	1	-	-	5
Female	20	-	-	-	-	-	1	1	-	2	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
New Mexico												
All races, both sexes	110	1	-	-	-	1	-	1	1	6	5	16
Male	64	-	-	-	-	-	-	1	-	3	3	14
Female	46	1	-	-	-	1	-	-	1	3	2	2
White, both sexes	77	1	-	-	-	1	-	-	-	5	2	12
Male	48	-	-	-	-	-	-	-	-	3	2	10
Female	29	1	-	-	-	1	-	-	-	2	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	31	-	-	-	-	-	-	1	1	1	3	4
Male	16	-	-	-	-	-	-	1	-	-	1	4
Female	15	-	-	-	-	-	-	-	1	1	2	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	437	6	1	-	1	1	1	2	7	9	15	31
Male	263	5	-	-	1	-	1	1	5	6	6	26
Female	174	1	1	-	-	1	-	1	2	3	9	5
White, both sexes	358	5	1	-	-	-	1	1	6	6	11	25
Male	211	4	-	-	-	-	1	-	4	4	4	21
Female	147	1	1	-	-	-	-	1	2	2	7	4
Black, both sexes	63	1	-	-	1	1	-	-	1	2	3	5
Male	43	1	-	-	1	-	-	-	1	1	2	5
Female	20	-	-	-	-	1	-	-	-	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	1	-	1	-	1
Male	8	-	-	-	-	-	-	1	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
North Carolina												
All races, both sexes	299	5	1	-	-	3	1	3	4	5	19	31
Male	172	3	-	-	-	2	-	1	1	4	8	22
Female	127	2	1	-	-	1	1	2	3	1	11	9
White, both sexes	241	4	-	-	-	2	1	3	2	4	15	26
Male	141	2	-	-	-	1	-	1	1	3	5	19
Female	100	2	-	-	-	1	1	2	1	1	10	7
Black, both sexes	53	1	1	-	-	1	-	-	2	1	4	5
Male	28	1	-	-	-	1	-	-	-	1	3	3
Female	25	-	1	-	-	-	-	-	2	-	1	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Ohio												
All races, both sexes	342	4	1	-	1	1	-	3	5	8	21	27
Male	186	4	1	-	-	-	-	-	3	4	10	13
Female	156	-	-	-	1	1	-	3	2	4	11	14
White, both sexes	294	3	1	-	1	-	-	2	4	6	16	20
Male	160	3	1	-	-	-	-	-	3	4	7	10
Female	134	-	-	-	1	-	-	2	1	2	9	10
Black, both sexes	44	1	-	-	-	1	-	1	1	1	4	7
Male	24	1	-	-	-	-	-	-	-	-	2	3
Female	20	-	-	-	-	1	-	1	1	1	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	132	1	2	1	1	2	-	1	1	3	11	8
Male	70	1	1	-	1	-	-	1	1	2	4	4
Female	62	-	1	1	-	2	-	-	-	1	7	4
White, both sexes	99	-	1	-	1	1	-	-	1	1	3	3
Male	52	-	-	-	1	-	-	-	1	1	2	2
Female	47	-	1	-	-	1	-	-	-	-	1	1
Black, both sexes	13	1	1	1	-	-	-	1	-	-	4	1
Male	8	1	1	-	-	-	-	1	-	-	2	-
Female	5	-	-	1	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	20	-	-	-	-	1	-	-	-	2	4	4
Male	10	-	-	-	-	-	-	-	-	1	-	2
Female	10	-	-	-	-	1	-	-	-	1	4	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Oregon												
All races, both sexes	76	-	-	-	-	-	1	-	1	-	-	7
Male	39	-	-	-	-	-	1	-	-	-	-	4
Female	37	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	69	-	-	-	-	-	1	-	1	-	-	7
Male	37	-	-	-	-	-	1	-	-	-	-	4
Female	32	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	606	2	-	-	-	1	1	4	3	19	34	56
Male	338	1	-	-	-	-	1	-	2	11	18	38
Female	268	1	-	-	-	1	-	4	1	8	16	18
White, both sexes	520	1	-	-	-	1	-	3	3	16	31	39
Male	292	-	-	-	-	-	-	-	2	10	17	28
Female	228	1	-	-	-	1	-	3	1	6	14	11
Black, both sexes	79	1	-	-	-	-	1	1	-	3	3	16
Male	42	1	-	-	-	-	1	-	-	1	1	10
Female	37	-	-	-	-	-	-	1	-	2	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Rhode Island												
All races, both sexes	34	1	-	-	-	-	-	1	-	2	2	2
Male	22	1	-	-	-	-	-	-	-	2	1	2
Female	12	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	31	-	-	-	-	-	-	1	-	2	2	2
Male	20	-	-	-	-	-	-	-	-	2	1	2
Female	11	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	179	3	3	-	1	-	1	1	2	1	13	18
Male	90	1	1	-	-	-	-	1	1	1	4	8
Female	89	2	2	-	1	-	1	-	1	-	9	10
White, both sexes	128	1	-	-	-	-	-	-	1	-	7	14
Male	69	-	-	-	-	-	-	-	1	-	3	7
Female	59	1	-	-	-	-	-	-	-	-	4	7
Black, both sexes	51	2	3	-	1	-	1	1	1	1	6	4
Male	21	1	1	-	-	-	-	1	-	1	1	1
Female	30	1	2	-	1	-	1	-	1	-	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
South Dakota												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	3
Male	11	-	-	-	-	-	-	-	-	1	-	3
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	141	2	-	-	-	-	-	1	2	2	9	13
Male	80	1	-	-	-	-	-	1	1	1	7	11
Female	61	1	-	-	-	-	-	-	1	1	2	2
White, both sexes	124	1	-	-	-	-	-	-	2	2	8	11
Male	69	1	-	-	-	-	-	-	1	1	6	9
Female	55	-	-	-	-	-	-	-	1	1	2	2
Black, both sexes	16	1	-	-	-	-	-	1	-	-	1	2
Male	10	-	-	-	-	-	-	1	-	-	1	2
Female	6	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Texas												
All races, both sexes	694	7	1	4	2	3	4	5	13	20	29	73
Male	389	2	-	1	1	1	1	3	8	10	17	48
Female	305	5	1	3	1	2	3	2	5	10	12	25
White, both sexes	609	6	1	4	-	2	2	5	11	17	25	58
Male	339	1	-	1	-	1	-	3	7	10	15	37
Female	270	5	1	3	-	1	2	2	4	7	10	21
Black, both sexes	74	-	-	-	2	1	2	-	2	3	2	14
Male	44	-	-	-	1	-	1	-	1	-	1	10
Female	30	-	-	-	1	1	1	-	1	3	1	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	53	-	-	-	-	-	-	-	-	2	2	7
Male	31	-	-	-	-	-	-	-	-	2	1	3
Female	22	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	52	-	-	-	-	-	-	-	-	2	2	7
Male	30	-	-	-	-	-	-	-	-	2	1	3
Female	22	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Vermont												
All races, both sexes	12	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	233	2	1	-	-	2	-	2	1	5	11	23
Male	137	1	-	-	-	1	-	-	-	2	6	14
Female	96	1	1	-	-	1	-	2	1	3	5	9
White, both sexes	182	1	1	-	-	2	-	1	1	4	6	18
Male	108	-	-	-	-	1	-	-	-	1	4	12
Female	74	1	1	-	-	1	-	1	1	3	2	6
Black, both sexes	49	1	-	-	-	-	-	1	-	1	5	5
Male	29	1	-	-	-	-	-	-	-	1	2	2
Female	20	-	-	-	-	-	-	1	-	-	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Washington												
All races, both sexes	110	3	1	-	-	-	1	-	1	2	6	6
Male	61	2	1	-	-	-	1	-	-	2	3	4
Female	49	1	-	-	-	-	-	-	1	-	3	2
White, both sexes	91	3	1	-	-	-	1	-	1	1	3	4
Male	50	2	1	-	-	-	1	-	-	1	2	2
Female	41	1	-	-	-	-	-	-	1	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	71	1	-	-	-	-	-	-	-	2	6	5
Male	41	1	-	-	-	-	-	-	-	1	4	1
Female	30	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	68	1	-	-	-	-	-	-	-	2	6	3
Male	40	1	-	-	-	-	-	-	-	1	4	1
Female	28	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Wisconsin												
All races, both sexes	169	2	-	-	-	-	1	1	-	2	8	11
Male	93	1	-	-	-	-	1	-	-	1	6	7
Female	76	1	-	-	-	-	-	1	-	1	2	4
White, both sexes	157	2	-	-	-	-	1	1	-	2	7	6
Male	85	1	-	-	-	-	1	-	-	1	5	4
Female	72	1	-	-	-	-	-	1	-	1	2	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	5
Male	5	-	-	-	-	-	-	-	-	-	-	3
Female	3	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	27	-	-	-	-	-	-	-	1	-	1	2
Male	15	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	25	-	-	-	-	-	-	-	1	-	1	2
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
United States												
All races, both sexes	1,206	1,207	972	902	856	787	710	381	202	54	9	3
Male	834	795	601	493	445	382	328	167	68	13	2	3
Female	372	412	371	409	411	405	382	214	134	41	7	-
White, both sexes	960	1,000	842	752	741	706	622	344	179	46	7	3
Male	672	663	528	414	398	349	289	148	65	11	1	3
Female	288	337	314	338	343	357	333	196	114	35	6	-
Black, both sexes	199	174	106	112	75	54	68	21	17	6	1	-
Male	135	108	61	60	34	20	29	9	1	1	-	-
Female	64	66	45	52	41	34	39	12	16	5	1	-
American Indian 1/, both sexes	29	15	11	15	13	7	4	1	-	1	-	-
Male	16	12	5	9	3	4	2	1	-	-	-	-
Female	13	3	6	6	10	3	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	18	18	13	23	27	20	16	15	6	1	1	-
Male	11	12	7	10	10	9	8	9	2	1	1	-
Female	7	6	6	13	17	11	8	6	4	-	-	-
Alabama												
All races, both sexes	24	32	21	16	13	20	16	6	2	2	-	-
Male	16	18	9	9	5	13	7	-	-	-	-	-
Female	8	14	12	7	8	7	9	6	2	2	-	-
White, both sexes	19	23	19	11	11	19	9	6	1	1	-	-
Male	11	13	9	7	5	13	4	-	-	-	-	-
Female	8	10	10	4	6	6	5	6	1	1	-	-
Black, both sexes	5	9	2	4	2	1	7	-	1	1	-	-
Male	5	5	-	2	-	-	3	-	-	-	-	-
Female	-	4	2	2	2	1	4	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Alaska												
All races, both sexes	3	2	4	2	1	1	1	-	1	-	-	-
Male	2	1	4	1	1	1	-	-	-	-	-	-
Female	1	1	-	1	-	-	1	-	1	-	-	-
White, both sexes	2	1	4	1	1	1	1	-	1	-	-	-
Male	1	-	4	-	1	1	-	-	-	-	-	-
Female	1	1	-	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	30	17	20	13	19	19	20	4	4	-	-	-
Male	21	11	8	7	10	7	12	2	2	-	-	-
Female	9	6	12	6	9	12	8	2	2	-	-	-
White, both sexes	26	16	16	11	16	18	19	4	4	-	-	-
Male	19	11	7	6	10	6	11	2	2	-	-	-
Female	7	5	9	5	6	12	8	2	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	4	2	3	1	1	-	-	-	-	-
Male	2	-	1	1	-	1	1	-	-	-	-	-
Female	1	1	3	1	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Arkansas												
All races, both sexes	23	14	15	12	9	8	7	4	3	1	-	-
Male	18	8	7	5	5	3	2	4	2	-	-	-
Female	5	6	8	7	4	5	5	-	1	1	-	-
White, both sexes	19	11	15	11	8	8	7	4	3	-	-	-
Male	15	6	7	5	5	3	2	4	2	-	-	-
Female	4	5	8	6	3	5	5	-	1	-	-	-
Black, both sexes	4	3	-	1	1	-	-	-	-	1	-	-
Male	3	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	103	118	88	65	59	54	64	27	7	3	-	-
Male	70	73	58	35	28	31	30	11	2	1	-	-
Female	33	45	30	30	31	23	34	16	5	2	-	-
White, both sexes	78	106	83	54	50	46	52	23	6	2	-	-
Male	53	70	56	32	24	26	24	9	2	-	-	-
Female	25	36	27	22	26	20	28	14	4	2	-	-
Black, both sexes	12	9	-	4	3	1	4	-	-	-	-	-
Male	8	3	-	1	1	-	2	-	-	-	-	-
Female	4	6	-	3	2	1	2	-	-	-	-	-
American Indian 1/, both sexes	5	-	1	3	-	-	-	-	-	-	-	-
Male	3	-	-	2	-	-	-	-	-	-	-	-
Female	2	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	3	4	4	6	7	8	4	1	1	-	-
Male	6	-	2	-	3	5	4	2	-	1	-	-
Female	2	3	2	4	3	2	4	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Colorado												
All races, both sexes	14	25	18	16	15	24	6	6	2	2	-	-
Male	11	20	9	9	5	14	3	1	1	2	-	-
Female	3	5	9	7	10	10	3	5	1	-	-	-
White, both sexes	13	25	17	15	15	21	6	6	2	2	-	-
Male	10	20	9	8	5	13	3	1	1	2	-	-
Female	3	5	8	7	10	8	3	5	1	-	-	-
Black, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	18	11	16	13	14	10	10	8	2	-	-
Male	12	11	5	9	8	7	5	3	1	-	-	-
Female	10	7	6	7	5	7	5	7	7	2	-	-
White, both sexes	16	16	11	14	13	12	10	10	8	2	-	-
Male	9	10	5	7	8	6	5	3	1	-	-	-
Female	7	6	6	7	5	6	5	7	7	2	-	-
Black, both sexes	6	1	-	2	-	2	-	-	-	-	-	-
Male	3	-	-	2	-	1	-	-	-	-	-	-
Female	3	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Delaware												
All races, both sexes	3	6	2	2	2	6	2	1	-	-	-	-
Male	1	5	1	1	2	3	1	1	-	-	-	-
Female	2	1	1	1	-	3	1	-	-	-	-	-
White, both sexes	2	5	2	1	2	5	2	1	-	-	-	-
Male	-	5	1	1	2	3	1	1	-	-	-	-
Female	2	-	1	-	-	2	1	-	-	-	-	-
Black, both sexes	1	1	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	5	1	1	2	2	1	1	-	-	-	-
Male	3	4	-	-	1	1	1	-	-	-	-	-
Female	1	1	1	1	1	1	-	1	-	-	-	-
White, both sexes	1	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	4	1	1	1	2	-	1	-	-	-	-
Male	3	4	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Florida												
All races, both sexes	69	81	54	53	57	43	37	22	15	5	-	-
Male	48	57	34	32	36	28	22	8	7	1	-	-
Female	21	24	20	21	21	15	15	14	8	4	-	-
White, both sexes	57	67	49	45	53	40	30	21	14	4	-	-
Male	38	49	31	28	34	26	17	8	7	1	-	-
Female	19	18	18	17	19	14	13	13	7	3	-	-
Black, both sexes	11	13	4	7	4	3	7	-	1	1	-	-
Male	9	7	2	3	2	2	5	-	-	-	-	-
Female	2	6	2	4	2	1	2	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	58	58	41	38	35	18	19	12	5	1	-	-
Male	40	42	25	20	22	9	9	2	-	-	-	-
Female	18	16	16	18	13	9	10	10	5	1	-	-
White, both sexes	39	43	33	26	28	15	13	11	3	1	-	-
Male	30	31	20	15	17	7	6	2	-	-	-	-
Female	9	12	13	11	11	8	7	9	3	1	-	-
Black, both sexes	19	15	6	12	6	2	6	1	2	-	-	-
Male	10	11	4	5	4	1	3	-	-	-	-	-
Female	9	4	2	7	2	1	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Hawaii												
All races, both sexes	6	3	4	2	8	6	7	4	2	1	-	-
Male	3	1	3	1	3	3	3	3	2	-	-	-
Female	3	2	1	1	5	3	4	1	-	1	-	-
White, both sexes	2	-	3	1	1	4	5	-	1	1	-	-
Male	1	-	2	1	-	3	2	-	1	-	-	-
Female	1	-	1	-	1	1	3	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	1	1	6	2	2	4	1	-	-	-
Male	2	1	1	-	3	-	1	3	1	-	-	-
Female	2	1	-	1	3	2	1	1	-	-	-	-
Idaho												
All races, both sexes	2	1	1	2	2	1	1	4	-	-	-	-
Male	-	1	-	1	1	1	1	3	-	-	-	-
Female	2	-	1	1	1	-	-	1	-	-	-	-
White, both sexes	2	1	1	2	2	1	1	4	-	-	-	-
Male	-	1	-	1	1	1	1	3	-	-	-	-
Female	2	-	1	1	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Illinois												
All races, both sexes	36	44	44	38	37	31	33	21	6	2	-	-
Male	23	30	32	17	17	13	16	8	2	-	-	-
Female	13	14	12	21	20	18	17	13	4	2	-	-
White, both sexes	31	34	36	31	31	23	27	21	6	2	-	-
Male	19	21	28	12	17	12	15	8	2	-	-	-
Female	12	13	8	19	14	11	12	13	4	2	-	-
Black, both sexes	4	8	8	6	5	6	6	-	-	-	-	-
Male	4	7	4	4	-	1	1	-	-	-	-	-
Female	-	1	4	2	5	5	5	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	2	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
Indiana												
All races, both sexes	32	28	22	24	14	15	11	9	2	-	-	-
Male	22	17	13	14	7	6	2	2	-	-	-	-
Female	10	11	9	10	7	9	9	7	2	-	-	-
White, both sexes	26	26	20	21	12	14	10	8	1	-	-	-
Male	17	17	12	13	7	6	2	1	-	-	-	-
Female	9	9	8	8	5	8	8	7	1	-	-	-
Black, both sexes	6	2	2	2	2	1	1	1	1	-	-	-
Male	5	-	1	1	-	-	-	1	-	-	-	-
Female	1	2	1	1	2	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Iowa												
All races, both sexes	6	6	1	6	4	5	6	2	3	1	-	-
Male	6	4	1	4	4	2	3	1	3	1	-	-
Female	-	2	-	2	-	3	3	1	-	-	-	-
White, both sexes	5	6	1	6	4	5	6	2	3	1	-	-
Male	5	4	1	4	4	2	3	1	3	1	-	-
Female	-	2	-	2	-	3	3	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	13	11	3	21	8	10	6	1	2	-	-	-
Male	6	8	3	14	3	2	3	-	-	-	-	-
Female	7	3	-	7	5	8	3	1	2	-	-	-
White, both sexes	13	9	3	17	7	8	6	1	2	-	-	-
Male	6	8	3	12	3	2	3	-	-	-	-	-
Female	7	1	-	5	4	6	3	1	2	-	-	-
Black, both sexes	-	2	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	2	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Kentucky												
All races, both sexes	23	17	20	13	17	10	12	3	4	1	-	-
Male	16	10	15	7	9	4	3	2	2	1	-	-
Female	7	7	5	6	8	6	9	1	2	-	-	-
White, both sexes	19	15	19	11	16	10	11	3	4	1	-	-
Male	12	9	14	7	8	4	3	2	2	1	-	-
Female	7	6	5	4	8	6	8	1	2	-	-	-
Black, both sexes	4	1	1	2	1	-	1	-	-	-	-	-
Male	4	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	25	21	26	22	27	22	8	8	1	1	-	-
Male	15	16	16	13	13	14	3	-	-	-	-	-
Female	10	5	10	9	14	8	5	8	1	1	-	-
White, both sexes	17	17	21	16	18	17	6	6	-	1	-	-
Male	11	12	13	10	8	10	3	-	-	-	-	-
Female	6	5	8	6	10	7	3	6	-	1	-	-
Black, both sexes	8	3	5	6	9	4	1	2	1	-	-	-
Male	4	3	3	3	5	3	-	-	-	-	-	-
Female	4	-	2	3	4	1	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Maine												
All races, both sexes	6	5	6	5	1	6	4	2	2	1	-	-
Male	6	4	5	2	1	5	1	2	2	-	-	-
Female	-	1	1	3	-	1	3	-	-	1	-	-
White, both sexes	6	5	6	5	1	6	4	2	2	1	-	-
Male	6	4	5	2	1	5	1	2	2	-	-	-
Female	-	1	1	3	-	1	3	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	20	22	15	13	14	7	16	7	-	-	1	-
Male	19	14	11	6	7	5	8	3	-	-	1	-
Female	1	8	4	7	7	2	8	4	-	-	-	-
White, both sexes	11	14	10	7	10	5	12	5	-	-	1	-
Male	11	9	7	3	7	3	6	3	-	-	1	-
Female	-	5	3	4	3	2	6	2	-	-	-	-
Black, both sexes	9	7	4	4	4	2	4	2	-	-	-	-
Male	8	5	3	3	-	2	2	-	-	-	-	-
Female	1	2	1	1	4	-	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Massachusetts												
All races, both sexes	38	52	22	23	27	22	16	11	5	1	-	-
Male	28	31	12	13	15	11	6	7	2	-	-	-
Female	10	21	10	10	12	11	10	4	3	1	-	-
White, both sexes	36	48	21	22	23	21	15	11	5	1	-	-
Male	27	28	12	13	14	10	5	7	2	-	-	-
Female	9	20	9	9	9	11	10	4	3	1	-	-
Black, both sexes	1	4	1	1	3	-	1	-	-	-	-	-
Male	1	3	-	-	1	-	1	-	-	-	-	-
Female	-	1	1	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	49	37	35	28	27	25	28	22	9	1	-	-
Male	31	26	26	14	14	11	11	12	1	-	-	-
Female	18	11	9	14	13	14	17	10	8	1	-	-
White, both sexes	37	27	29	21	24	20	27	17	5	1	-	-
Male	23	21	21	9	13	10	10	9	-	-	-	-
Female	14	6	8	12	11	10	17	8	5	1	-	-
Black, both sexes	12	10	6	7	2	5	1	4	4	-	-	-
Male	8	5	5	5	1	1	1	3	1	-	-	-
Female	4	5	1	2	1	4	-	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Minnesota												
All races, both sexes	16	10	10	7	10	10	16	6	6	-	-	-
Male	13	7	5	1	3	5	9	4	1	-	-	-
Female	3	3	5	6	7	5	7	2	5	-	-	-
White, both sexes	16	7	9	7	9	10	16	5	6	-	-	-
Male	13	5	4	1	3	5	9	3	1	-	-	-
Female	3	2	5	6	6	5	7	2	5	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	22	17	16	14	6	9	5	-	2	-	-
Male	11	13	6	10	9	3	3	2	-	1	-	-
Female	6	9	11	6	5	3	6	3	-	1	-	-
White, both sexes	8	16	14	11	12	6	9	4	-	-	-	-
Male	5	10	5	7	8	3	3	2	-	-	-	-
Female	3	6	9	4	4	3	6	2	-	-	-	-
Black, both sexes	7	6	3	5	1	-	-	1	-	2	-	-
Male	5	3	1	3	1	-	-	-	-	1	-	-
Female	2	3	2	2	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Missouri												
All races, both sexes	25	17	14	13	25	16	20	12	5	1	-	-
Male	18	11	6	8	14	10	8	3	3	-	-	-
Female	7	6	8	5	11	6	12	9	2	1	-	-
White, both sexes	21	15	11	11	23	15	18	12	4	1	-	-
Male	16	10	5	7	13	9	7	3	3	-	-	-
Female	5	5	6	4	10	6	11	9	1	1	-	-
Black, both sexes	4	2	2	2	2	-	2	-	1	-	-	-
Male	2	1	1	1	1	-	1	-	-	-	-	-
Female	2	1	1	1	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	5	-	4	1	3	1	2	-	-	-	-
Male	2	3	-	1	1	1	-	2	-	-	-	-
Female	1	2	-	3	-	2	1	-	-	-	-	-
White, both sexes	3	5	-	2	1	3	-	2	-	-	-	-
Male	2	3	-	-	1	1	-	2	-	-	-	-
Female	1	2	-	2	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Nebraska												
All races, both sexes	9	11	6	4	6	8	2	4	-	-	1	-
Male	8	8	4	2	3	6	1	-	-	-	-	-
Female	1	3	2	2	3	2	1	4	-	-	1	-
White, both sexes	8	7	6	4	6	8	2	4	-	-	1	-
Male	7	5	4	2	3	6	1	-	-	-	-	-
Female	1	2	2	2	3	2	1	4	-	-	1	-
Black, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	15	15	22	12	12	10	9	5	3	-	-	-
Male	8	9	15	8	8	9	4	2	2	-	-	-
Female	7	6	7	4	4	1	5	3	1	-	-	-
White, both sexes	14	11	21	10	12	10	9	5	2	-	-	-
Male	8	7	15	6	8	9	4	2	2	-	-	-
Female	6	4	6	4	4	1	5	3	-	-	-	-
Black, both sexes	1	2	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	2	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	-	-	-	1	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
New Hampshire												
All races, both sexes	5	6	3	3	1	2	4	4	1	-	1	-
Male	1	2	1	1	1	1	2	4	-	-	-	-
Female	4	4	2	2	-	1	2	-	1	-	1	-
White, both sexes	5	5	3	3	1	2	4	4	1	-	1	-
Male	1	2	1	1	1	1	2	4	-	-	-	-
Female	4	3	2	2	-	1	2	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	43	34	21	32	18	23	30	7	7	1	1	-
Male	33	19	14	18	10	9	17	3	2	1	-	-
Female	10	15	7	14	8	14	13	4	5	-	1	-
White, both sexes	31	31	16	24	18	21	27	6	4	1	-	-
Male	25	18	11	13	10	8	14	2	2	1	-	-
Female	6	13	5	11	8	13	13	4	2	-	-	-
Black, both sexes	11	2	4	7	-	2	2	-	2	-	1	-
Male	8	-	2	4	-	1	2	-	-	-	-	-
Female	3	2	2	3	-	1	-	-	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	-	-	1	1	1	-	-	-
Male	-	1	1	1	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
New Mexico												
All races, both sexes	17	11	7	9	15	4	10	4	1	1	-	-
Male	10	7	6	8	4	3	3	2	-	-	-	-
Female	7	4	1	1	11	1	7	2	1	1	-	-
White, both sexes	9	7	6	8	9	3	9	4	1	-	-	-
Male	7	3	6	7	3	2	3	2	-	-	-	-
Female	2	4	-	1	6	1	6	2	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	4	1	1	5	1	1	-	-	1	-	-
Male	3	4	-	1	1	1	-	-	-	-	-	-
Female	4	-	1	-	4	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	59	47	50	49	43	47	29	22	15	1	1	-
Male	45	35	29	25	18	27	15	12	6	-	-	-
Female	14	12	21	24	25	20	14	10	9	1	1	-
White, both sexes	46	40	38	42	34	41	25	21	13	1	1	-
Male	34	29	21	22	13	25	12	11	6	-	-	-
Female	12	11	17	20	21	16	13	10	7	1	1	-
Black, both sexes	11	7	12	3	8	3	2	1	2	-	-	-
Male	9	6	8	-	5	2	1	1	-	-	-	-
Female	2	1	4	3	3	1	1	-	2	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	1	3	2	-	-	-	-	-
Male	1	-	-	3	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	3	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
North Carolina												
All races, both sexes	34	37	36	30	26	25	22	12	5	-	-	-
Male	19	23	26	20	15	10	9	7	2	-	-	-
Female	15	14	10	10	11	15	13	5	3	-	-	-
White, both sexes	25	29	28	24	22	22	19	10	5	-	-	-
Male	18	19	18	16	14	8	8	6	2	-	-	-
Female	7	10	10	8	8	14	11	4	3	-	-	-
Black, both sexes	7	8	8	6	4	3	2	-	-	-	-	-
Male	-	4	8	4	1	2	-	-	-	-	-	-
Female	7	4	-	2	3	1	2	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	1	2	-	1	1	-	2	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	1	1	-	2	1	-
White, both sexes	-	1	1	1	2	-	1	1	-	2	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	1	1	-	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Ohio												
All races, both sexes	43	38	42	29	40	32	21	13	8	5	-	-
Male	29	22	29	15	17	15	13	7	4	-	-	-
Female	14	16	13	14	23	17	8	6	4	5	-	-
White, both sexes	34	32	37	28	36	31	20	11	8	4	-	-
Male	21	18	26	14	16	15	13	5	4	-	-	-
Female	13	14	11	14	20	16	7	6	4	4	-	-
Black, both sexes	9	6	4	1	3	1	1	2	-	1	-	-
Male	8	4	2	1	1	-	-	2	-	-	-	-
Female	1	2	2	-	2	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	18	17	13	12	10	14	5	8	2	2	-	-
Male	14	11	11	8	4	3	1	3	-	-	-	-
Female	4	6	2	4	6	11	4	5	2	2	-	-
White, both sexes	15	14	11	9	9	13	5	8	2	2	-	-
Male	12	8	9	6	3	3	1	3	-	-	-	-
Female	3	6	2	3	6	10	4	5	2	2	-	-
Black, both sexes	-	-	1	3	-	-	-	-	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	3	1	-	1	1	-	-	-	-	-	-
Male	2	3	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Oregon												
All races, both sexes	8	12	9	9	8	8	7	4	1	1	-	-
Male	5	9	4	5	4	1	2	2	1	1	-	-
Female	3	3	5	4	4	7	5	2	-	-	-	-
White, both sexes	7	11	9	7	6	7	7	4	1	1	-	-
Male	5	8	4	4	4	1	2	2	1	1	-	-
Female	2	3	5	3	2	6	5	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	72	78	60	53	55	60	54	32	18	1	1	2
Male	49	52	35	31	32	21	27	13	5	-	-	2
Female	23	26	25	22	23	39	27	19	13	1	1	-
White, both sexes	55	69	49	46	55	53	51	29	15	1	1	2
Male	37	45	30	26	32	21	26	12	4	-	-	2
Female	18	24	19	20	23	32	25	17	11	1	1	-
Black, both sexes	16	9	11	6	-	6	3	2	1	-	-	-
Male	11	7	5	4	-	-	1	-	-	-	-	-
Female	5	2	6	2	-	6	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	-	1	2	-	-	-
Male	1	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Rhode Island												
All races, both sexes	1	5	5	1	4	6	2	1	1	-	-	-
Male	1	3	4	-	2	4	1	1	-	-	-	-
Female	-	2	1	1	2	2	1	-	1	-	-	-
White, both sexes	1	4	5	1	4	5	2	1	1	-	-	-
Male	1	2	4	-	2	4	1	1	-	-	-	-
Female	-	2	1	1	2	1	1	-	1	-	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	24	32	15	23	13	9	12	4	3	1	-	-
Male	15	22	8	13	9	3	1	2	-	-	-	-
Female	9	10	7	10	4	6	11	2	3	1	-	-
White, both sexes	19	23	11	18	9	9	10	2	3	1	-	-
Male	12	16	7	13	5	3	1	1	-	-	-	-
Female	7	7	4	5	4	6	9	1	3	1	-	-
Black, both sexes	5	9	4	5	4	-	2	2	-	-	-	-
Male	3	6	1	-	4	-	-	1	-	-	-	-
Female	2	3	3	5	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
South Dakota												
All races, both sexes	2	3	3	1	-	3	-	1	-	1	1	-
Male	1	2	2	-	-	1	-	-	-	1	-	-
Female	1	1	1	1	-	2	-	1	-	-	1	-
White, both sexes	1	2	1	1	-	3	-	1	-	1	1	-
Male	1	2	1	-	-	1	-	-	-	1	-	-
Female	-	-	-	1	-	2	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	13	21	16	10	14	11	18	6	3	-	-	-
Male	9	13	11	5	7	5	4	2	2	-	-	-
Female	4	8	5	5	7	6	14	4	1	-	-	-
White, both sexes	10	17	15	10	14	11	15	5	3	-	-	-
Male	7	9	10	5	7	5	4	2	2	-	-	-
Female	3	8	5	5	7	6	11	3	1	-	-	-
Black, both sexes	3	3	1	-	-	-	3	1	-	-	-	-
Male	2	3	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Texas												
All races, both sexes	95	85	69	72	69	51	45	22	19	5	1	-
Male	71	55	40	30	40	22	20	11	5	2	1	-
Female	24	30	29	42	29	29	25	11	14	3	-	-
White, both sexes	86	75	64	61	60	48	39	21	19	5	-	-
Male	66	48	37	22	35	21	18	10	5	2	-	-
Female	20	27	27	39	25	27	21	11	14	3	-	-
Black, both sexes	9	10	5	8	7	3	5	1	-	-	-	-
Male	5	7	3	7	4	1	2	1	-	-	-	-
Female	4	3	2	1	3	2	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	-	1	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
Utah												
All races, both sexes	3	9	6	7	5	5	2	2	1	1	-	1
Male	3	4	3	6	2	2	2	1	-	1	-	1
Female	-	5	3	1	3	3	-	1	1	-	-	-
White, both sexes	3	8	6	7	5	5	2	2	1	1	-	1
Male	3	3	3	6	2	2	2	1	-	1	-	1
Female	-	5	3	1	3	3	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Vermont												
All races, both sexes	1	1	1	3	2	1	1	-	-	-	-	-
Male	-	1	1	1	2	1	1	-	-	-	-	-
Female	1	-	-	2	-	-	-	-	-	-	-	-
White, both sexes	1	1	1	3	2	1	1	-	-	-	-	-
Male	-	1	1	1	2	1	1	-	-	-	-	-
Female	1	-	-	2	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	40	22	30	28	13	22	22	1	5	3	-	-
Male	28	17	19	19	6	11	12	-	1	-	-	-
Female	12	5	11	9	7	11	10	1	4	3	-	-
White, both sexes	34	14	23	24	11	19	15	1	4	3	-	-
Male	25	12	15	16	4	9	8	-	1	-	-	-
Female	9	2	8	8	7	10	7	1	3	3	-	-
Black, both sexes	5	8	7	3	2	3	7	-	1	-	-	-
Male	3	5	4	3	2	2	4	-	-	-	-	-
Female	2	3	3	-	-	1	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Washington												
All races, both sexes	10	11	12	11	18	8	11	3	6	-	-	-
Male	8	10	6	5	6	1	7	2	3	-	-	-
Female	2	1	6	6	12	7	4	1	3	-	-	-
White, both sexes	8	10	10	10	13	8	10	2	6	-	-	-
Male	7	9	6	4	3	1	7	1	3	-	-	-
Female	1	1	4	6	10	7	3	1	3	-	-	-
Black, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	4	-	1	1	-	-	-	-
Male	-	-	-	1	2	-	-	1	-	-	-	-
Female	-	-	1	-	2	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	7	10	9	8	6	8	7	1	1	-	-	-
Male	2	10	6	5	2	4	3	1	1	-	-	-
Female	5	-	3	3	4	4	4	-	-	-	-	-
White, both sexes	7	9	9	8	6	8	7	1	1	-	-	-
Male	2	9	6	5	2	4	3	1	1	-	-	-
Female	5	-	3	3	4	4	4	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Wisconsin												
All races, both sexes	14	17	18	22	13	23	16	12	8	1	-	-
Male	12	11	9	13	7	9	9	6	1	-	-	-
Female	2	6	9	9	6	14	7	6	7	1	-	-
White, both sexes	13	16	16	21	13	22	16	12	8	1	-	-
Male	11	10	8	13	7	8	9	6	1	-	-	-
Female	2	6	8	8	6	14	7	6	7	1	-	-
Black, both sexes	-	1	2	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	7	3	2	2	3	3	-	-	-	-	-
Male	2	4	3	1	1	1	2	-	-	-	-	-
Female	1	3	-	1	1	2	1	-	-	-	-	-
White, both sexes	3	5	3	2	2	3	3	-	-	-	-	-
Male	2	3	3	1	1	1	2	-	-	-	-	-
Female	1	2	-	1	1	2	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
United States												
All races, both sexes	3,072	-	-	1	3	1	7	4	12	16	26	39
Male	1,363	-	-	-	1	-	5	3	5	7	10	13
Female	1,709	-	-	1	2	1	2	1	7	9	16	26
White, both sexes	2,655	-	-	1	3	1	5	4	9	11	17	24
Male	1,193	-	-	-	1	-	4	3	5	6	7	7
Female	1,462	-	-	1	2	1	1	1	4	5	10	17
Black, both sexes	325	-	-	-	-	-	2	-	3	4	8	13
Male	135	-	-	-	-	-	1	-	-	1	3	6
Female	190	-	-	-	-	-	1	-	3	3	5	7
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	1	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	65	-	-	-	-	-	-	-	1	1	1	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	1	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	54	-	-	-	-	-	1	-	-	-	2	-
Male	28	-	-	-	-	-	1	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	50	-	-	-	-	-	1	-	-	-	2	-
Male	26	-	-	-	-	-	1	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Arkansas												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	373	-	-	-	2	-	-	1	3	2	5	5
Male	172	-	-	-	1	-	-	1	3	2	2	2
Female	201	-	-	-	1	-	-	-	-	-	3	3
White, both sexes	311	-	-	-	2	-	-	1	3	2	4	5
Male	151	-	-	-	1	-	-	1	3	2	2	2
Female	160	-	-	-	1	-	-	-	-	-	2	3
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Colorado												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Florida												
All races, both sexes	200	-	-	-	-	-	-	-	-	1	1	3
Male	84	-	-	-	-	-	-	-	-	-	-	1
Female	116	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	170	-	-	-	-	-	-	-	-	1	1	2
Male	72	-	-	-	-	-	-	-	-	-	-	-
Female	98	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	63	-	-	-	-	-	-	-	-	1	1	2
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	47	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Illinois												
All races, both sexes	117	-	-	-	-	1	1	-	1	-	1	-
Male	50	-	-	-	-	-	1	-	1	-	1	-
Female	67	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	101	-	-	-	-	1	-	-	1	-	1	-
Male	39	-	-	-	-	-	-	-	1	-	1	-
Female	62	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Iowa												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	36	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Kentucky												
All races, both sexes	45	-	-	-	-	-	-	-	-	1	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	41	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	-	1
Male	28	-	-	-	-	-	-	-	-	1	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Maine												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	41	-	-	-	-	-	1	-	-	-	2	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	1	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Massachusetts												
All races, both sexes	54	-	-	-	-	-	-	-	-	1	-	-
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	-	-
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	67	-	-	-	-	-	-	-	1	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	1	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Minnesota												
All races, both sexes	56	-	-	-	1	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	1	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	32	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Missouri												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	1	2
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	64	-	-	-	-	-	-	-	-	-	1	2
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Nebraska												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	-	-	-	-	-	1	-	1	-	-	1
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	18	-	-	-	-	-	1	-	1	-	-	1
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	96	-	-	-	-	-	-	-	1	1	-	2
Male	46	-	-	-	-	-	-	-	1	1	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	85	-	-	-	-	-	-	-	1	1	-	1
Male	41	-	-	-	-	-	-	-	1	1	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
New Mexico												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	183	-	-	-	-	-	-	1	-	-	1	5
Male	79	-	-	-	-	-	-	-	-	-	1	1
Female	104	-	-	-	-	-	-	1	-	-	-	4
White, both sexes	145	-	-	-	-	-	-	1	-	-	-	1
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	78	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	29	-	-	-	-	-	-	-	-	-	1	4
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
North Carolina												
All races, both sexes	105	-	-	-	-	-	-	1	1	1	-	1
Male	42	-	-	-	-	-	-	1	-	-	-	-
Female	63	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	83	-	-	-	-	-	-	1	-	-	-	1
Male	35	-	-	-	-	-	-	1	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	22	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Ohio												
All races, both sexes	116	-	-	-	-	-	-	1	-	1	1	-
Male	50	-	-	-	-	-	-	1	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	102	-	-	-	-	-	-	1	-	1	-	-
Male	46	-	-	-	-	-	-	1	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	52	-	-	-	-	-	1	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	45	-	-	-	-	-	1	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Oregon												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	164	-	-	-	-	-	-	-	-	1	2	2
Male	68	-	-	-	-	-	-	-	-	-	-	1
Female	96	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	156	-	-	-	-	-	-	-	-	-	1	1
Male	66	-	-	-	-	-	-	-	-	-	-	1
Female	90	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Rhode Island												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
South Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Texas												
All races, both sexes	208	-	-	1	-	-	1	-	-	1	3	1
Male	92	-	-	-	-	-	1	-	-	1	1	-
Female	116	-	-	1	-	-	-	-	-	-	2	1
White, both sexes	182	-	-	1	-	-	1	-	-	1	3	1
Male	78	-	-	-	-	-	1	-	-	1	1	-
Female	104	-	-	1	-	-	-	-	-	-	2	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	75	-	-	-	-	-	-	-	-	-	-	-
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Washington												
All races, both sexes	75	-	-	-	-	-	-	-	1	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	1	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Wisconsin												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
United States												
All races, both sexes	74	95	122	170	278	428	627	585	402	152	30	-
Male	46	45	73	107	164	220	275	232	117	34	6	-
Female	28	50	49	63	114	208	352	353	285	118	24	-
White, both sexes	50	75	100	128	231	375	561	530	366	135	29	-
Male	30	38	58	85	138	197	256	211	110	32	5	-
Female	20	37	42	43	93	178	305	319	256	103	24	-
Black, both sexes	18	18	17	36	36	42	51	39	26	11	1	-
Male	11	6	11	18	20	22	15	14	5	1	1	-
Female	7	12	6	18	16	20	36	25	21	10	-	-
American Indian 1/, both sexes	1	1	4	2	2	1	6	4	1	2	-	-
Male	1	-	3	1	1	-	2	2	-	-	-	-
Female	-	1	1	1	1	1	4	2	1	2	-	-
Asian or Pacific Islander 2/, both sexes	5	1	1	4	9	10	9	12	9	4	-	-
Male	4	1	1	3	5	1	2	5	2	1	-	-
Female	1	-	-	1	4	9	7	7	7	3	-	-
Alabama												
All races, both sexes	-	2	3	4	5	11	15	15	5	2	-	-
Male	-	1	2	-	3	5	8	6	1	1	-	-
Female	-	1	1	4	2	6	7	9	4	1	-	-
White, both sexes	-	1	2	3	3	10	14	15	5	2	-	-
Male	-	1	2	-	2	5	8	6	1	1	-	-
Female	-	-	-	3	1	5	6	9	4	1	-	-
Black, both sexes	-	1	1	1	2	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Alaska												
All races, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	5	9	5	5	10	5	8	2	1	-
Male	-	1	3	8	2	2	6	2	2	-	-	-
Female	-	-	2	1	3	3	4	3	6	2	1	-
White, both sexes	-	1	4	9	5	5	8	5	7	2	1	-
Male	-	1	3	8	2	2	4	2	2	-	-	-
Female	-	-	1	1	3	3	4	3	5	2	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Arkansas												
All races, both sexes	-	3	4	3	6	8	9	7	3	-	-	-
Male	-	2	2	-	3	2	4	2	1	-	-	-
Female	-	1	2	3	3	6	5	5	2	-	-	-
White, both sexes	-	2	3	3	4	7	8	7	2	-	-	-
Male	-	1	1	-	2	1	4	2	-	-	-	-
Female	-	1	2	3	2	6	4	5	2	-	-	-
Black, both sexes	-	1	1	-	2	1	1	-	1	-	-	-
Male	-	1	1	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	16	16	23	36	46	71	72	50	18	2	-
Male	2	6	10	12	23	23	36	28	16	5	-	-
Female	3	10	6	11	13	23	35	44	34	13	2	-
White, both sexes	5	11	13	17	30	37	61	62	43	13	2	-
Male	2	5	8	9	19	21	31	25	15	5	-	-
Female	3	6	5	8	11	16	30	37	28	8	2	-
Black, both sexes	-	4	2	3	2	3	6	2	2	1	-	-
Male	-	1	1	1	1	1	3	-	-	-	-	-
Female	-	3	1	2	1	2	3	2	2	1	-	-
American Indian 1/, both sexes	-	1	1	1	2	-	1	-	-	1	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	6	3	8	5	3	-	-
Male	-	-	-	2	2	1	2	3	1	-	-	-
Female	-	-	-	-	-	5	1	5	4	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Colorado												
All races, both sexes	-	-	2	1	3	5	7	6	6	2	-	-
Male	-	-	1	1	1	2	6	2	1	-	-	-
Female	-	-	1	-	2	3	1	4	5	2	-	-
White, both sexes	-	-	2	1	3	5	7	6	6	2	-	-
Male	-	-	1	1	1	2	6	2	1	-	-	-
Female	-	-	1	-	2	3	1	4	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	2	-	2	8	12	5	1	1	1	-
Male	-	-	-	-	1	4	-	2	-	-	-	-
Female	-	-	2	-	1	4	12	3	1	1	1	-
White, both sexes	-	-	2	-	2	7	11	5	1	1	1	-
Male	-	-	-	-	1	3	-	2	-	-	-	-
Female	-	-	2	-	1	4	11	3	1	1	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Delaware												
All races, both sexes	1	-	-	-	-	-	3	1	-	-	-	-
Male	1	-	-	-	-	-	3	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	3	-	-	-	-	-
Male	1	-	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	1	-	-	1	1	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	1	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Florida												
All races, both sexes	1	5	9	11	19	30	45	30	26	14	5	-
Male	1	3	7	6	10	14	19	12	6	3	2	-
Female	-	2	2	5	9	16	26	18	20	11	3	-
White, both sexes	-	5	7	6	16	25	39	29	22	12	5	-
Male	-	3	5	3	8	13	19	11	5	3	2	-
Female	-	2	2	3	8	12	20	18	17	9	3	-
Black, both sexes	1	-	2	5	2	3	6	1	3	2	-	-
Male	1	-	2	3	1	1	-	1	-	-	-	-
Female	-	-	-	2	1	2	6	-	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Georgia												
All races, both sexes	-	2	3	3	6	8	13	12	9	3	-	-
Male	-	1	2	3	5	4	5	3	4	2	-	-
Female	-	1	1	-	1	4	8	9	5	1	-	-
White, both sexes	-	-	2	3	5	5	12	9	7	3	-	-
Male	-	-	1	3	4	2	5	3	4	2	-	-
Female	-	-	1	-	1	3	7	6	3	1	-	-
Black, both sexes	-	2	1	-	1	3	1	3	2	-	-	-
Male	-	1	1	-	1	2	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	1	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Idaho												
All races, both sexes	1	1	-	-	-	2	6	1	1	-	-	-
Male	1	1	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	5	-	1	-	-	-
White, both sexes	1	1	-	-	-	2	6	1	1	-	-	-
Male	1	1	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	5	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Illinois												
All races, both sexes	4	4	2	5	9	13	30	27	11	6	2	-
Male	2	1	2	3	6	7	11	10	4	1	-	-
Female	2	3	-	2	3	6	19	17	7	5	2	-
White, both sexes	3	3	-	3	8	11	25	27	10	6	2	-
Male	1	-	-	2	5	6	9	10	3	1	-	-
Female	2	3	-	1	3	5	16	17	7	5	2	-
Black, both sexes	1	1	1	2	1	2	3	-	1	-	-	-
Male	1	1	1	1	1	1	2	-	1	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Indiana												
All races, both sexes	2	2	1	7	3	7	17	15	2	1	-	-
Male	-	-	1	4	3	3	11	4	-	1	-	-
Female	2	2	-	3	-	4	6	11	2	-	-	-
White, both sexes	2	2	1	6	3	7	13	15	2	1	-	-
Male	-	-	1	3	3	3	10	4	-	1	-	-
Female	2	2	-	3	-	4	3	11	2	-	-	-
Black, both sexes	-	-	-	1	-	-	4	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Iowa												
All races, both sexes	-	2	-	-	1	3	5	5	5	5	1	-
Male	-	1	-	-	1	2	1	3	2	1	-	-
Female	-	1	-	-	-	1	4	2	3	4	1	-
White, both sexes	-	2	-	-	1	3	5	5	5	5	1	-
Male	-	1	-	-	1	2	1	3	2	1	-	-
Female	-	1	-	-	-	1	4	2	3	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	1	2	3	4	9	6	6	3	-	-
Male	-	-	1	1	-	3	3	3	1	-	-	-
Female	1	-	-	1	3	1	6	3	5	3	-	-
White, both sexes	1	-	1	2	2	3	9	6	6	3	-	-
Male	-	-	1	1	-	2	3	3	1	-	-	-
Female	1	-	-	1	2	1	6	3	5	3	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Kentucky												
All races, both sexes	1	1	4	1	6	4	17	6	3	-	-	-
Male	1	1	2	1	2	2	6	3	2	-	-	-
Female	-	-	2	-	4	2	11	3	1	-	-	-
White, both sexes	-	1	3	1	5	4	17	6	3	-	-	-
Male	-	1	2	1	2	2	6	3	2	-	-	-
Female	-	-	1	-	3	2	11	3	1	-	-	-
Black, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	5	3	6	12	12	11	7	3	-	-
Male	1	1	2	2	2	7	6	3	2	-	-	-
Female	-	1	3	1	4	5	6	8	5	3	-	-
White, both sexes	-	1	4	2	4	7	8	6	6	3	-	-
Male	-	-	2	1	1	4	3	1	2	-	-	-
Female	-	1	2	1	3	3	5	5	4	3	-	-
Black, both sexes	1	1	1	1	2	5	4	5	1	-	-	-
Male	1	1	-	1	1	3	3	2	-	-	-	-
Female	-	-	1	-	1	2	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Maine												
All races, both sexes	1	1	-	1	2	8	2	5	3	-	-	-
Male	1	-	-	-	1	5	-	3	1	-	-	-
Female	-	1	-	1	1	3	2	2	2	-	-	-
White, both sexes	1	1	-	1	2	8	2	5	3	-	-	-
Male	1	-	-	-	1	5	-	3	1	-	-	-
Female	-	1	-	1	1	3	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	4	2	5	7	5	9	3	2	-	-
Male	1	-	4	-	4	5	3	3	2	-	-	-
Female	-	-	-	2	1	2	2	6	1	2	-	-
White, both sexes	-	-	3	-	2	7	5	8	2	1	-	-
Male	-	-	3	-	1	5	3	3	1	-	-	-
Female	-	-	-	-	1	2	2	5	1	1	-	-
Black, both sexes	1	-	1	2	3	-	-	-	1	1	-	-
Male	1	-	1	-	3	-	-	-	1	-	-	-
Female	-	-	-	2	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Massachusetts												
All races, both sexes	-	-	2	-	4	12	12	8	11	2	2	-
Male	-	-	1	-	3	4	6	2	4	2	1	-
Female	-	-	1	-	1	8	6	6	7	-	1	-
White, both sexes	-	-	2	-	4	12	11	8	10	2	2	-
Male	-	-	1	-	3	4	6	2	4	2	1	-
Female	-	-	1	-	1	8	5	6	6	-	1	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	4	4	8	6	8	9	16	7	3	1	-
Male	-	1	2	6	3	6	1	5	4	-	-	-
Female	-	3	2	2	3	2	8	11	3	3	1	-
White, both sexes	-	3	2	5	6	3	7	14	7	3	1	-
Male	-	1	1	4	3	2	1	5	4	-	-	-
Female	-	2	1	1	3	1	6	9	3	3	1	-
Black, both sexes	-	1	1	3	-	5	2	2	-	-	-	-
Male	-	-	-	2	-	4	-	-	-	-	-	-
Female	-	1	1	1	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Minnesota												
All races, both sexes	2	2	1	4	6	6	8	13	10	2	-	-
Male	1	1	1	4	4	4	4	7	2	1	-	-
Female	1	1	-	-	2	2	4	6	8	1	-	-
White, both sexes	2	2	1	4	5	6	8	13	10	2	-	-
Male	1	1	1	4	4	4	4	7	2	1	-	-
Female	1	1	-	-	1	2	4	6	8	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	1	-	3	3	3	4	7	5	2	-	-
Male	2	1	-	1	2	-	1	2	2	1	-	-
Female	-	-	-	2	1	3	3	5	3	1	-	-
White, both sexes	1	1	-	2	2	2	3	4	5	2	-	-
Male	1	1	-	1	1	-	1	2	2	1	-	-
Female	-	-	-	1	1	2	2	2	3	1	-	-
Black, both sexes	1	-	-	1	1	1	1	3	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Missouri												
All races, both sexes	2	2	2	-	5	14	12	10	12	5	1	-
Male	2	1	2	-	3	7	3	4	4	1	1	-
Female	-	1	-	-	2	7	9	6	8	4	-	-
White, both sexes	1	2	2	-	5	14	11	10	11	5	-	-
Male	1	1	2	-	3	7	3	4	4	1	-	-
Female	-	1	-	-	2	7	8	6	7	4	-	-
Black, both sexes	1	-	-	-	-	-	1	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	1	-	1	4	1	1	1	-	-
Male	2	-	-	-	-	1	3	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	1	1	-	-
White, both sexes	2	-	-	1	-	1	4	1	1	1	-	-
Male	2	-	-	-	-	1	3	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Nebraska												
All races, both sexes	-	-	2	-	3	4	3	2	2	4	-	-
Male	-	-	1	-	2	2	2	1	-	1	-	-
Female	-	-	1	-	1	2	1	1	2	3	-	-
White, both sexes	-	-	2	-	3	4	3	2	2	4	-	-
Male	-	-	1	-	2	2	2	1	-	1	-	-
Female	-	-	1	-	1	2	1	1	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	1	3	2	1	2	2	2	1	-	-
Male	-	1	1	3	1	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	2	1	-	-
White, both sexes	-	1	1	3	2	1	2	2	2	1	-	-
Male	-	1	1	3	1	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
New Hampshire												
All races, both sexes	-	-	1	-	-	2	5	5	-	-	-	-
Male	-	-	-	-	-	1	2	3	-	-	-	-
Female	-	-	1	-	-	1	3	2	-	-	-	-
White, both sexes	-	-	1	-	-	2	5	5	-	-	-	-
Male	-	-	-	-	-	1	2	3	-	-	-	-
Female	-	-	1	-	-	1	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	4	3	3	8	11	29	13	12	7	1	-
Male	1	1	2	3	4	4	15	6	5	1	1	-
Female	-	3	1	-	4	7	14	7	7	6	-	-
White, both sexes	1	2	3	2	7	10	27	12	11	6	1	-
Male	1	1	2	2	4	4	13	5	5	1	1	-
Female	-	1	1	-	3	6	14	7	6	5	-	-
Black, both sexes	-	2	-	-	1	-	2	1	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	2	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
New Mexico												
All races, both sexes	3	1	1	1	4	3	3	4	2	1	-	-
Male	2	-	1	1	4	3	2	2	-	-	-	-
Female	1	1	-	-	-	-	1	2	2	1	-	-
White, both sexes	3	1	-	-	4	3	3	4	2	1	-	-
Male	2	-	-	-	4	3	2	2	-	-	-	-
Female	1	1	-	-	-	-	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	5	7	13	18	28	28	40	22	6	3	-
Male	4	-	5	6	7	15	16	18	5	1	-	-
Female	2	5	2	7	11	13	12	22	17	5	3	-
White, both sexes	4	4	5	8	15	24	21	32	21	6	3	-
Male	3	-	3	5	6	14	14	16	5	1	-	-
Female	1	4	2	3	9	10	7	16	16	5	3	-
Black, both sexes	1	1	2	4	2	4	4	5	1	-	-	-
Male	-	-	2	1	1	1	2	-	-	-	-	-
Female	1	1	-	3	1	3	2	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	3	2	-	-	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
North Carolina												
All races, both sexes	-	4	4	8	12	12	27	12	14	7	1	-
Male	-	2	2	4	8	5	10	4	5	1	-	-
Female	-	2	2	4	4	7	17	8	9	6	1	-
White, both sexes	-	3	3	4	10	11	24	7	12	6	1	-
Male	-	2	2	2	7	5	9	1	5	1	-	-
Female	-	1	1	2	3	6	15	6	7	5	1	-
Black, both sexes	-	1	1	4	2	1	3	5	2	1	-	-
Male	-	-	-	2	1	-	1	3	-	-	-	-
Female	-	1	1	2	1	1	2	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	1	-	3	4	2	1	-	-
Male	-	-	-	-	1	-	3	1	-	-	-	-
Female	-	-	-	1	-	-	-	3	2	1	-	-
White, both sexes	-	-	-	1	1	-	2	4	2	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	-	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Ohio												
All races, both sexes	2	6	4	9	11	13	25	19	18	6	-	-
Male	-	4	1	7	5	10	8	8	5	1	-	-
Female	2	2	3	2	6	3	17	11	13	5	-	-
White, both sexes	1	5	4	7	7	13	22	18	18	5	-	-
Male	-	4	1	5	4	10	8	7	5	1	-	-
Female	1	1	3	2	3	3	14	11	13	4	-	-
Black, both sexes	1	1	-	2	4	-	3	1	-	1	-	-
Male	-	-	-	2	1	-	-	1	-	-	-	-
Female	1	1	-	-	3	-	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	1	3	4	11	11	9	6	2	-	-
Male	-	1	1	3	3	4	6	5	1	-	-	-
Female	1	-	-	-	1	7	5	4	5	2	-	-
White, both sexes	1	1	1	2	3	11	10	6	6	2	-	-
Male	-	1	1	2	3	4	6	3	1	-	-	-
Female	1	-	-	-	-	7	4	3	5	2	-	-
Black, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Oregon												
All races, both sexes	1	-	1	1	3	5	6	15	5	4	-	-
Male	1	-	-	-	2	4	2	2	3	2	-	-
Female	-	-	1	1	1	1	4	13	2	2	-	-
White, both sexes	1	-	1	1	3	5	6	15	5	4	-	-
Male	1	-	-	-	2	4	2	2	3	2	-	-
Female	-	-	1	1	1	1	4	13	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	2	6	7	16	26	28	39	25	6	3	-
Male	1	2	4	5	12	9	13	16	4	-	1	-
Female	-	-	2	2	4	17	15	23	21	6	2	-
White, both sexes	1	2	6	6	16	24	27	38	25	6	3	-
Male	1	2	4	4	12	8	13	16	4	-	1	-
Female	-	-	2	2	4	16	14	22	21	6	2	-
Black, both sexes	-	-	-	1	-	2	1	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Rhode Island												
All races, both sexes	-	-	1	-	2	1	5	2	2	1	-	-
Male	-	-	-	-	2	1	3	-	2	-	-	-
Female	-	-	1	-	-	-	2	2	-	1	-	-
White, both sexes	-	-	1	-	2	1	5	2	2	1	-	-
Male	-	-	-	-	2	1	3	-	2	-	-	-
Female	-	-	1	-	-	-	2	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	-	2	9	4	8	9	12	2	-	-
Male	1	2	-	1	6	3	1	3	4	1	-	-
Female	-	-	-	1	3	1	7	6	8	1	-	-
White, both sexes	-	2	-	1	6	3	7	6	8	2	-	-
Male	-	2	-	1	4	2	1	2	3	1	-	-
Female	-	-	-	-	2	1	6	4	5	1	-	-
Black, both sexes	1	-	-	1	3	1	1	3	4	-	-	-
Male	1	-	-	-	2	1	-	1	1	-	-	-
Female	-	-	-	1	1	-	1	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
South Dakota												
All races, both sexes	3	1	-	-	-	-	4	5	1	-	-	-
Male	2	-	-	-	-	-	4	1	-	-	-	-
Female	1	1	-	-	-	-	-	4	1	-	-	-
White, both sexes	2	1	-	-	-	-	4	5	1	-	-	-
Male	1	-	-	-	-	-	4	1	-	-	-	-
Female	1	1	-	-	-	-	-	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	1	4	2	9	14	14	15	10	3	-	-
Male	3	-	2	1	4	10	6	8	4	1	-	-
Female	2	1	2	1	5	4	8	7	6	2	-	-
White, both sexes	4	1	4	1	8	13	13	15	8	3	-	-
Male	3	-	2	1	3	9	6	8	4	1	-	-
Female	1	1	2	-	5	4	7	7	4	2	-	-
Black, both sexes	1	-	-	1	1	1	1	-	2	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Texas												
All races, both sexes	13	9	7	15	15	36	26	34	34	9	3	-
Male	7	5	1	11	10	18	13	13	9	2	-	-
Female	6	4	6	4	5	18	13	21	25	7	3	-
White, both sexes	7	8	7	12	11	32	25	31	32	7	3	-
Male	4	4	1	9	7	17	13	11	8	1	-	-
Female	3	4	6	3	4	15	12	20	24	6	3	-
Black, both sexes	4	1	-	3	4	4	1	3	2	2	-	-
Male	2	1	-	2	3	1	-	2	1	1	-	-
Female	2	-	-	1	1	3	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	2	-	-	9	5	5	1	-	-
Male	-	-	-	2	-	-	2	1	2	-	-	-
Female	-	-	-	-	-	-	7	4	3	1	-	-
White, both sexes	-	-	-	2	-	-	9	5	5	1	-	-
Male	-	-	-	2	-	-	2	1	2	-	-	-
Female	-	-	-	-	-	-	7	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Vermont												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	2	4	2	6	10	12	19	10	5	-	-
Male	3	1	4	2	5	6	5	8	-	1	-	-
Female	2	1	-	-	1	4	7	11	10	4	-	-
White, both sexes	2	1	3	2	4	8	10	18	8	3	-	-
Male	1	-	3	2	4	5	5	7	-	1	-	-
Female	1	1	-	-	-	3	5	11	8	2	-	-
Black, both sexes	2	-	1	-	-	1	2	1	2	2	-	-
Male	1	-	1	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	-	-	2	-	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	2	1	-	-	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Washington												
All races, both sexes	1	2	2	2	7	7	19	18	10	4	1	-
Male	-	2	2	1	2	4	7	12	2	-	-	-
Female	1	-	-	1	5	3	12	6	8	4	1	-
White, both sexes	1	2	2	2	5	6	18	18	9	4	1	-
Male	-	2	2	1	1	4	7	12	2	-	-	-
Female	1	-	-	1	4	2	11	6	7	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	-	1	-	2	4	5	9	6	4	1	1	-
Male	-	1	-	2	2	3	4	1	1	1	-	-
Female	-	-	-	-	2	2	5	5	3	-	1	-
White, both sexes	-	1	-	2	4	4	9	6	4	1	1	-
Male	-	1	-	2	2	3	4	1	1	1	-	-
Female	-	-	-	-	2	1	5	5	3	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Wisconsin												
All races, both sexes	2	1	2	2	2	8	11	10	7	2	1	-
Male	2	-	-	2	1	3	2	6	3	1	-	-
Female	-	1	2	-	1	5	9	4	4	1	1	-
White, both sexes	1	1	2	2	2	8	11	10	7	2	1	-
Male	1	-	-	2	1	3	2	6	3	1	-	-
Female	-	1	2	-	1	5	9	4	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
United States												
All races, both sexes	4,882	15	6	2	3	8	26	43	66	129	210	249
Male	2,590	10	5	-	1	5	17	33	43	85	148	172
Female	2,292	5	1	2	2	3	9	10	23	44	62	77
White, both sexes	4,106	10	3	2	1	7	16	33	48	94	169	180
Male	2,157	7	2	-	1	5	10	28	31	61	119	122
Female	1,949	3	1	2	-	2	6	5	17	33	50	58
Black, both sexes	639	5	2	-	2	-	10	8	14	31	35	60
Male	355	3	2	-	-	-	7	4	9	22	25	42
Female	284	2	-	-	2	-	3	4	5	9	10	18
American Indian 1/, both sexes	28	-	-	-	-	-	-	1	-	2	2	5
Male	14	-	-	-	-	-	-	1	-	1	2	4
Female	14	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	109	-	1	-	-	1	-	1	4	2	4	4
Male	64	-	1	-	-	-	-	-	3	1	2	4
Female	45	-	-	-	-	1	-	1	1	1	2	-
Alabama												
All races, both sexes	104	-	-	-	-	-	-	1	1	3	5	7
Male	57	-	-	-	-	-	-	1	1	2	4	6
Female	47	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	83	-	-	-	-	-	-	1	1	2	4	5
Male	46	-	-	-	-	-	-	1	1	1	3	4
Female	37	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	21	-	-	-	-	-	-	-	-	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	1	2
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Alaska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	83	-	-	-	-	1	-	2	1	6	4	3
Male	45	-	-	-	-	-	-	2	1	3	3	2
Female	38	-	-	-	-	1	-	-	-	3	1	1
White, both sexes	77	-	-	-	-	1	-	1	1	5	4	2
Male	42	-	-	-	-	-	-	1	1	3	3	1
Female	35	-	-	-	-	1	-	-	-	2	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Arkansas												
All races, both sexes	58	-	-	-	-	-	-	-	1	1	1	5
Male	36	-	-	-	-	-	-	-	1	1	1	5
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	1	1	4
Male	33	-	-	-	-	-	-	-	-	1	1	4
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	468	3	1	-	1	4	3	5	9	18	20	27
Male	261	3	1	-	-	4	1	5	7	10	14	15
Female	207	-	-	-	1	-	2	-	2	8	6	12
White, both sexes	381	2	1	-	-	4	2	4	7	16	17	19
Male	207	2	1	-	-	4	1	4	5	8	12	9
Female	174	-	-	-	-	-	1	-	2	8	5	10
Black, both sexes	44	1	-	-	1	-	1	1	-	2	1	5
Male	26	1	-	-	-	-	-	1	-	2	1	3
Female	18	-	-	-	1	-	1	-	-	-	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	2	-	2	2
Male	26	-	-	-	-	-	-	-	2	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Colorado												
All races, both sexes	51	-	-	-	-	1	-	-	1	-	1	1
Male	28	-	-	-	-	1	-	-	1	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	46	-	-	-	-	1	-	-	1	-	1	-
Male	25	-	-	-	-	1	-	-	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	58	-	-	-	-	-	-	-	-	2	2	2
Male	32	-	-	-	-	-	-	-	-	2	2	2
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	2	2	2
Male	29	-	-	-	-	-	-	-	-	2	2	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Florida												
All races, both sexes	315	2	1	-	-	-	1	3	3	9	8	19
Male	169	1	1	-	-	-	1	-	1	8	5	12
Female	146	1	-	-	-	-	-	3	2	1	3	7
White, both sexes	277	1	-	-	-	-	-	2	2	7	7	14
Male	147	-	-	-	-	-	-	-	1	7	4	9
Female	130	1	-	-	-	-	-	2	1	-	3	5
Black, both sexes	34	1	-	-	-	-	1	1	1	2	1	5
Male	18	1	-	-	-	-	1	-	-	1	1	3
Female	16	-	-	-	-	-	-	1	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	139	1	-	-	-	-	1	1	1	4	5	10
Male	77	1	-	-	-	-	-	1	1	4	4	7
Female	62	-	-	-	-	-	1	-	-	-	1	3
White, both sexes	104	1	-	-	-	-	1	1	1	2	3	6
Male	56	1	-	-	-	-	-	1	1	2	3	5
Female	48	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	33	-	-	-	-	-	-	-	-	2	2	4
Male	19	-	-	-	-	-	-	-	-	2	1	2
Female	14	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Hawaii												
All races, both sexes	23	-	-	-	-	1	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	1	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	1	-	-	-	-	1	-
Idaho												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Illinois												
All races, both sexes	189	1	-	-	-	-	3	4	5	5	14	8
Male	91	1	-	-	-	-	1	2	1	3	8	6
Female	98	-	-	-	-	-	2	2	4	2	6	2
White, both sexes	156	1	-	-	-	-	1	2	5	3	9	5
Male	75	1	-	-	-	-	1	2	1	2	5	3
Female	81	-	-	-	-	-	-	-	4	1	4	2
Black, both sexes	32	-	-	-	-	-	2	1	-	2	5	3
Male	16	-	-	-	-	-	-	-	-	1	3	3
Female	16	-	-	-	-	-	2	1	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	102	-	-	1	1	-	2	1	2	3	8	3
Male	55	-	-	-	1	-	2	1	-	2	7	2
Female	47	-	-	1	-	-	-	-	2	1	1	1
White, both sexes	92	-	-	1	1	-	1	1	2	3	6	3
Male	49	-	-	-	1	-	1	1	-	2	6	2
Female	43	-	-	1	-	-	-	-	2	1	-	1
Black, both sexes	10	-	-	-	-	-	1	-	-	-	2	-
Male	6	-	-	-	-	-	1	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Iowa												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	1	2
Male	22	-	-	-	-	-	-	-	-	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	1	1	2
Male	21	-	-	-	-	-	-	-	-	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	43	-	-	-	-	-	-	-	1	-	-	1
Male	25	-	-	-	-	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	41	-	-	-	-	-	-	-	1	-	-	1
Male	23	-	-	-	-	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Kentucky												
All races, both sexes	80	-	1	-	-	-	-	2	1	-	7	2
Male	46	-	-	-	-	-	-	2	-	-	4	2
Female	34	-	1	-	-	-	-	-	1	-	3	-
White, both sexes	74	-	1	-	-	-	-	2	-	-	7	2
Male	43	-	-	-	-	-	-	2	-	-	4	2
Female	31	-	1	-	-	-	-	-	-	-	3	-
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	81	-	-	-	-	-	-	1	2	5	2	6
Male	43	-	-	-	-	-	-	1	1	1	1	5
Female	38	-	-	-	-	-	-	-	1	4	1	1
White, both sexes	51	-	-	-	-	-	-	1	1	2	1	4
Male	27	-	-	-	-	-	-	1	1	-	-	3
Female	24	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	27	-	-	-	-	-	-	-	1	3	1	2
Male	14	-	-	-	-	-	-	-	-	1	1	2
Female	13	-	-	-	-	-	-	-	1	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	80	-	-	-	-	-	-	1	-	1	2	4
Male	42	-	-	-	-	-	-	1	-	1	1	1
Female	38	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	56	-	-	-	-	-	-	-	-	-	2	1
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	23	-	-	-	-	-	-	1	-	1	-	3
Male	12	-	-	-	-	-	-	1	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Massachusetts												
All races, both sexes	105	-	-	-	-	-	-	-	1	-	2	4
Male	49	-	-	-	-	-	-	-	1	-	1	4
Female	56	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	97	-	-	-	-	-	-	-	1	-	2	4
Male	45	-	-	-	-	-	-	-	1	-	1	4
Female	52	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	173	-	-	-	-	-	-	-	2	5	5	10
Male	87	-	-	-	-	-	-	-	-	1	3	7
Female	86	-	-	-	-	-	-	-	2	4	2	3
White, both sexes	137	-	-	-	-	-	-	-	2	3	4	9
Male	69	-	-	-	-	-	-	-	-	-	2	7
Female	68	-	-	-	-	-	-	-	2	3	2	2
Black, both sexes	34	-	-	-	-	-	-	-	-	2	1	1
Male	17	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Minnesota												
All races, both sexes	89	-	-	-	-	-	-	-	-	3	7	2
Male	45	-	-	-	-	-	-	-	-	2	5	1
Female	44	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	86	-	-	-	-	-	-	-	-	3	6	1
Male	43	-	-	-	-	-	-	-	-	2	5	-
Female	43	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	50	-	-	-	-	-	3	1	4	2	2	4
Male	32	-	-	-	-	-	3	1	2	1	1	4
Female	18	-	-	-	-	-	-	-	2	1	1	-
White, both sexes	29	-	-	-	-	-	2	-	1	-	2	2
Male	18	-	-	-	-	-	2	-	1	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	21	-	-	-	-	-	1	1	3	2	-	2
Male	14	-	-	-	-	-	1	1	1	1	-	2
Female	7	-	-	-	-	-	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Missouri												
All races, both sexes	113	-	-	-	-	1	-	1	-	3	8	5
Male	60	-	-	-	-	-	-	-	-	2	6	4
Female	53	-	-	-	-	1	-	1	-	1	2	1
White, both sexes	102	-	-	-	-	1	-	1	-	2	7	3
Male	53	-	-	-	-	-	-	-	-	1	6	2
Female	49	-	-	-	-	1	-	1	-	1	1	1
Black, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	27	-	-	-	-	-	-	-	-	1	2	3
Male	11	-	-	-	-	-	-	-	-	-	2	3
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	1	2	3
Male	11	-	-	-	-	-	-	-	-	-	2	3
Female	15	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Nebraska												
All races, both sexes	36	-	1	-	-	-	-	1	1	-	2	1
Male	16	-	1	-	-	-	-	-	1	-	1	-
Female	20	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	33	-	-	-	-	-	-	-	1	-	2	1
Male	15	-	-	-	-	-	-	-	1	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	1	-	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	49	-	-	-	-	-	1	-	1	1	4	1
Male	31	-	-	-	-	-	1	-	-	1	3	1
Female	18	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	40	-	-	-	-	-	-	-	-	1	3	-
Male	25	-	-	-	-	-	-	-	-	1	2	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	1	-	-	-	1	-
Male	3	-	-	-	-	-	1	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
New Hampshire												
All races, both sexes	23	-	-	-	-	-	-	-	1	1	-	1
Male	8	-	-	-	-	-	-	-	1	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	1	1	-	1
Male	8	-	-	-	-	-	-	-	1	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	123	1	-	1	-	-	2	-	2	3	3	9
Male	61	1	-	-	-	-	2	-	2	2	1	5
Female	62	-	-	1	-	-	-	-	-	1	2	4
White, both sexes	102	1	-	1	-	-	1	-	1	3	3	8
Male	52	1	-	-	-	-	1	-	1	2	1	5
Female	50	-	-	1	-	-	-	-	-	1	2	3
Black, both sexes	20	-	-	-	-	-	1	-	1	-	-	1
Male	9	-	-	-	-	-	1	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
New Mexico												
All races, both sexes	34	-	-	-	1	-	-	1	1	-	4	2
Male	17	-	-	-	-	-	-	1	1	-	2	2
Female	17	-	-	-	1	-	-	-	-	-	2	-
White, both sexes	29	-	-	-	-	-	-	1	-	-	3	2
Male	15	-	-	-	-	-	-	1	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	290	2	1	-	-	-	1	2	2	11	7	16
Male	140	1	1	-	-	-	1	1	2	6	6	13
Female	150	1	-	-	-	-	-	1	-	5	1	3
White, both sexes	238	1	-	-	-	-	-	2	2	6	6	11
Male	114	1	-	-	-	-	-	1	2	3	5	8
Female	124	-	-	-	-	-	-	1	-	3	1	3
Black, both sexes	48	1	1	-	-	-	1	-	-	4	1	5
Male	24	-	1	-	-	-	1	-	-	3	1	5
Female	24	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
North Carolina												
All races, both sexes	161	1	-	-	-	-	1	2	2	2	4	7
Male	87	1	-	-	-	-	-	2	2	2	3	7
Female	74	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	120	1	-	-	-	-	1	2	1	2	2	5
Male	62	1	-	-	-	-	-	2	1	2	1	5
Female	58	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	37	-	-	-	-	-	-	-	1	-	2	2
Male	23	-	-	-	-	-	-	-	1	-	2	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	1	-	1	1	-
Male	8	-	-	-	-	-	-	1	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	1	-	1	-	-
Male	7	-	-	-	-	-	-	1	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Ohio												
All races, both sexes	246	-	-	-	-	-	2	2	4	6	18	17
Male	124	-	-	-	-	-	1	2	1	4	11	11
Female	122	-	-	-	-	-	1	-	3	2	7	6
White, both sexes	210	-	-	-	-	-	2	2	4	2	14	10
Male	102	-	-	-	-	-	1	2	1	1	8	5
Female	108	-	-	-	-	-	1	-	3	1	6	5
Black, both sexes	32	-	-	-	-	-	-	-	-	2	4	7
Male	21	-	-	-	-	-	-	-	-	2	3	6
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	85	-	-	-	-	-	1	-	-	4	6	2
Male	44	-	-	-	-	-	1	-	-	4	3	-
Female	41	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	75	-	-	-	-	-	1	-	-	3	6	2
Male	40	-	-	-	-	-	1	-	-	3	3	-
Female	35	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Oregon												
All races, both sexes	57	-	-	-	-	-	-	1	3	1	4	5
Male	35	-	-	-	-	-	-	1	3	1	3	1
Female	22	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	55	-	-	-	-	-	-	1	3	1	3	4
Male	34	-	-	-	-	-	-	1	3	1	2	1
Female	21	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	236	-	-	-	-	-	1	1	2	4	4	8
Male	114	-	-	-	-	-	1	1	2	1	3	4
Female	122	-	-	-	-	-	-	-	-	3	1	4
White, both sexes	205	-	-	-	-	-	-	-	1	4	2	6
Male	99	-	-	-	-	-	-	-	1	1	2	4
Female	106	-	-	-	-	-	-	-	-	3	-	2
Black, both sexes	30	-	-	-	-	-	1	1	1	-	2	2
Male	15	-	-	-	-	-	1	1	1	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Rhode Island												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	97	-	-	-	-	-	-	1	2	5	6	3
Male	57	-	-	-	-	-	-	1	2	3	5	2
Female	40	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	62	-	-	-	-	-	-	1	1	2	4	1
Male	35	-	-	-	-	-	-	1	1	1	4	1
Female	27	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	35	-	-	-	-	-	-	-	1	3	2	2
Male	22	-	-	-	-	-	-	-	1	2	1	1
Female	13	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	132	-	-	-	-	-	-	-	5	2	7	7
Male	82	-	-	-	-	-	-	-	3	2	6	3
Female	50	-	-	-	-	-	-	-	2	-	1	4
White, both sexes	112	-	-	-	-	-	-	-	4	2	4	5
Male	66	-	-	-	-	-	-	-	2	2	4	2
Female	46	-	-	-	-	-	-	-	2	-	-	3
Black, both sexes	20	-	-	-	-	-	-	-	1	-	3	2
Male	16	-	-	-	-	-	-	-	1	-	2	1
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Texas												
All races, both sexes	321	1	-	-	-	-	2	6	3	9	16	16
Male	184	-	-	-	-	-	1	5	3	8	14	13
Female	137	1	-	-	-	-	1	1	-	1	2	3
White, both sexes	280	-	-	-	-	-	2	5	1	8	14	14
Male	164	-	-	-	-	-	1	5	1	7	12	12
Female	116	-	-	-	-	-	1	-	-	1	2	2
Black, both sexes	33	1	-	-	-	-	-	1	2	1	1	2
Male	15	-	-	-	-	-	-	-	2	1	1	1
Female	18	1	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	38	-	-	-	-	-	-	1	1	2	-	3
Male	23	-	-	-	-	-	-	1	1	2	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	1	1	1	-	3
Male	21	-	-	-	-	-	-	1	1	1	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	125	-	-	-	-	-	-	-	1	2	3	9
Male	66	-	-	-	-	-	-	-	-	1	2	6
Female	59	-	-	-	-	-	-	-	1	1	1	3
White, both sexes	95	-	-	-	-	-	-	-	1	2	2	5
Male	52	-	-	-	-	-	-	-	-	1	1	4
Female	43	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	28	-	-	-	-	-	-	-	-	-	1	4
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	15	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Washington												
All races, both sexes	76	-	1	-	-	-	-	-	-	2	5	6
Male	41	-	1	-	-	-	-	-	-	1	4	3
Female	35	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	66	-	1	-	-	-	-	-	-	1	4	5
Male	33	-	1	-	-	-	-	-	-	-	3	2
Female	33	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	50	3	-	-	-	-	1	-	-	-	-	1
Male	22	1	-	-	-	-	1	-	-	-	-	1
Female	28	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	2	-	-	-	-	1	-	-	-	-	1
Male	19	-	-	-	-	-	1	-	-	-	-	1
Female	27	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Wisconsin												
All races, both sexes	90	-	-	-	-	-	1	-	-	1	6	1
Male	41	-	-	-	-	-	-	-	-	1	5	1
Female	49	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	89	-	-	-	-	-	1	-	-	1	6	1
Male	41	-	-	-	-	-	-	-	-	1	5	1
Female	48	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
United States												
All races, both sexes	337	321	333	419	413	555	678	584	337	122	26	-
Male	223	225	211	236	225	298	308	224	92	24	5	-
Female	114	96	122	183	188	257	370	360	245	98	21	-
White, both sexes	254	248	270	353	355	487	603	534	308	108	23	-
Male	167	177	172	204	190	262	278	212	85	19	5	-
Female	87	71	98	149	165	225	325	322	223	89	18	-
Black, both sexes	76	62	52	53	42	55	58	39	22	11	2	-
Male	50	41	30	27	24	28	22	9	6	4	-	-
Female	26	21	22	26	18	27	36	30	16	7	2	-
American Indian 1/, both sexes	1	4	3	4	2	1	-	2	1	-	-	-
Male	-	2	3	1	-	-	-	-	-	-	-	-
Female	1	2	-	3	2	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	7	8	9	14	12	17	9	6	3	1	-
Male	6	5	6	4	11	8	8	3	1	1	-	-
Female	-	2	2	5	3	4	9	6	5	2	1	-
Alabama												
All races, both sexes	7	7	8	10	10	13	10	9	8	3	2	-
Male	4	6	5	6	5	7	4	1	3	2	-	-
Female	3	1	3	4	5	6	6	8	5	1	2	-
White, both sexes	6	7	6	7	10	9	9	6	6	2	2	-
Male	3	6	4	5	5	5	4	1	2	1	-	-
Female	3	1	2	2	5	4	5	5	4	1	2	-
Black, both sexes	1	-	2	3	-	4	1	3	2	1	-	-
Male	1	-	1	1	-	2	-	-	1	1	-	-
Female	-	-	1	2	-	2	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Alaska												
All races, both sexes	-	1	1	-	2	1	2	-	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	2	-	1	-	1	-	-	-
White, both sexes	-	-	1	-	2	1	2	-	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	7	8	10	11	6	8	5	4	2	1	-
Male	1	5	3	6	7	4	3	1	3	1	-	-
Female	3	2	5	4	4	2	5	4	1	1	1	-
White, both sexes	4	7	7	9	11	6	8	5	3	2	1	-
Male	1	5	3	6	7	4	3	1	2	1	-	-
Female	3	2	4	3	4	2	5	4	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Arkansas												
All races, both sexes	2	9	2	5	8	6	9	6	3	-	-	-
Male	1	6	1	3	5	4	6	1	1	-	-	-
Female	1	3	1	2	3	2	3	5	2	-	-	-
White, both sexes	2	6	1	5	8	4	8	6	3	-	-	-
Male	1	6	1	3	5	3	6	1	1	-	-	-
Female	1	-	-	2	3	1	2	5	2	-	-	-
Black, both sexes	-	2	1	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	36	27	39	43	41	47	51	56	27	9	1	-
Male	24	21	27	25	22	29	24	20	7	2	-	-
Female	12	6	12	18	19	18	27	36	20	7	1	-
White, both sexes	26	22	29	36	34	37	43	49	24	8	1	-
Male	18	16	20	20	17	24	18	19	7	2	-	-
Female	8	6	9	16	17	13	25	30	17	6	1	-
Black, both sexes	9	4	5	4	2	3	2	3	-	-	-	-
Male	5	4	2	3	1	1	2	-	-	-	-	-
Female	4	-	3	1	1	2	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	3	5	7	6	4	2	1	-	-
Male	1	1	4	2	4	4	4	1	-	-	-	-
Female	-	-	-	1	1	3	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Colorado												
All races, both sexes	6	6	4	3	3	9	9	5	1	1	-	-
Male	3	6	4	1	1	2	4	2	1	1	-	-
Female	3	-	-	2	2	7	5	3	-	-	-	-
White, both sexes	5	6	3	1	3	9	9	5	1	1	-	-
Male	2	6	3	1	1	2	4	2	1	1	-	-
Female	3	-	-	-	2	7	5	3	-	-	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	5	1	6	2	6	8	13	6	2	1	-
Male	2	4	-	3	-	5	5	4	2	-	1	-
Female	-	1	1	3	2	1	3	9	4	2	-	-
White, both sexes	2	3	1	5	2	4	6	12	6	2	1	-
Male	2	3	-	3	-	4	4	4	2	-	1	-
Female	-	-	1	2	2	-	2	8	4	2	-	-
Black, both sexes	-	2	-	1	-	2	2	1	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Delaware												
All races, both sexes	1	2	-	1	-	2	2	2	1	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	1	-	2	1	1	1	-	-	-
White, both sexes	1	2	-	1	-	2	2	2	1	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	1	-	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	2	-	-	2	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	2	-	-	2	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Florida												
All races, both sexes	19	20	23	26	19	38	43	44	28	6	3	-
Male	11	13	14	15	12	23	25	19	6	2	-	-
Female	8	7	9	11	7	15	18	25	22	4	3	-
White, both sexes	16	20	21	22	16	33	40	43	25	6	2	-
Male	9	13	13	13	9	20	24	19	4	2	-	-
Female	7	7	8	9	7	13	16	24	21	4	2	-
Black, both sexes	1	-	2	4	3	5	3	1	2	-	1	-
Male	-	-	1	2	3	3	1	-	1	-	-	-
Female	1	-	1	2	-	2	2	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	20	14	9	16	7	10	26	9	4	1	-	-
Male	12	10	5	11	3	5	9	2	2	-	-	-
Female	8	4	4	5	4	5	17	7	2	1	-	-
White, both sexes	13	8	6	16	4	8	21	8	4	1	-	-
Male	8	4	4	11	1	3	8	2	2	-	-	-
Female	5	4	2	5	3	5	13	6	2	1	-	-
Black, both sexes	7	6	3	-	2	1	5	1	-	-	-	-
Male	4	6	1	-	1	1	1	-	-	-	-	-
Female	3	-	2	-	1	-	4	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Hawaii												
All races, both sexes	-	-	1	2	3	1	5	3	2	2	-	-
Male	-	-	1	1	2	1	2	1	-	2	-	-
Female	-	-	-	1	1	-	3	2	2	-	-	-
White, both sexes	-	-	-	1	1	1	1	-	-	1	-	-
Male	-	-	-	1	-	1	-	-	-	1	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	-	4	3	2	1	-	-
Male	-	-	1	-	2	-	2	1	-	1	-	-
Female	-	-	-	1	-	-	2	2	2	-	-	-
Idaho												
All races, both sexes	1	3	-	5	2	2	3	4	-	-	-	-
Male	1	3	-	4	-	1	2	3	-	-	-	-
Female	-	-	-	1	2	1	1	1	-	-	-	-
White, both sexes	1	3	-	5	2	2	3	4	-	-	-	-
Male	1	3	-	4	-	1	2	3	-	-	-	-
Female	-	-	-	1	2	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Illinois												
All races, both sexes	3	15	14	9	22	13	28	25	13	4	3	-
Male	2	10	10	5	11	8	11	9	3	-	-	-
Female	1	5	4	4	11	5	17	16	10	4	3	-
White, both sexes	2	9	12	9	19	11	28	22	11	4	3	-
Male	1	6	9	5	9	7	11	9	3	-	-	-
Female	1	3	3	4	10	4	17	13	8	4	3	-
Black, both sexes	1	6	2	-	3	2	-	3	2	-	-	-
Male	1	4	1	-	2	1	-	-	-	-	-	-
Female	-	2	1	-	1	1	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	9	6	8	8	5	5	14	12	9	3	2	-
Male	6	5	4	4	5	3	7	5	-	-	1	-
Female	3	1	4	4	-	2	7	7	9	3	1	-
White, both sexes	6	5	7	8	5	5	14	11	8	3	2	-
Male	4	4	3	4	5	3	7	5	-	-	1	-
Female	2	1	4	4	-	2	7	6	8	3	1	-
Black, both sexes	3	1	1	-	-	-	-	1	1	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Iowa												
All races, both sexes	3	1	2	2	4	6	9	9	4	-	-	-
Male	2	1	1	1	3	5	2	4	-	-	-	-
Female	1	-	1	1	1	1	7	5	4	-	-	-
White, both sexes	3	1	2	2	4	5	9	9	4	-	-	-
Male	2	1	1	1	3	4	2	4	-	-	-	-
Female	1	-	1	1	1	1	7	5	4	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	5	3	2	1	7	6	9	5	1	-	-
Male	1	3	2	1	-	4	6	6	1	-	-	-
Female	1	2	1	1	1	3	-	3	4	1	-	-
White, both sexes	2	5	3	1	1	7	5	9	5	1	-	-
Male	1	3	2	-	-	4	5	6	1	-	-	-
Female	1	2	1	1	1	3	-	3	4	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Kentucky												
All races, both sexes	6	4	9	10	4	7	12	10	2	2	1	-
Male	5	3	6	5	3	3	4	8	-	-	1	-
Female	1	1	3	5	1	4	8	2	2	2	-	-
White, both sexes	5	4	8	9	4	7	11	9	2	2	1	-
Male	4	3	5	5	3	3	4	7	-	-	1	-
Female	1	1	3	4	1	4	7	2	2	2	-	-
Black, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	9	9	7	5	7	9	10	1	2	-	-
Male	4	7	6	3	3	1	4	5	-	1	-	-
Female	2	2	3	4	2	6	5	5	1	1	-	-
White, both sexes	3	3	7	3	3	5	6	10	1	1	-	-
Male	3	2	5	1	3	1	2	5	-	-	-	-
Female	-	1	2	2	-	4	4	5	1	1	-	-
Black, both sexes	3	4	2	4	2	2	2	-	-	1	-	-
Male	1	3	1	2	-	-	2	-	-	1	-	-
Female	2	1	1	2	2	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Maine												
All races, both sexes	1	1	1	4	1	2	1	5	2	2	-	-
Male	1	-	-	2	1	1	-	4	1	1	-	-
Female	-	1	1	2	-	1	1	1	1	1	-	-
White, both sexes	1	1	1	4	1	2	1	5	2	2	-	-
Male	1	-	-	2	1	1	-	4	1	1	-	-
Female	-	1	1	2	-	1	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	8	8	8	6	7	14	11	6	-	-	-
Male	3	6	4	4	4	5	8	2	2	-	-	-
Female	1	2	4	4	2	2	6	9	4	-	-	-
White, both sexes	3	5	4	5	4	6	11	10	5	-	-	-
Male	2	4	1	2	3	5	7	2	2	-	-	-
Female	1	1	3	3	1	1	4	8	3	-	-	-
Black, both sexes	-	3	4	3	2	1	3	1	1	-	-	-
Male	-	2	3	2	1	-	1	-	-	-	-	-
Female	-	1	1	1	1	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Massachusetts												
All races, both sexes	9	7	10	7	12	12	13	7	13	6	2	-
Male	6	4	5	5	7	6	5	2	3	-	-	-
Female	3	3	5	2	5	6	8	5	10	6	2	-
White, both sexes	9	4	10	7	12	10	12	7	12	5	2	-
Male	6	2	5	5	7	4	5	2	3	-	-	-
Female	3	2	5	2	5	6	7	5	9	5	2	-
Black, both sexes	-	3	-	-	-	1	1	-	1	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Michigan												
All races, both sexes	11	14	15	16	14	23	25	17	11	5	-	-
Male	9	7	11	9	8	12	8	10	1	1	-	-
Female	2	7	4	7	6	11	17	7	10	4	-	-
White, both sexes	8	7	10	14	12	20	21	13	10	4	-	-
Male	7	3	7	8	7	10	7	9	1	1	-	-
Female	1	4	3	6	5	10	14	4	9	3	-	-
Black, both sexes	3	6	5	2	2	3	3	4	1	1	-	-
Male	2	3	4	1	1	2	1	1	-	-	-	-
Female	1	3	1	1	1	1	2	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Minnesota												
All races, both sexes	5	12	5	4	4	8	9	19	6	5	-	-
Male	3	8	4	2	4	6	3	5	2	-	-	-
Female	2	4	1	2	-	2	6	14	4	5	-	-
White, both sexes	5	12	5	4	3	8	9	19	6	5	-	-
Male	3	8	4	2	3	6	3	5	2	-	-	-
Female	2	4	1	2	-	2	6	14	4	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	2	2	4	3	6	1	3	4	3	-	-
Male	6	2	1	3	1	2	1	1	2	1	-	-
Female	-	-	1	1	2	4	-	2	2	2	-	-
White, both sexes	2	2	1	2	3	2	1	3	4	2	-	-
Male	2	2	1	1	1	-	1	1	2	1	-	-
Female	-	-	-	1	2	2	-	2	2	1	-	-
Black, both sexes	4	-	1	2	-	4	-	-	-	1	-	-
Male	4	-	-	2	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	2	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Missouri												
All races, both sexes	8	3	4	11	11	12	13	22	9	2	-	-
Male	7	2	1	6	5	8	9	7	2	1	-	-
Female	1	1	3	5	6	4	4	15	7	1	-	-
White, both sexes	6	2	3	10	10	11	13	22	9	2	-	-
Male	6	1	1	5	5	7	9	7	2	1	-	-
Female	-	1	2	5	5	4	4	15	7	1	-	-
Black, both sexes	2	1	1	1	1	1	-	-	-	-	-	-
Male	1	1	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	1	-	2	4	-	7	3	2	-	-	-
Male	1	-	-	-	2	-	1	1	1	-	-	-
Female	1	1	-	2	2	-	6	2	1	-	-	-
White, both sexes	2	1	-	2	4	-	7	2	2	-	-	-
Male	1	-	-	-	2	-	1	1	1	-	-	-
Female	1	1	-	2	2	-	6	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Nebraska												
All races, both sexes	2	3	-	1	3	5	6	2	4	4	-	-
Male	1	3	-	-	2	2	1	1	2	1	-	-
Female	1	-	-	1	1	3	5	1	2	3	-	-
White, both sexes	2	3	-	1	3	5	5	2	4	4	-	-
Male	1	3	-	-	2	2	1	1	2	1	-	-
Female	1	-	-	1	1	3	4	1	2	3	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	4	4	8	4	8	4	5	-	-	-	-
Male	3	3	4	6	3	3	2	1	-	-	-	-
Female	1	1	-	2	1	5	2	4	-	-	-	-
White, both sexes	3	4	3	7	4	7	3	5	-	-	-	-
Male	2	3	3	5	3	3	2	1	-	-	-	-
Female	1	1	-	2	1	4	1	4	-	-	-	-
Black, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
New Hampshire												
All races, both sexes	1	-	1	1	1	6	2	5	1	2	-	-
Male	1	-	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	5	1	5	1	2	-	-
White, both sexes	1	-	1	1	1	6	2	5	1	2	-	-
Male	1	-	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	5	1	5	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	13	6	4	6	8	19	19	13	9	4	1	-
Male	9	5	3	2	4	9	9	3	3	-	1	-
Female	4	1	1	4	4	10	10	10	6	4	-	-
White, both sexes	10	5	3	3	6	17	16	12	7	4	1	-
Male	6	4	2	2	3	9	8	3	3	-	1	-
Female	4	1	1	1	3	8	8	9	4	4	-	-
Black, both sexes	3	1	1	3	1	2	3	1	2	-	-	-
Male	3	1	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	3	-	2	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
New Mexico												
All races, both sexes	-	4	-	3	1	3	10	2	-	2	-	-
Male	-	3	-	1	-	1	5	1	-	-	-	-
Female	-	1	-	2	1	2	5	1	-	2	-	-
White, both sexes	-	4	-	2	-	3	10	2	-	2	-	-
Male	-	3	-	1	-	1	5	1	-	-	-	-
Female	-	1	-	1	-	2	5	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	15	12	21	28	29	41	39	26	22	13	2	-
Male	10	7	12	11	15	21	16	8	7	2	-	-
Female	5	5	9	17	14	20	23	18	15	11	2	-
White, both sexes	7	8	19	22	27	36	31	24	22	12	2	-
Male	5	5	12	9	14	18	14	8	7	2	-	-
Female	2	3	7	13	13	18	17	16	15	10	2	-
Black, both sexes	8	4	2	6	1	4	8	1	-	1	-	-
Male	5	2	-	2	-	2	2	-	-	-	-	-
Female	3	2	2	4	1	2	6	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
North Carolina												
All races, both sexes	17	13	15	21	11	23	16	13	10	2	1	-
Male	13	9	9	13	2	11	8	2	2	1	-	-
Female	4	4	6	8	9	12	8	11	8	1	1	-
White, both sexes	9	10	10	19	7	18	12	12	7	2	-	-
Male	7	8	6	12	-	8	5	2	1	1	-	-
Female	2	2	4	7	7	10	7	10	6	1	-	-
Black, both sexes	8	3	3	2	3	4	4	1	3	-	1	-
Male	6	1	2	1	2	2	3	-	1	-	-	-
Female	2	2	1	1	1	2	1	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	-	-	1	-	1	2	-	2	1	-
Male	1	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	1	1	-
White, both sexes	1	1	-	-	1	-	1	2	-	2	1	-
Male	1	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Ohio												
All races, both sexes	17	10	12	26	22	23	39	34	12	2	-	-
Male	14	5	7	11	12	10	15	17	2	1	-	-
Female	3	5	5	15	10	13	24	17	10	1	-	-
White, both sexes	14	9	10	24	21	20	34	30	12	2	-	-
Male	11	5	7	11	11	8	14	14	2	1	-	-
Female	3	4	3	13	10	12	20	16	10	1	-	-
Black, both sexes	3	1	2	2	1	2	4	4	-	-	-	-
Male	3	-	-	-	1	2	1	3	-	-	-	-
Female	-	1	2	2	-	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Oklahoma												
All races, both sexes	7	5	7	7	3	9	10	16	6	2	-	-
Male	3	4	3	3	2	6	6	8	1	-	-	-
Female	4	1	4	4	1	3	4	8	5	2	-	-
White, both sexes	6	3	6	5	3	9	9	15	5	2	-	-
Male	3	3	2	2	2	6	6	8	1	-	-	-
Female	3	-	4	3	1	3	3	7	4	2	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	2	-	-	-	1	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Oregon												
All races, both sexes	4	4	6	3	3	4	9	7	2	1	-	-
Male	4	3	4	2	2	1	5	5	-	-	-	-
Female	-	1	2	1	1	3	4	2	2	1	-	-
White, both sexes	4	4	6	3	3	4	9	7	2	1	-	-
Male	4	3	4	2	2	1	5	5	-	-	-	-
Female	-	1	2	1	1	3	4	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	14	3	14	9	20	31	45	36	28	15	1	-
Male	10	3	10	8	13	16	18	14	7	2	1	-
Female	4	-	4	1	7	15	27	22	21	13	-	-
White, both sexes	10	2	11	8	17	28	43	33	27	12	1	-
Male	7	2	8	7	11	16	18	14	6	1	1	-
Female	3	-	3	1	6	12	25	19	21	11	-	-
Black, both sexes	4	1	3	1	3	3	1	3	1	3	-	-
Male	3	1	2	1	2	-	-	-	1	1	-	-
Female	1	-	1	-	1	3	1	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Rhode Island												
All races, both sexes	2	1	-	1	2	5	7	4	3	-	-	-
Male	1	1	-	1	1	2	2	1	-	-	-	-
Female	1	-	-	-	1	3	5	3	3	-	-	-
White, both sexes	2	1	-	1	2	5	7	4	3	-	-	-
Male	1	1	-	1	1	2	2	1	-	-	-	-
Female	1	-	-	-	1	3	5	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	11	9	7	7	13	9	8	2	2	-	-
Male	7	6	5	7	5	7	4	3	-	-	-	-
Female	5	5	4	-	2	6	5	5	2	2	-	-
White, both sexes	4	6	6	4	6	9	8	7	2	1	-	-
Male	3	3	3	4	4	4	4	2	-	-	-	-
Female	1	3	3	-	2	5	4	5	2	1	-	-
Black, both sexes	8	5	3	3	1	4	1	1	-	1	-	-
Male	4	3	2	3	1	3	-	1	-	-	-	-
Female	4	2	1	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
South Dakota												
All races, both sexes	1	1	-	-	2	1	3	2	-	-	-	-
Male	1	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-
White, both sexes	1	1	-	-	1	1	3	2	-	-	-	-
Male	1	1	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	9	12	12	10	23	15	10	9	1	-	-
Male	4	7	9	7	5	16	9	5	6	-	-	-
Female	6	2	3	5	5	7	6	5	3	1	-	-
White, both sexes	8	7	10	10	8	23	12	10	8	1	-	-
Male	3	5	7	6	3	16	6	5	5	-	-	-
Female	5	2	3	4	5	7	6	5	3	1	-	-
Black, both sexes	2	2	2	2	2	-	3	-	1	-	-	-
Male	1	2	2	1	2	-	3	-	1	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Texas												
All races, both sexes	22	22	18	32	35	36	44	31	18	7	3	-
Male	14	16	15	19	17	20	21	13	5	-	-	-
Female	8	6	3	13	18	16	23	18	13	7	3	-
White, both sexes	18	19	16	27	30	36	41	25	16	6	2	-
Male	12	14	13	17	14	20	20	11	5	-	-	-
Female	6	5	3	10	16	16	21	14	11	6	2	-
Black, both sexes	4	2	2	2	4	-	3	5	2	1	-	-
Male	2	1	2	1	2	-	1	1	-	-	-	-
Female	2	1	-	1	2	-	2	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	1	-	-	1	-	-	1	-
Male	-	1	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	2	1	6	2	4	1	8	3	4	-	-	-
Male	1	1	2	1	2	1	6	1	1	-	-	-
Female	1	-	4	1	2	-	2	2	3	-	-	-
White, both sexes	2	1	6	2	3	1	8	3	4	-	-	-
Male	1	1	2	1	1	1	6	1	1	-	-	-
Female	1	-	4	1	2	-	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Vermont												
All races, both sexes	2	-	1	-	1	1	2	2	2	-	-	-
Male	1	-	1	-	1	-	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	2	1	-	-	-
White, both sexes	2	-	1	-	1	1	2	2	2	-	-	-
Male	1	-	1	-	1	-	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	7	7	12	18	16	22	16	7	-	-	-
Male	3	5	4	9	10	6	11	6	3	-	-	-
Female	2	2	3	3	8	10	11	10	4	-	-	-
White, both sexes	5	4	5	8	11	14	17	16	5	-	-	-
Male	3	3	3	7	7	5	9	6	3	-	-	-
Female	2	1	2	1	4	9	8	10	2	-	-	-
Black, both sexes	-	3	2	4	7	2	3	-	2	-	-	-
Male	-	2	1	2	3	1	1	-	-	-	-	-
Female	-	1	1	2	4	1	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Washington												
All races, both sexes	4	9	3	4	9	12	8	6	6	1	-	-
Male	3	4	1	2	6	6	4	2	4	-	-	-
Female	1	5	2	2	3	6	4	4	2	1	-	-
White, both sexes	3	5	3	4	9	11	7	6	6	1	-	-
Male	2	2	1	2	6	5	3	2	4	-	-	-
Female	1	3	2	2	3	6	4	4	2	1	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	2	2	4	3	13	5	5	8	-	-	-
Male	1	2	2	2	1	8	-	1	2	-	-	-
Female	2	-	-	2	2	5	5	4	6	-	-	-
White, both sexes	3	2	2	3	3	13	5	3	8	-	-	-
Male	1	2	2	1	1	8	-	-	2	-	-	-
Female	2	-	-	2	2	5	5	3	6	-	-	-
Black, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Wisconsin												
All races, both sexes	6	4	3	10	8	6	17	16	10	-	1	-
Male	3	2	1	5	5	4	7	5	2	-	-	-
Female	3	2	2	5	3	2	10	11	8	-	1	-
White, both sexes	6	4	3	10	8	5	17	16	10	-	1	-
Male	3	2	1	5	5	4	7	5	2	-	-	-
Female	3	2	2	5	3	1	10	11	8	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	1	2	-	1	-	2	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
White, both sexes	-	-	1	1	2	-	1	-	2	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
United States												
All races, both sexes	3,373	-	2	1	2	7	23	38	60	111	186	214
Male	1,885	-	2	-	-	5	15	29	40	75	133	150
Female	1,488	-	-	1	2	2	8	9	20	36	53	64
White, both sexes	2,815	-	-	1	-	6	15	29	44	81	152	154
Male	1,560	-	-	-	-	5	9	24	28	53	109	106
Female	1,255	-	-	1	-	1	6	5	16	28	43	48
Black, both sexes	472	-	1	-	2	-	8	7	12	27	30	53
Male	276	-	1	-	-	-	6	4	9	20	22	38
Female	196	-	-	-	2	-	2	3	3	7	8	15
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	-	2	-	4
Male	9	-	-	-	-	-	-	1	-	1	-	3
Female	11	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	66	-	1	-	-	1	-	1	4	1	4	3
Male	40	-	1	-	-	-	-	-	3	1	2	3
Female	26	-	-	-	-	1	-	1	1	-	2	-
Alabama												
All races, both sexes	84	-	-	-	-	-	-	1	1	3	5	6
Male	51	-	-	-	-	-	-	1	1	2	4	5
Female	33	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	66	-	-	-	-	-	-	1	1	2	4	4
Male	40	-	-	-	-	-	-	1	1	1	3	3
Female	26	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	18	-	-	-	-	-	-	-	-	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	64	-	-	-	-	1	-	2	1	4	3	3
Male	37	-	-	-	-	-	-	2	1	2	2	2
Female	27	-	-	-	-	1	-	-	-	2	1	1
White, both sexes	61	-	-	-	-	1	-	1	1	4	3	2
Male	35	-	-	-	-	-	-	1	1	2	2	1
Female	26	-	-	-	-	1	-	-	-	2	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Arkansas												
All races, both sexes	46	-	-	-	-	-	-	-	1	1	1	5
Male	33	-	-	-	-	-	-	-	1	1	1	5
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	1	1	4
Male	30	-	-	-	-	-	-	-	-	1	1	4
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	313	-	-	-	1	4	3	5	9	16	17	25
Male	182	-	-	-	-	4	1	5	7	9	13	14
Female	131	-	-	-	1	-	2	-	2	7	4	11
White, both sexes	251	-	-	-	-	4	2	4	7	14	14	17
Male	142	-	-	-	-	4	1	4	5	7	11	8
Female	109	-	-	-	-	-	1	-	2	7	3	9
Black, both sexes	37	-	-	-	1	-	1	1	-	2	1	5
Male	23	-	-	-	-	-	-	1	-	2	1	3
Female	14	-	-	-	1	-	1	-	-	-	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	2	-	2	2
Male	15	-	-	-	-	-	-	-	2	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Colorado												
All races, both sexes	30	-	-	-	-	1	-	-	1	-	1	1
Male	17	-	-	-	-	1	-	-	1	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	26	-	-	-	-	1	-	-	1	-	1	-
Male	15	-	-	-	-	1	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	41	-	-	-	-	-	-	-	-	2	2	2
Male	25	-	-	-	-	-	-	-	-	2	2	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	2	2	2
Male	24	-	-	-	-	-	-	-	-	2	2	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Delaware												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Florida												
All races, both sexes	218	-	1	-	-	-	1	2	3	8	8	17
Male	122	-	1	-	-	-	1	-	1	7	5	11
Female	96	-	-	-	-	-	-	2	2	1	3	6
White, both sexes	190	-	-	-	-	-	-	2	2	7	7	13
Male	105	-	-	-	-	-	-	-	1	7	4	8
Female	85	-	-	-	-	-	-	2	1	-	3	5
Black, both sexes	25	-	-	-	-	-	1	-	1	1	1	4
Male	14	-	-	-	-	-	1	-	-	-	1	3
Female	11	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	112	-	-	-	-	-	1	1	1	4	5	9
Male	65	-	-	-	-	-	-	1	1	4	4	6
Female	47	-	-	-	-	-	1	-	-	-	1	3
White, both sexes	79	-	-	-	-	-	1	1	1	2	3	5
Male	45	-	-	-	-	-	-	1	1	2	3	4
Female	34	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	31	-	-	-	-	-	-	-	-	2	2	4
Male	18	-	-	-	-	-	-	-	-	2	1	2
Female	13	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Hawaii												
All races, both sexes	13	-	-	-	-	1	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	1	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	1	-	-	-	-	1	-
Idaho												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Illinois												
All races, both sexes	136	-	-	-	-	-	2	4	4	5	12	6
Male	66	-	-	-	-	-	1	2	1	3	6	4
Female	70	-	-	-	-	-	1	2	3	2	6	2
White, both sexes	111	-	-	-	-	-	1	2	4	3	8	4
Male	55	-	-	-	-	-	1	2	1	2	4	2
Female	56	-	-	-	-	-	-	-	3	1	4	2
Black, both sexes	24	-	-	-	-	-	1	1	-	2	4	2
Male	11	-	-	-	-	-	-	-	-	1	2	2
Female	13	-	-	-	-	-	1	1	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	76	-	-	-	-	-	2	1	2	2	7	3
Male	42	-	-	-	-	-	2	1	-	2	6	2
Female	34	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	68	-	-	-	-	-	1	1	2	2	5	3
Male	37	-	-	-	-	-	1	1	-	2	5	2
Female	31	-	-	-	-	-	-	-	2	-	-	1
Black, both sexes	8	-	-	-	-	-	1	-	-	-	2	-
Male	5	-	-	-	-	-	1	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Iowa												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Kentucky												
All races, both sexes	62	-	-	-	-	-	-	2	-	-	7	2
Male	38	-	-	-	-	-	-	2	-	-	4	2
Female	24	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	59	-	-	-	-	-	-	2	-	-	7	2
Male	36	-	-	-	-	-	-	2	-	-	4	2
Female	23	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	57	-	-	-	-	-	-	-	2	4	2	6
Male	35	-	-	-	-	-	-	-	1	1	1	5
Female	22	-	-	-	-	-	-	-	1	3	1	1
White, both sexes	35	-	-	-	-	-	-	-	1	2	1	4
Male	22	-	-	-	-	-	-	-	1	-	-	3
Female	13	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	20	-	-	-	-	-	-	-	1	2	1	2
Male	11	-	-	-	-	-	-	-	-	1	1	2
Female	9	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Maine												
All races, both sexes	15	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	53	-	-	-	-	-	-	1	-	1	2	3
Male	31	-	-	-	-	-	-	1	-	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	37	-	-	-	-	-	-	-	-	-	2	1
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	15	-	-	-	-	-	-	1	-	1	-	2
Male	8	-	-	-	-	-	-	1	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Massachusetts												
All races, both sexes	71	-	-	-	-	-	-	-	1	-	1	4
Male	37	-	-	-	-	-	-	-	1	-	-	4
Female	34	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	67	-	-	-	-	-	-	-	1	-	1	4
Male	35	-	-	-	-	-	-	-	1	-	-	4
Female	32	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	109	-	-	-	-	-	-	-	2	5	5	7
Male	55	-	-	-	-	-	-	-	-	1	3	5
Female	54	-	-	-	-	-	-	-	2	4	2	2
White, both sexes	87	-	-	-	-	-	-	-	2	3	4	6
Male	44	-	-	-	-	-	-	-	-	-	2	5
Female	43	-	-	-	-	-	-	-	2	3	2	1
Black, both sexes	20	-	-	-	-	-	-	-	-	2	1	1
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Minnesota												
All races, both sexes	56	-	-	-	-	-	-	-	-	2	6	1
Male	29	-	-	-	-	-	-	-	-	1	4	1
Female	27	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	54	-	-	-	-	-	-	-	-	2	5	-
Male	28	-	-	-	-	-	-	-	-	1	4	-
Female	26	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	37	-	-	-	-	-	3	1	3	2	2	3
Male	23	-	-	-	-	-	3	1	2	1	1	3
Female	14	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	23	-	-	-	-	-	2	-	1	-	2	1
Male	12	-	-	-	-	-	2	-	1	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	1	1	2	2	-	2
Male	11	-	-	-	-	-	1	1	1	1	-	2
Female	3	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Missouri												
All races, both sexes	81	-	-	-	-	-	-	1	-	2	8	2
Male	42	-	-	-	-	-	-	-	-	2	6	1
Female	39	-	-	-	-	-	-	1	-	-	2	1
White, both sexes	74	-	-	-	-	-	-	1	-	1	7	2
Male	39	-	-	-	-	-	-	-	-	1	6	1
Female	35	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	2	3
Male	9	-	-	-	-	-	-	-	-	-	2	3
Female	13	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	2	3
Male	9	-	-	-	-	-	-	-	-	-	2	3
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Nebraska												
All races, both sexes	26	-	1	-	-	-	-	1	1	-	1	1
Male	14	-	1	-	-	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	1	-	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	1	-	1	1	4	1
Male	23	-	-	-	-	-	1	-	-	1	3	1
Female	14	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	30	-	-	-	-	-	-	-	-	1	3	-
Male	18	-	-	-	-	-	-	-	-	1	2	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	1	-	-	-	1	-
Male	3	-	-	-	-	-	1	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	1	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	1	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	84	-	-	1	-	-	2	-	1	2	3	8
Male	49	-	-	-	-	-	2	-	1	2	1	4
Female	35	-	-	1	-	-	-	-	-	-	2	4
White, both sexes	68	-	-	1	-	-	1	-	-	2	3	7
Male	41	-	-	-	-	-	1	-	-	2	1	4
Female	27	-	-	1	-	-	-	-	-	-	2	3
Black, both sexes	15	-	-	-	-	-	1	-	1	-	-	1
Male	8	-	-	-	-	-	1	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
New Mexico												
All races, both sexes	18	-	-	-	1	-	-	1	1	-	2	2
Male	10	-	-	-	-	-	-	1	1	-	1	2
Female	8	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	1	-	-	2	2
Male	9	-	-	-	-	-	-	1	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	180	-	-	-	-	-	-	2	1	10	6	13
Male	92	-	-	-	-	-	-	1	1	6	5	12
Female	88	-	-	-	-	-	-	1	-	4	1	1
White, both sexes	143	-	-	-	-	-	-	2	1	6	5	9
Male	72	-	-	-	-	-	-	1	1	3	4	8
Female	71	-	-	-	-	-	-	1	-	3	1	1
Black, both sexes	35	-	-	-	-	-	-	-	-	4	1	4
Male	19	-	-	-	-	-	-	-	-	3	1	4
Female	16	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
North Carolina												
All races, both sexes	115	-	-	-	-	-	1	2	2	2	3	7
Male	67	-	-	-	-	-	-	2	2	2	2	7
Female	48	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	87	-	-	-	-	-	1	2	1	2	2	5
Male	47	-	-	-	-	-	-	2	1	2	1	5
Female	40	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	26	-	-	-	-	-	-	-	1	-	1	2
Male	19	-	-	-	-	-	-	-	1	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Ohio												
All races, both sexes	173	-	-	-	-	-	2	1	4	6	18	14
Male	90	-	-	-	-	-	1	1	1	4	11	9
Female	83	-	-	-	-	-	1	-	3	2	7	5
White, both sexes	147	-	-	-	-	-	2	1	4	2	14	8
Male	71	-	-	-	-	-	1	1	1	1	8	4
Female	76	-	-	-	-	-	1	-	3	1	6	4
Black, both sexes	24	-	-	-	-	-	-	-	-	2	4	6
Male	18	-	-	-	-	-	-	-	-	2	3	5
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	69	-	-	-	-	-	1	-	-	4	6	2
Male	35	-	-	-	-	-	1	-	-	4	3	-
Female	34	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	61	-	-	-	-	-	1	-	-	3	6	2
Male	33	-	-	-	-	-	1	-	-	3	3	-
Female	28	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Oregon												
All races, both sexes	35	-	-	-	-	-	-	1	3	1	3	4
Male	23	-	-	-	-	-	-	1	3	1	3	-
Female	12	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	33	-	-	-	-	-	-	1	3	1	2	3
Male	22	-	-	-	-	-	-	1	3	1	2	-
Female	11	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	150	-	-	-	-	-	1	1	2	3	3	6
Male	82	-	-	-	-	-	1	1	2	-	3	3
Female	68	-	-	-	-	-	-	-	-	3	-	3
White, both sexes	129	-	-	-	-	-	-	-	1	3	2	4
Male	70	-	-	-	-	-	-	-	1	-	2	3
Female	59	-	-	-	-	-	-	-	-	3	-	1
Black, both sexes	21	-	-	-	-	-	1	1	1	-	1	2
Male	12	-	-	-	-	-	1	1	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	80	-	-	-	-	-	-	1	2	5	6	2
Male	49	-	-	-	-	-	-	1	2	3	5	1
Female	31	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	48	-	-	-	-	-	-	1	1	2	4	-
Male	29	-	-	-	-	-	-	1	1	1	4	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	32	-	-	-	-	-	-	-	1	3	2	2
Male	20	-	-	-	-	-	-	-	1	2	1	1
Female	12	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	98	-	-	-	-	-	-	-	5	2	6	6
Male	59	-	-	-	-	-	-	-	3	2	6	2
Female	39	-	-	-	-	-	-	-	2	-	-	4
White, both sexes	83	-	-	-	-	-	-	-	4	2	4	4
Male	47	-	-	-	-	-	-	-	2	2	4	1
Female	36	-	-	-	-	-	-	-	2	-	-	3
Black, both sexes	15	-	-	-	-	-	-	-	1	-	2	2
Male	12	-	-	-	-	-	-	-	1	-	2	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Texas												
All races, both sexes	219	-	-	-	-	-	2	4	3	6	15	14
Male	137	-	-	-	-	-	1	3	3	5	14	13
Female	82	-	-	-	-	-	1	1	-	1	1	1
White, both sexes	195	-	-	-	-	-	2	3	1	5	13	13
Male	123	-	-	-	-	-	1	3	1	4	12	12
Female	72	-	-	-	-	-	1	-	-	1	1	1
Black, both sexes	20	-	-	-	-	-	-	1	2	1	1	1
Male	12	-	-	-	-	-	-	-	2	1	1	1
Female	8	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	26	-	-	-	-	-	-	1	1	1	-	3
Male	17	-	-	-	-	-	-	1	1	1	-	3
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	1	1	1	-	3
Male	16	-	-	-	-	-	-	1	1	1	-	3
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Vermont												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	85	-	-	-	-	-	-	-	1	1	2	8
Male	40	-	-	-	-	-	-	-	-	1	1	6
Female	45	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	66	-	-	-	-	-	-	-	1	1	2	4
Male	34	-	-	-	-	-	-	-	-	1	1	4
Female	32	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	4
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Washington												
All races, both sexes	51	-	-	-	-	-	-	-	-	2	2	4
Male	30	-	-	-	-	-	-	-	-	1	2	2
Female	21	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	42	-	-	-	-	-	-	-	-	1	1	3
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Wisconsin												
All races, both sexes	63	-	-	-	-	-	1	-	-	1	6	1
Male	31	-	-	-	-	-	-	-	-	1	5	1
Female	32	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	63	-	-	-	-	-	1	-	-	1	6	1
Male	31	-	-	-	-	-	-	-	-	1	5	1
Female	32	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
United States												
All races, both sexes	293	262	236	305	294	361	421	317	164	63	13	-
Male	200	181	151	170	168	194	191	120	45	13	3	-
Female	93	81	85	135	126	167	230	197	119	50	10	-
White, both sexes	223	204	195	259	249	316	381	289	151	54	12	-
Male	151	143	126	148	140	170	176	116	43	10	3	-
Female	72	61	69	111	109	146	205	173	108	44	9	-
Black, both sexes	64	49	36	38	34	38	34	21	11	7	-	-
Male	44	33	20	20	20	19	12	3	2	3	-	-
Female	20	16	16	18	14	19	22	18	9	4	-	-
American Indian 1/, both sexes	1	3	3	3	1	-	-	2	-	-	-	-
Male	-	1	3	-	-	-	-	-	-	-	-	-
Female	1	2	-	3	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	6	2	5	10	7	6	5	2	2	1	-
Male	5	4	2	2	8	5	3	1	-	-	-	-
Female	-	2	-	3	2	2	3	4	2	2	1	-
Alabama												
All races, both sexes	6	6	8	8	8	12	7	6	4	2	1	-
Male	4	5	5	5	5	6	4	1	1	2	-	-
Female	2	1	3	3	3	6	3	5	3	-	1	-
White, both sexes	5	6	6	5	8	8	6	5	3	1	1	-
Male	3	5	4	4	5	4	4	1	-	1	-	-
Female	2	1	2	1	3	4	2	4	3	-	1	-
Black, both sexes	1	-	2	3	-	4	1	1	1	1	-	-
Male	1	-	1	1	-	2	-	-	1	1	-	-
Female	-	-	1	2	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Alaska												
All races, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	7	6	7	9	5	5	3	3	1	-	-
Male	1	5	2	5	6	4	2	-	2	1	-	-
Female	3	2	4	2	3	1	3	3	1	-	-	-
White, both sexes	4	7	6	6	9	5	5	3	3	1	-	-
Male	1	5	2	5	6	4	2	-	2	1	-	-
Female	3	2	4	1	3	1	3	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Arkansas												
All races, both sexes	2	8	2	3	7	5	7	3	1	-	-	-
Male	1	6	1	2	5	4	5	-	1	-	-	-
Female	1	2	1	1	2	1	2	3	-	-	-	-
White, both sexes	2	6	1	3	7	3	6	3	1	-	-	-
Male	1	6	1	2	5	3	5	-	1	-	-	-
Female	1	-	-	1	2	-	1	3	-	-	-	-
Black, both sexes	-	1	1	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	29	21	25	30	29	30	23	27	13	6	-	-
Male	21	16	18	19	15	18	10	8	3	1	-	-
Female	8	5	7	11	14	12	13	19	10	5	-	-
White, both sexes	22	16	19	24	25	23	19	24	12	5	-	-
Male	16	11	14	15	12	15	7	8	3	1	-	-
Female	6	5	5	9	13	8	12	16	9	4	-	-
Black, both sexes	7	4	4	3	2	3	1	2	-	-	-	-
Male	5	4	2	2	1	1	1	-	-	-	-	-
Female	2	-	2	1	1	2	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	2	4	3	1	1	1	-	-
Male	-	1	1	2	2	2	2	-	-	-	-	-
Female	-	-	-	1	-	2	1	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Colorado												
All races, both sexes	5	4	3	2	2	4	4	2	-	-	-	-
Male	3	4	3	-	1	-	2	1	-	-	-	-
Female	2	-	-	2	1	4	2	1	-	-	-	-
White, both sexes	4	4	3	-	2	4	4	2	-	-	-	-
Male	2	4	3	-	1	-	2	1	-	-	-	-
Female	2	-	-	-	1	4	2	1	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	4	1	5	1	4	6	8	4	1	-	-
Male	1	3	-	3	-	3	4	3	2	-	-	-
Female	-	1	1	2	1	1	2	5	2	1	-	-
White, both sexes	1	3	1	4	1	3	4	7	4	1	-	-
Male	1	3	-	3	-	3	3	3	2	-	-	-
Female	-	-	1	1	1	-	1	4	2	1	-	-
Black, both sexes	-	1	-	1	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Delaware												
All races, both sexes	-	2	-	1	-	2	1	1	1	-	-	-
Male	-	2	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	-	1	1	-	-	-
White, both sexes	-	2	-	1	-	2	1	1	1	-	-	-
Male	-	2	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Florida												
All races, both sexes	18	17	18	20	13	24	28	20	16	2	2	-
Male	10	11	10	12	8	15	16	9	4	1	-	-
Female	8	6	8	8	5	9	12	11	12	1	2	-
White, both sexes	15	17	16	17	10	21	26	19	14	2	2	-
Male	8	11	9	11	5	12	16	9	3	1	-	-
Female	7	6	7	6	5	9	10	10	11	1	2	-
Black, both sexes	1	-	2	3	3	3	2	1	2	-	-	-
Male	-	-	1	1	3	3	-	-	1	-	-	-
Female	1	-	1	2	-	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	20	13	8	12	6	5	20	5	2	-	-	-
Male	12	9	5	8	3	2	7	2	1	-	-	-
Female	8	4	3	4	3	3	13	3	1	-	-	-
White, both sexes	13	7	6	12	3	4	15	4	2	-	-	-
Male	8	3	4	8	1	1	6	2	1	-	-	-
Female	5	4	2	4	2	3	9	2	1	-	-	-
Black, both sexes	7	6	2	-	2	-	5	1	-	-	-	-
Male	4	6	1	-	1	-	1	-	-	-	-	-
Female	3	-	1	-	1	-	4	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Hawaii												
All races, both sexes	-	-	1	1	3	-	2	3	-	-	-	-
Male	-	-	1	1	2	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	-	1	3	-	-	-	-
Male	-	-	1	-	2	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Idaho												
All races, both sexes	1	3	-	3	1	2	3	1	-	-	-	-
Male	1	3	-	3	-	1	2	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	1	3	-	3	1	2	3	1	-	-	-	-
Male	1	3	-	3	-	1	2	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Illinois												
All races, both sexes	3	12	10	6	16	10	22	13	7	2	2	-
Male	2	8	7	2	8	6	10	5	1	-	-	-
Female	1	4	3	4	8	4	12	8	6	2	2	-
White, both sexes	2	7	8	6	14	9	22	11	6	2	2	-
Male	1	5	6	2	7	6	10	5	1	-	-	-
Female	1	2	2	4	7	3	12	6	5	2	2	-
Black, both sexes	1	5	2	-	2	1	-	2	1	-	-	-
Male	1	3	1	-	1	-	-	-	-	-	-	-
Female	-	2	1	-	1	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	9	6	4	6	4	2	9	10	5	2	2	-
Male	6	5	1	2	4	2	3	5	-	-	1	-
Female	3	1	3	4	-	-	6	5	5	2	1	-
White, both sexes	6	5	4	6	4	2	9	10	4	2	2	-
Male	4	4	1	2	4	2	3	5	-	-	1	-
Female	2	1	3	4	-	-	6	5	4	2	1	-
Black, both sexes	3	1	-	-	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Iowa												
All races, both sexes	2	1	1	-	1	4	4	4	1	-	-	-
Male	1	1	-	-	1	3	1	1	-	-	-	-
Female	1	-	1	-	-	1	3	3	1	-	-	-
White, both sexes	2	1	1	-	1	3	4	4	1	-	-	-
Male	1	1	-	-	1	2	1	1	-	-	-	-
Female	1	-	1	-	-	1	3	3	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	4	2	2	1	3	3	6	3	1	-	-
Male	1	2	1	1	-	2	3	5	-	-	-	-
Female	1	2	1	1	1	1	-	1	3	1	-	-
White, both sexes	2	4	2	1	1	3	3	6	3	1	-	-
Male	1	2	1	-	-	2	3	5	-	-	-	-
Female	1	2	1	1	1	1	-	1	3	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Kentucky												
All races, both sexes	6	3	9	9	3	4	9	6	1	1	-	-
Male	5	3	6	4	2	2	3	5	-	-	-	-
Female	1	-	3	5	1	2	6	1	1	1	-	-
White, both sexes	5	3	8	8	3	4	9	6	1	1	-	-
Male	4	3	5	4	2	2	3	5	-	-	-	-
Female	1	-	3	4	1	2	6	1	1	1	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	8	6	4	3	4	7	5	-	1	-	-
Male	4	6	6	1	2	1	3	3	-	1	-	-
Female	1	2	-	3	1	3	4	2	-	-	-	-
White, both sexes	3	2	5	2	2	2	6	5	-	-	-	-
Male	3	1	5	1	2	1	2	3	-	-	-	-
Female	-	1	-	1	-	1	4	2	-	-	-	-
Black, both sexes	2	4	1	2	1	2	1	-	-	1	-	-
Male	1	3	1	-	-	-	1	-	-	1	-	-
Female	1	1	-	2	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Maine												
All races, both sexes	1	1	-	2	1	2	-	4	-	2	-	-
Male	1	-	-	1	1	1	-	3	-	1	-	-
Female	-	1	-	1	-	1	-	1	-	1	-	-
White, both sexes	1	1	-	2	1	2	-	4	-	2	-	-
Male	1	-	-	1	1	1	-	3	-	1	-	-
Female	-	1	-	1	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	7	2	6	4	6	9	5	3	-	-	-
Male	3	5	1	3	3	4	4	2	2	-	-	-
Female	1	2	1	3	1	2	5	3	1	-	-	-
White, both sexes	3	4	1	3	3	5	8	4	3	-	-	-
Male	2	3	-	1	3	4	4	2	2	-	-	-
Female	1	1	1	2	-	1	4	2	1	-	-	-
Black, both sexes	-	3	1	3	1	1	1	1	-	-	-	-
Male	-	2	1	2	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Massachusetts												
All races, both sexes	8	6	8	5	9	6	8	5	5	4	1	-
Male	6	3	4	4	5	4	3	1	2	-	-	-
Female	2	3	4	1	4	2	5	4	3	4	1	-
White, both sexes	8	4	8	5	9	5	8	5	5	3	1	-
Male	6	2	4	4	5	3	3	1	2	-	-	-
Female	2	2	4	1	4	2	5	4	3	3	1	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Michigan												
All races, both sexes	11	8	10	13	8	13	13	8	4	2	-	-
Male	9	4	6	6	6	6	3	5	-	1	-	-
Female	2	4	4	7	2	7	10	3	4	1	-	-
White, both sexes	8	4	7	11	7	12	12	6	3	2	-	-
Male	7	1	4	5	5	6	3	5	-	1	-	-
Female	1	3	3	6	2	6	9	1	3	1	-	-
Black, both sexes	3	3	3	2	1	1	-	2	1	-	-	-
Male	2	2	2	1	1	-	-	-	-	-	-	-
Female	1	1	1	1	-	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Minnesota												
All races, both sexes	5	10	4	2	2	5	7	7	3	2	-	-
Male	3	6	3	1	2	3	3	1	1	-	-	-
Female	2	4	1	1	-	2	4	6	2	2	-	-
White, both sexes	5	10	4	2	2	5	7	7	3	2	-	-
Male	3	6	3	1	2	3	3	1	1	-	-	-
Female	2	4	1	1	-	2	4	6	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	1	1	2	2	4	1	3	3	2	-	-
Male	4	1	1	1	-	2	1	1	1	-	-	-
Female	-	-	-	1	2	2	-	2	2	2	-	-
White, both sexes	2	1	1	1	2	2	1	3	3	1	-	-
Male	2	1	1	-	-	-	1	1	1	-	-	-
Female	-	-	-	1	2	2	-	2	2	1	-	-
Black, both sexes	2	-	-	1	-	2	-	-	-	1	-	-
Male	2	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Missouri												
All races, both sexes	7	1	4	6	9	8	9	19	4	1	-	-
Male	6	1	1	3	4	6	6	5	1	-	-	-
Female	1	-	3	3	5	2	3	14	3	1	-	-
White, both sexes	6	1	3	5	8	7	9	19	4	1	-	-
Male	6	1	1	2	4	5	6	5	1	-	-	-
Female	-	-	2	3	4	2	3	14	3	1	-	-
Black, both sexes	1	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	1	-	2	3	-	6	1	1	-	-	-
Male	1	-	-	-	2	-	-	-	1	-	-	-
Female	1	1	-	2	1	-	6	1	-	-	-	-
White, both sexes	2	1	-	2	3	-	6	-	1	-	-	-
Male	1	-	-	-	2	-	-	-	1	-	-	-
Female	1	1	-	2	1	-	6	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Nebraska												
All races, both sexes	2	2	-	1	3	4	4	1	2	2	-	-
Male	1	2	-	-	2	2	1	-	2	1	-	-
Female	1	-	-	1	1	2	3	1	-	1	-	-
White, both sexes	2	2	-	1	3	4	3	1	2	2	-	-
Male	1	2	-	-	2	2	1	-	2	1	-	-
Female	1	-	-	1	1	2	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	4	3	4	3	5	3	4	-	-	-	-
Male	2	3	3	3	2	1	2	1	-	-	-	-
Female	1	1	-	1	1	4	1	3	-	-	-	-
White, both sexes	2	4	2	4	3	5	2	4	-	-	-	-
Male	1	3	2	3	2	1	2	1	-	-	-	-
Female	1	1	-	1	1	4	-	3	-	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
New Hampshire												
All races, both sexes	1	-	-	1	-	5	1	1	1	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	4	1	1	1	-	-	-
White, both sexes	1	-	-	1	-	5	1	1	1	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	4	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	12	2	4	4	6	12	12	7	5	2	1	-
Male	9	1	3	2	4	8	7	2	2	-	1	-
Female	3	1	1	2	2	4	5	5	3	2	-	-
White, both sexes	9	2	3	3	4	10	10	6	4	2	1	-
Male	6	1	2	2	3	8	6	2	2	-	1	-
Female	3	1	1	1	1	2	4	4	2	2	-	-
Black, both sexes	3	-	1	1	1	2	2	1	1	-	-	-
Male	3	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
New Mexico												
All races, both sexes	-	3	-	3	-	1	2	1	-	1	-	-
Male	-	2	-	1	-	-	2	-	-	-	-	-
Female	-	1	-	2	-	1	-	1	-	1	-	-
White, both sexes	-	3	-	2	-	1	2	1	-	1	-	-
Male	-	2	-	1	-	-	2	-	-	-	-	-
Female	-	1	-	1	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	11	13	23	18	23	20	10	9	7	1	-
Male	9	7	8	10	8	12	8	1	3	1	-	-
Female	4	4	5	13	10	11	12	9	6	6	1	-
White, both sexes	6	8	11	18	16	20	16	8	9	7	1	-
Male	5	5	8	8	7	10	7	1	3	1	-	-
Female	1	3	3	10	9	10	9	7	6	6	1	-
Black, both sexes	7	3	2	5	1	3	4	1	-	-	-	-
Male	4	2	-	2	-	2	1	-	-	-	-	-
Female	3	1	2	3	1	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
North Carolina												
All races, both sexes	13	12	8	16	7	16	10	8	6	2	-	-
Male	11	9	6	9	2	7	5	1	1	1	-	-
Female	2	3	2	7	5	9	5	7	5	1	-	-
White, both sexes	7	9	6	15	4	11	7	8	5	2	-	-
Male	6	8	4	8	-	4	3	1	1	1	-	-
Female	1	1	2	7	4	7	4	7	4	1	-	-
Black, both sexes	6	3	2	1	2	4	3	-	1	-	-	-
Male	5	1	2	1	2	2	2	-	-	-	-	-
Female	1	2	-	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	1	-	-	2	-	2	-	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
White, both sexes	1	-	-	-	1	-	-	2	-	2	-	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Ohio												
All races, both sexes	14	7	7	16	15	17	24	19	8	1	-	-
Male	12	5	4	5	7	8	10	10	2	-	-	-
Female	2	2	3	11	8	9	14	9	6	1	-	-
White, both sexes	11	7	5	15	14	15	22	18	8	1	-	-
Male	9	5	4	5	6	6	9	9	2	-	-	-
Female	2	2	1	10	8	9	13	9	6	1	-	-
Black, both sexes	3	-	2	1	1	2	2	1	-	-	-	-
Male	3	-	-	-	1	2	1	1	-	-	-	-
Female	-	-	2	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7	4	6	5	3	8	9	10	2	2	-	-
Male	3	3	3	2	2	5	5	4	-	-	-	-
Female	4	1	3	3	1	3	4	6	2	2	-	-
White, both sexes	6	3	5	4	3	8	8	9	1	2	-	-
Male	3	3	2	2	2	5	5	4	-	-	-	-
Female	3	-	3	2	1	3	3	5	1	2	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Oregon												
All races, both sexes	3	3	4	3	2	2	4	1	1	-	-	-
Male	3	2	2	2	1	1	3	1	-	-	-	-
Female	-	1	2	1	1	1	1	-	1	-	-	-
White, both sexes	3	3	4	3	2	2	4	1	1	-	-	-
Male	3	2	2	2	1	1	3	1	-	-	-	-
Female	-	1	2	1	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	11	3	12	8	17	18	26	21	11	6	1	-
Male	8	3	9	7	12	9	11	9	2	1	1	-
Female	3	-	3	1	5	9	15	12	9	5	-	-
White, both sexes	9	2	9	8	14	16	26	20	11	3	1	-
Male	6	2	7	7	10	9	11	9	2	-	1	-
Female	3	-	2	1	4	7	15	11	9	3	-	-
Black, both sexes	2	1	3	-	3	2	-	1	-	3	-	-
Male	2	1	2	-	2	-	-	-	-	1	-	-
Female	-	-	1	-	1	2	-	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Rhode Island												
All races, both sexes	1	1	-	1	1	5	4	1	1	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	1	-	-	-	1	3	3	-	1	-	-	-
White, both sexes	1	1	-	1	1	5	4	1	1	-	-	-
Male	-	1	-	1	-	2	1	1	-	-	-	-
Female	1	-	-	-	1	3	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	11	10	9	7	5	8	7	4	1	2	-	-
Male	6	5	5	7	5	5	2	2	-	-	-	-
Female	5	5	4	-	-	3	5	2	1	2	-	-
White, both sexes	3	6	6	4	4	6	6	3	1	1	-	-
Male	2	3	3	4	4	3	2	1	-	-	-	-
Female	1	3	3	-	-	3	4	2	1	1	-	-
Black, both sexes	8	4	3	3	1	2	1	1	-	1	-	-
Male	4	2	2	3	1	2	-	1	-	-	-	-
Female	4	2	1	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
South Dakota												
All races, both sexes	1	-	-	-	2	1	2	1	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	1	-	-	-	1	1	2	1	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	7	10	10	9	15	9	6	3	1	-	-
Male	3	5	7	5	5	10	4	5	2	-	-	-
Female	6	2	3	5	4	5	5	1	1	1	-	-
White, both sexes	7	6	9	8	7	15	7	6	3	1	-	-
Male	2	4	6	4	3	10	2	5	2	-	-	-
Female	5	2	3	4	4	5	5	1	1	1	-	-
Black, both sexes	2	1	1	2	2	-	2	-	-	-	-	-
Male	1	1	1	1	2	-	2	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Texas												
All races, both sexes	18	17	13	22	21	20	33	18	8	3	2	-
Male	13	12	11	14	12	10	16	8	2	-	-	-
Female	5	5	2	8	9	10	17	10	6	3	2	-
White, both sexes	15	15	12	20	17	20	32	16	7	3	1	-
Male	11	11	10	13	9	10	16	8	2	-	-	-
Female	4	4	2	7	8	10	16	8	5	3	1	-
Black, both sexes	3	2	1	1	3	-	1	2	1	-	-	-
Male	2	1	1	1	2	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	2	1	4	1	3	-	6	2	1	-	-	-
Male	1	1	2	-	2	-	4	1	-	-	-	-
Female	1	-	2	1	1	-	2	1	1	-	-	-
White, both sexes	2	1	4	1	2	-	6	2	1	-	-	-
Male	1	1	2	-	1	-	4	1	-	-	-	-
Female	1	-	2	1	1	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Vermont												
All races, both sexes	2	-	1	-	-	1	2	-	2	-	-	-
Male	1	-	1	-	-	-	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	2	-	1	-	-	1	2	-	2	-	-	-
Male	1	-	1	-	-	-	1	-	1	-	-	-
Female	1	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	7	4	7	15	13	11	9	4	-	-	-
Male	2	5	2	6	7	4	4	1	1	-	-	-
Female	1	2	2	1	8	9	7	8	3	-	-	-
White, both sexes	3	4	3	6	10	12	9	9	2	-	-	-
Male	2	3	2	5	6	4	4	1	1	-	-	-
Female	1	1	1	1	4	8	5	8	1	-	-	-
Black, both sexes	-	3	1	1	5	1	1	-	2	-	-	-
Male	-	2	-	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	4	1	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Washington												
All races, both sexes	4	8	3	3	7	9	4	2	3	-	-	-
Male	3	3	1	2	6	4	3	1	2	-	-	-
Female	1	5	2	1	1	5	1	1	1	-	-	-
White, both sexes	3	4	3	3	7	9	3	2	3	-	-	-
Male	2	1	1	2	6	4	2	1	2	-	-	-
Female	1	3	2	1	1	5	1	1	1	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	1	3	1	9	3	4	5	-	-	-
Male	1	1	1	2	-	5	-	-	-	-	-	-
Female	-	-	-	1	1	4	3	4	5	-	-	-
White, both sexes	1	1	1	2	1	9	3	3	5	-	-	-
Male	1	1	1	1	-	5	-	-	-	-	-	-
Female	-	-	-	1	1	4	3	3	5	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Wisconsin												
All races, both sexes	6	4	-	9	7	4	10	8	6	-	-	-
Male	3	2	-	4	4	3	4	2	2	-	-	-
Female	3	2	-	5	3	1	6	6	4	-	-	-
White, both sexes	6	4	-	9	7	4	10	8	6	-	-	-
Male	3	2	-	4	4	3	4	2	2	-	-	-
Female	3	2	-	5	3	1	6	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	1	1	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
White, both sexes	-	-	1	1	1	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
United States												
All races, both sexes	1,509	15	4	1	1	1	3	5	6	18	24	35
Male	705	10	3	-	1	-	2	4	3	10	15	22
Female	804	5	1	1	-	1	1	1	3	8	9	13
White, both sexes	1,291	10	3	1	1	1	1	4	4	13	17	26
Male	597	7	2	-	1	-	1	4	3	8	10	16
Female	694	3	1	1	-	1	-	-	1	5	7	10
Black, both sexes	167	5	1	-	-	-	2	1	2	4	5	7
Male	79	3	1	-	-	-	1	-	-	2	3	4
Female	88	2	-	-	-	-	1	1	2	2	2	3
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	2	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	1	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Alaska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	19	-	-	-	-	-	-	-	-	2	1	-
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Arkansas												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	155	3	1	-	-	-	-	-	-	2	3	2
Male	79	3	1	-	-	-	-	-	-	1	1	1
Female	76	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	130	2	1	-	-	-	-	-	-	2	3	2
Male	65	2	1	-	-	-	-	-	-	1	1	1
Female	65	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Colorado												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Florida												
All races, both sexes	97	2	-	-	-	-	-	1	-	1	-	2
Male	47	1	-	-	-	-	-	-	-	1	-	1
Female	50	1	-	-	-	-	-	1	-	-	-	1
White, both sexes	87	1	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	45	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	1	-	1	-	1
Male	4	1	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	27	1	-	-	-	-	-	-	-	-	-	1
Male	12	1	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	1	-	-	-	-	-	-	-	-	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Hawaii												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Illinois												
All races, both sexes	53	1	-	-	-	-	1	-	1	-	2	2
Male	25	1	-	-	-	-	-	-	-	-	2	2
Female	28	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	45	1	-	-	-	-	-	-	1	-	1	1
Male	20	1	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	8	-	-	-	-	-	1	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	26	-	-	1	1	-	-	-	-	1	1	-
Male	13	-	-	-	1	-	-	-	-	-	1	-
Female	13	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	24	-	-	1	1	-	-	-	-	1	1	-
Male	12	-	-	-	1	-	-	-	-	-	1	-
Female	12	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Kentucky												
All races, both sexes	18	-	1	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	15	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	24	-	-	-	-	-	-	1	-	1	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Massachusetts												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	-	3
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	32	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	3
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Minnesota												
All races, both sexes	33	-	-	-	-	-	-	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	13	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Missouri												
All races, both sexes	32	-	-	-	-	1	-	-	-	1	-	3
Male	18	-	-	-	-	-	-	-	-	-	-	3
Female	14	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	28	-	-	-	-	1	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Nebraska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
New Hampshire												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	39	1	-	-	-	-	-	-	1	1	-	1
Male	12	1	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	34	1	-	-	-	-	-	-	1	1	-	1
Male	11	1	-	-	-	-	-	-	1	-	-	1
Female	23	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
New Mexico												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	110	2	1	-	-	-	1	-	1	1	1	3
Male	48	1	1	-	-	-	1	-	1	-	1	1
Female	62	1	-	-	-	-	-	-	-	1	-	2
White, both sexes	95	1	-	-	-	-	-	-	1	-	1	2
Male	42	1	-	-	-	-	-	-	1	-	1	-
Female	53	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	13	1	1	-	-	-	1	-	-	-	-	1
Male	5	-	1	-	-	-	1	-	-	-	-	1
Female	8	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
North Carolina												
All races, both sexes	46	1	-	-	-	-	-	-	-	-	1	-
Male	20	1	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	1	-	-	-	-	-	-	-	-	-	-
Male	15	1	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Ohio												
All races, both sexes	73	-	-	-	-	-	-	1	-	-	-	3
Male	34	-	-	-	-	-	-	1	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	63	-	-	-	-	-	-	1	-	-	-	2
Male	31	-	-	-	-	-	-	1	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Oregon												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	86	-	-	-	-	-	-	-	-	1	1	2
Male	32	-	-	-	-	-	-	-	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	76	-	-	-	-	-	-	-	-	1	-	2
Male	29	-	-	-	-	-	-	-	-	1	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Texas												
All races, both sexes	102	1	-	-	-	-	-	2	-	3	1	2
Male	47	-	-	-	-	-	-	2	-	3	-	-
Female	55	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	85	-	-	-	-	-	-	2	-	3	1	1
Male	41	-	-	-	-	-	-	2	-	3	-	-
Female	44	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	13	1	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	40	-	-	-	-	-	-	-	-	1	1	1
Male	26	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Washington												
All races, both sexes	25	-	1	-	-	-	-	-	-	-	3	2
Male	11	-	1	-	-	-	-	-	-	-	2	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	24	-	1	-	-	-	-	-	-	-	3	2
Male	10	-	1	-	-	-	-	-	-	-	2	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	21	3	-	-	-	-	1	-	-	-	-	-
Male	11	1	-	-	-	-	1	-	-	-	-	-
Female	10	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	2	-	-	-	-	1	-	-	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	10	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Wisconsin												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
United States												
All races, both sexes	44	59	97	114	119	194	257	267	173	59	13	-
Male	23	44	60	66	57	104	117	104	47	11	2	-
Female	21	15	37	48	62	90	140	163	126	48	11	-
White, both sexes	31	44	75	94	106	171	222	245	157	54	11	-
Male	16	34	46	56	50	92	102	96	42	9	2	-
Female	15	10	29	38	56	79	120	149	115	45	9	-
Black, both sexes	12	13	16	15	8	17	24	18	11	4	2	-
Male	6	8	10	7	4	9	10	6	4	1	-	-
Female	6	5	6	8	4	8	14	12	7	3	2	-
American Indian 1/, both sexes	-	1	-	1	1	1	-	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	6	4	4	5	11	4	4	1	-	-
Male	1	1	4	2	3	3	5	2	1	1	-	-
Female	-	-	2	2	1	2	6	2	3	-	-	-
Alabama												
All races, both sexes	1	1	-	2	2	1	3	3	4	1	1	-
Male	-	1	-	1	-	1	-	-	2	-	-	-
Female	1	-	-	1	2	-	3	3	2	1	1	-
White, both sexes	1	1	-	2	2	1	3	1	3	1	1	-
Male	-	1	-	1	-	1	-	-	2	-	-	-
Female	1	-	-	1	2	-	3	1	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Alaska												
All races, both sexes	-	-	1	-	2	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	2	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	2	3	2	1	3	2	1	1	1	-
Male	-	-	1	1	1	-	1	1	1	-	-	-
Female	-	-	1	2	1	1	2	1	-	1	1	-
White, both sexes	-	-	1	3	2	1	3	2	-	1	1	-
Male	-	-	1	1	1	-	1	1	-	-	-	-
Female	-	-	-	2	1	1	2	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Arkansas												
All races, both sexes	-	1	-	2	1	1	2	3	2	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	1	-	1	1	1	1	2	2	-	-	-
White, both sexes	-	-	-	2	1	1	2	3	2	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	1	1	1	1	2	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	6	14	13	12	17	28	29	14	3	1	-
Male	3	5	9	6	7	11	14	12	4	1	-	-
Female	4	1	5	7	5	6	14	17	10	2	1	-
White, both sexes	4	6	10	12	9	14	24	25	12	3	1	-
Male	2	5	6	5	5	9	11	11	4	1	-	-
Female	2	1	4	7	4	5	13	14	8	2	1	-
Black, both sexes	2	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	2	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	-	3	3	3	3	1	-	-	-
Male	1	-	3	-	2	2	2	1	-	-	-	-
Female	-	-	-	-	1	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Colorado												
All races, both sexes	1	2	1	1	1	5	5	3	1	1	-	-
Male	-	2	1	1	-	2	2	1	1	1	-	-
Female	1	-	-	-	1	3	3	2	-	-	-	-
White, both sexes	1	2	-	1	1	5	5	3	1	1	-	-
Male	-	2	-	1	-	2	2	1	1	1	-	-
Female	1	-	-	-	1	3	3	2	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	-	1	1	2	2	5	2	1	1	-
Male	1	1	-	-	-	2	1	1	-	-	1	-
Female	-	-	-	1	1	-	1	4	2	1	-	-
White, both sexes	1	-	-	1	1	1	2	5	2	1	1	-
Male	1	-	-	-	-	1	1	1	-	-	1	-
Female	-	-	-	1	1	-	1	4	2	1	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Delaware												
All races, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Florida												
All races, both sexes	1	3	5	6	6	14	15	24	12	4	1	-
Male	1	2	4	3	4	8	9	10	2	1	-	-
Female	-	1	1	3	2	6	6	14	10	3	1	-
White, both sexes	1	3	5	5	6	12	14	24	11	4	-	-
Male	1	2	4	2	4	8	8	10	1	1	-	-
Female	-	1	1	3	2	4	6	14	10	3	-	-
Black, both sexes	-	-	-	1	-	2	1	-	-	-	1	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	1	4	1	5	6	4	2	1	-	-
Male	-	1	-	3	-	3	2	-	1	-	-	-
Female	-	-	1	1	1	2	4	4	1	1	-	-
White, both sexes	-	1	-	4	1	4	6	4	2	1	-	-
Male	-	1	-	3	-	2	2	-	1	-	-	-
Female	-	-	-	1	1	2	4	4	1	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Hawaii												
All races, both sexes	-	-	-	1	-	1	3	-	2	2	-	-
Male	-	-	-	-	-	1	1	-	-	2	-	-
Female	-	-	-	1	-	-	2	-	2	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	-	2	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	-	2	-	2	-	-	-
Idaho												
All races, both sexes	-	-	-	2	1	-	-	3	-	-	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	1	-	-	3	-	-	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Illinois												
All races, both sexes	-	3	4	3	6	3	6	12	6	2	1	-
Male	-	2	3	3	3	2	1	4	2	-	-	-
Female	-	1	1	-	3	1	5	8	4	2	1	-
White, both sexes	-	2	4	3	5	2	6	11	5	2	1	-
Male	-	1	3	3	2	1	1	4	2	-	-	-
Female	-	1	1	-	3	1	5	7	3	2	1	-
Black, both sexes	-	1	-	-	1	1	-	1	1	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	4	2	1	3	5	2	4	1	-	-
Male	-	-	3	2	1	1	4	-	-	-	-	-
Female	-	-	1	-	-	2	1	2	4	1	-	-
White, both sexes	-	-	3	2	1	3	5	1	4	1	-	-
Male	-	-	2	2	1	1	4	-	-	-	-	-
Female	-	-	1	-	-	2	1	1	4	1	-	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Iowa												
All races, both sexes	1	-	1	2	3	2	5	5	3	-	-	-
Male	1	-	1	1	2	2	1	3	-	-	-	-
Female	-	-	-	1	1	-	4	2	3	-	-	-
White, both sexes	1	-	1	2	3	2	5	5	3	-	-	-
Male	1	-	1	1	2	2	1	3	-	-	-	-
Female	-	-	-	1	1	-	4	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	-	-	4	3	3	2	-	-	-
Male	-	1	1	-	-	2	3	1	1	-	-	-
Female	-	-	-	-	-	2	-	2	1	-	-	-
White, both sexes	-	1	1	-	-	4	2	3	2	-	-	-
Male	-	1	1	-	-	2	2	1	1	-	-	-
Female	-	-	-	-	-	2	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Kentucky												
All races, both sexes	-	1	-	1	1	3	3	4	1	1	1	-
Male	-	-	-	1	1	1	1	3	-	-	1	-
Female	-	1	-	-	-	2	2	1	1	1	-	-
White, both sexes	-	1	-	1	1	3	2	3	1	1	1	-
Male	-	-	-	1	1	1	1	2	-	-	1	-
Female	-	1	-	-	-	2	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	3	3	2	3	2	5	1	1	-	-
Male	-	1	-	2	1	-	1	2	-	-	-	-
Female	1	-	3	1	1	3	1	3	1	1	-	-
White, both sexes	-	1	2	1	1	3	-	5	1	1	-	-
Male	-	1	-	-	1	-	-	2	-	-	-	-
Female	-	-	2	1	-	3	-	3	1	1	-	-
Black, both sexes	1	-	1	2	1	-	1	-	-	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	1	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Maine												
All races, both sexes	-	-	1	2	-	-	1	1	2	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	2	-	-	1	1	2	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	1	6	2	2	1	5	6	3	-	-	-
Male	-	1	3	1	1	1	4	-	-	-	-	-
Female	-	-	3	1	1	-	1	6	3	-	-	-
White, both sexes	-	1	3	2	1	1	3	6	2	-	-	-
Male	-	1	1	1	-	1	3	-	-	-	-	-
Female	-	-	2	1	1	-	-	6	2	-	-	-
Black, both sexes	-	-	3	-	1	-	2	-	1	-	-	-
Male	-	-	2	-	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Massachusetts												
All races, both sexes	1	1	2	2	3	6	5	2	8	2	1	-
Male	-	1	1	1	2	2	2	1	1	-	-	-
Female	1	-	1	1	1	4	3	1	7	2	1	-
White, both sexes	1	-	2	2	3	5	4	2	7	2	1	-
Male	-	-	1	1	2	1	2	1	1	-	-	-
Female	1	-	1	1	1	4	2	1	6	2	1	-
Black, both sexes	-	1	-	-	-	1	1	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	6	5	3	6	10	12	9	7	3	-	-
Male	-	3	5	3	2	6	5	5	1	-	-	-
Female	-	3	-	-	4	4	7	4	6	3	-	-
White, both sexes	-	3	3	3	5	8	9	7	7	2	-	-
Male	-	2	3	3	2	4	4	4	1	-	-	-
Female	-	1	-	-	3	4	5	3	6	2	-	-
Black, both sexes	-	3	2	-	1	2	3	2	-	1	-	-
Male	-	1	2	-	-	2	1	1	-	-	-	-
Female	-	2	-	-	1	-	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Minnesota												
All races, both sexes	-	2	1	2	2	3	2	12	3	3	-	-
Male	-	2	1	1	2	3	-	4	1	-	-	-
Female	-	-	-	1	-	-	2	8	2	3	-	-
White, both sexes	-	2	1	2	1	3	2	12	3	3	-	-
Male	-	2	1	1	1	3	-	4	1	-	-	-
Female	-	-	-	1	-	-	2	8	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	1	1	2	1	2	-	-	1	1	-	-
Male	2	1	-	2	1	-	-	-	1	1	-	-
Female	-	-	1	-	-	2	-	-	-	-	-	-
White, both sexes	-	1	-	1	1	-	-	-	1	1	-	-
Male	-	1	-	1	1	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	1	-	2	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Missouri												
All races, both sexes	1	2	-	5	2	4	4	3	5	1	-	-
Male	1	1	-	3	1	2	3	2	1	1	-	-
Female	-	1	-	2	1	2	1	1	4	-	-	-
White, both sexes	-	1	-	5	2	4	4	3	5	1	-	-
Male	-	-	-	3	1	2	3	2	1	1	-	-
Female	-	1	-	2	1	2	1	1	4	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Nebraska												
All races, both sexes	-	1	-	-	-	1	2	1	2	2	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	2	-	2	2	-	-
White, both sexes	-	1	-	-	-	1	2	1	2	2	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	2	-	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	1	4	1	3	1	1	-	-	-	-
Male	1	-	1	3	1	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
White, both sexes	1	-	1	3	1	2	1	1	-	-	-	-
Male	1	-	1	2	1	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
New Hampshire												
All races, both sexes	-	-	1	-	1	1	1	4	-	2	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	4	-	2	-	-
White, both sexes	-	-	1	-	1	1	1	4	-	2	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	4	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	4	-	2	2	7	7	6	4	2	-	-
Male	-	4	-	-	-	1	2	1	1	-	-	-
Female	1	-	-	2	2	6	5	5	3	2	-	-
White, both sexes	1	3	-	-	2	7	6	6	3	2	-	-
Male	-	3	-	-	-	1	2	1	1	-	-	-
Female	1	-	-	-	2	6	4	5	2	2	-	-
Black, both sexes	-	1	-	2	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
New Mexico												
All races, both sexes	-	1	-	-	1	2	8	1	-	1	-	-
Male	-	1	-	-	-	1	3	1	-	-	-	-
Female	-	-	-	-	1	1	5	-	-	1	-	-
White, both sexes	-	1	-	-	-	2	8	1	-	1	-	-
Male	-	1	-	-	-	1	3	1	-	-	-	-
Female	-	-	-	-	-	1	5	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	8	5	11	18	19	16	13	6	1	-
Male	1	-	4	1	7	9	8	7	4	1	-	-
Female	1	1	4	4	4	9	11	9	9	5	1	-
White, both sexes	1	-	8	4	11	16	15	16	13	5	1	-
Male	-	-	4	1	7	8	7	7	4	1	-	-
Female	1	-	4	3	4	8	8	9	9	4	1	-
Black, both sexes	1	1	-	1	-	1	4	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
North Carolina												
All races, both sexes	4	1	7	5	4	7	6	5	4	-	1	-
Male	2	-	3	4	-	4	3	1	1	-	-	-
Female	2	1	4	1	4	3	3	4	3	-	1	-
White, both sexes	2	1	4	4	3	7	5	4	2	-	-	-
Male	1	-	2	4	-	4	2	1	-	-	-	-
Female	1	1	2	-	3	3	3	3	2	-	-	-
Black, both sexes	2	-	1	1	1	-	1	1	2	-	1	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	1	1	1	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	1	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	1	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Ohio												
All races, both sexes	3	3	5	10	7	6	15	15	4	1	-	-
Male	2	-	3	6	5	2	5	7	-	1	-	-
Female	1	3	2	4	2	4	10	8	4	-	-	-
White, both sexes	3	2	5	9	7	5	12	12	4	1	-	-
Male	2	-	3	6	5	2	5	5	-	1	-	-
Female	1	2	2	3	2	3	7	7	4	-	-	-
Black, both sexes	-	1	-	1	-	-	2	3	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	1	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	1	2	-	1	1	6	4	-	-	-
Male	-	1	-	1	-	1	1	4	1	-	-	-
Female	-	-	1	1	-	-	-	2	3	-	-	-
White, both sexes	-	-	1	1	-	1	1	6	4	-	-	-
Male	-	-	-	-	-	1	1	4	1	-	-	-
Female	-	-	1	1	-	-	-	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Oregon												
All races, both sexes	1	1	2	-	1	2	5	6	1	1	-	-
Male	1	1	2	-	1	-	2	4	-	-	-	-
Female	-	-	-	-	-	2	3	2	1	1	-	-
White, both sexes	1	1	2	-	1	2	5	6	1	1	-	-
Male	1	1	2	-	1	-	2	4	-	-	-	-
Female	-	-	-	-	-	2	3	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	2	1	3	13	19	15	17	9	-	-
Male	2	-	1	1	1	7	7	5	5	1	-	-
Female	1	-	1	-	2	6	12	10	12	8	-	-
White, both sexes	1	-	2	-	3	12	17	13	16	9	-	-
Male	1	-	1	-	1	7	7	5	4	1	-	-
Female	-	-	1	-	2	5	10	8	12	8	-	-
Black, both sexes	2	-	-	1	-	1	1	2	1	-	-	-
Male	1	-	-	1	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Rhode Island												
All races, both sexes	1	-	-	-	1	-	3	3	2	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	2	-	-	-
White, both sexes	1	-	-	-	1	-	3	3	2	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	-	-	2	5	2	4	1	-	-	-
Male	1	1	-	-	-	2	2	1	-	-	-	-
Female	-	-	-	-	2	3	-	3	1	-	-	-
White, both sexes	1	-	-	-	2	3	2	4	1	-	-	-
Male	1	-	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	-	2	2	-	3	1	-	-	-
Black, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
South Dakota												
All races, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	2	2	2	1	8	6	4	6	-	-	-
Male	1	2	2	2	-	6	5	-	4	-	-	-
Female	-	-	-	-	1	2	1	4	2	-	-	-
White, both sexes	1	1	1	2	1	8	5	4	5	-	-	-
Male	1	1	1	2	-	6	4	-	3	-	-	-
Female	-	-	-	-	1	2	1	4	2	-	-	-
Black, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	1	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Texas												
All races, both sexes	4	5	5	10	14	16	11	13	10	4	1	-
Male	1	4	4	5	5	10	5	5	3	-	-	-
Female	3	1	1	5	9	6	6	8	7	4	1	-
White, both sexes	3	4	4	7	13	16	9	9	9	3	1	-
Male	1	3	3	4	5	10	4	3	3	-	-	-
Female	2	1	1	3	8	6	5	6	6	3	1	-
Black, both sexes	1	-	1	1	1	-	2	3	1	1	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	1	-	-	1	1	-	1	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	2	1	1	1	2	1	3	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	-	-	2	-	1	-	-	1	2	-	-	-
White, both sexes	-	-	2	1	1	1	2	1	3	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	-	-	2	-	1	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Vermont												
All races, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	3	5	3	3	11	7	3	-	-	-
Male	1	-	2	3	3	2	7	5	2	-	-	-
Female	1	-	1	2	-	1	4	2	1	-	-	-
White, both sexes	2	-	2	2	1	2	8	7	3	-	-	-
Male	1	-	1	2	1	1	5	5	2	-	-	-
Female	1	-	1	-	-	1	3	2	1	-	-	-
Black, both sexes	-	-	1	3	2	1	2	-	-	-	-	-
Male	-	-	1	1	2	1	1	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Washington												
All races, both sexes	-	1	-	1	2	3	4	4	3	1	-	-
Male	-	1	-	-	-	2	1	1	2	-	-	-
Female	-	-	-	1	2	1	3	3	1	1	-	-
White, both sexes	-	1	-	1	2	2	4	4	3	1	-	-
Male	-	1	-	-	-	1	1	1	2	-	-	-
Female	-	-	-	1	2	1	3	3	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	1	1	1	2	4	2	1	3	-	-	-
Male	-	1	1	-	1	3	-	1	2	-	-	-
Female	2	-	-	1	1	1	2	-	1	-	-	-
White, both sexes	2	1	1	1	2	4	2	-	3	-	-	-
Male	-	1	1	-	1	3	-	-	2	-	-	-
Female	2	-	-	1	1	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Wisconsin												
All races, both sexes	-	-	3	1	1	2	7	8	4	-	1	-
Male	-	-	1	1	1	1	3	3	-	-	-	-
Female	-	-	2	-	-	1	4	5	4	-	1	-
White, both sexes	-	-	3	1	1	1	7	8	4	-	1	-
Male	-	-	1	1	1	1	3	3	-	-	-	-
Female	-	-	2	-	-	-	4	5	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
United States												
All races, both sexes	8,023	44	20	3	1	6	7	17	15	40	84	153
Male	3,649	24	13	3	-	4	6	10	9	30	54	103
Female	4,374	20	7	-	1	2	1	7	6	10	30	50
White, both sexes	6,980	29	13	2	1	5	4	11	10	32	62	116
Male	3,141	17	8	2	-	3	4	7	7	24	40	79
Female	3,839	12	5	-	1	2	-	4	3	8	22	37
Black, both sexes	900	15	7	1	-	1	3	6	3	8	17	31
Male	427	7	5	1	-	1	2	3	1	6	9	21
Female	473	8	2	-	-	-	1	3	2	2	8	10
American Indian 1/, both sexes	40	-	-	-	-	-	-	-	-	-	3	3
Male	21	-	-	-	-	-	-	-	-	-	3	2
Female	19	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	103	-	-	-	-	-	-	-	2	-	2	3
Male	60	-	-	-	-	-	-	-	1	-	2	1
Female	43	-	-	-	-	-	-	-	1	-	-	2
Alabama												
All races, both sexes	176	2	-	-	-	-	-	-	1	-	5	5
Male	81	2	-	-	-	-	-	-	1	-	5	3
Female	95	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	131	1	-	-	-	-	-	-	1	-	4	3
Male	64	1	-	-	-	-	-	-	1	-	4	3
Female	67	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	45	1	-	-	-	-	-	-	-	-	1	2
Male	17	1	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Alaska												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	132	1	1	-	-	-	-	1	-	-	2	2
Male	64	1	1	-	-	-	-	1	-	-	2	1
Female	68	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	114	-	1	-	-	-	-	-	-	-	1	-
Male	55	-	1	-	-	-	-	-	-	-	1	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	1	-	-	-	-	-	1	-	-	-	-
Male	4	1	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Arkansas												
All races, both sexes	125	-	-	-	-	-	-	-	-	-	5	1
Male	54	-	-	-	-	-	-	-	-	-	3	-
Female	71	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	107	-	-	-	-	-	-	-	-	-	3	1
Male	46	-	-	-	-	-	-	-	-	-	2	-
Female	61	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	314	7	-	-	-	1	-	3	2	1	5	3
Male	157	3	-	-	-	1	-	2	2	1	2	1
Female	157	4	-	-	-	-	-	1	-	-	3	2
White, both sexes	282	6	-	-	-	-	-	3	2	1	5	2
Male	138	2	-	-	-	-	-	2	2	1	2	1
Female	144	4	-	-	-	-	-	1	-	-	3	1
Black, both sexes	14	1	-	-	-	1	-	-	-	-	-	-
Male	9	1	-	-	-	1	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Colorado												
All races, both sexes	114	1	-	-	-	-	-	-	-	1	-	1
Male	54	-	-	-	-	-	-	-	-	-	-	1
Female	60	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	110	-	-	-	-	-	-	-	-	1	-	1
Male	54	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	147	2	1	-	-	-	-	-	-	4	-	1
Male	66	1	1	-	-	-	-	-	-	3	-	1
Female	81	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	141	1	-	-	-	-	-	-	-	2	-	1
Male	62	1	-	-	-	-	-	-	-	2	-	1
Female	79	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	1	1	-	-	-	-	-	-	2	-	-
Male	4	-	1	-	-	-	-	-	-	1	-	-
Female	2	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Delaware												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Florida												
All races, both sexes	360	2	2	-	-	-	-	-	-	-	2	6
Male	167	2	-	-	-	-	-	-	-	-	2	5
Female	193	-	2	-	-	-	-	-	-	-	-	1
White, both sexes	322	-	2	-	-	-	-	-	-	-	-	3
Male	146	-	-	-	-	-	-	-	-	-	-	3
Female	176	-	2	-	-	-	-	-	-	-	-	-
Black, both sexes	36	2	-	-	-	-	-	-	-	-	2	3
Male	20	2	-	-	-	-	-	-	-	-	2	2
Female	16	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	253	2	1	-	1	-	-	2	-	4	8	9
Male	115	1	1	-	-	-	-	1	-	3	3	7
Female	138	1	-	-	1	-	-	1	-	1	5	2
White, both sexes	174	1	-	-	1	-	-	1	-	1	5	5
Male	82	1	-	-	-	-	-	1	-	1	2	5
Female	92	-	-	-	1	-	-	-	-	-	3	-
Black, both sexes	77	1	1	-	-	-	-	1	-	3	3	3
Male	33	-	1	-	-	-	-	-	-	2	1	2
Female	44	1	-	-	-	-	-	1	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Hawaii												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	2	-
Male	28	-	-	-	-	-	-	-	-	-	2	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Illinois												
All races, both sexes	363	-	1	-	-	-	-	-	2	-	5	9
Male	165	-	-	-	-	-	-	-	1	-	3	6
Female	198	-	1	-	-	-	-	-	1	-	2	3
White, both sexes	322	-	1	-	-	-	-	-	2	-	5	8
Male	141	-	-	-	-	-	-	-	1	-	3	5
Female	181	-	1	-	-	-	-	-	1	-	2	3
Black, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	196	2	1	-	-	-	-	-	-	-	2	5
Male	98	1	1	-	-	-	-	-	-	-	2	4
Female	98	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	182	2	1	-	-	-	-	-	-	-	2	4
Male	91	1	1	-	-	-	-	-	-	-	2	3
Female	91	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Iowa												
All races, both sexes	87	-	-	-	-	-	-	1	1	-	-	1
Male	32	-	-	-	-	-	-	1	1	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	84	-	-	-	-	-	-	1	-	-	-	-
Male	30	-	-	-	-	-	-	1	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	87	-	-	1	-	1	-	-	-	2	-	-
Male	38	-	-	1	-	-	-	-	-	1	-	-
Female	49	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	83	-	-	1	-	1	-	-	-	2	-	-
Male	36	-	-	1	-	-	-	-	-	1	-	-
Female	47	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Kentucky												
All races, both sexes	125	-	1	-	-	-	1	-	-	2	1	6
Male	61	-	-	-	-	-	1	-	-	2	1	4
Female	64	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	118	-	1	-	-	-	1	-	-	2	-	6
Male	56	-	-	-	-	-	1	-	-	2	-	4
Female	62	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	146	-	-	-	-	-	-	-	-	-	3	6
Male	76	-	-	-	-	-	-	-	-	-	3	6
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	109	-	-	-	-	-	-	-	-	-	2	4
Male	52	-	-	-	-	-	-	-	-	-	2	4
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	37	-	-	-	-	-	-	-	-	-	1	2
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Maine												
All races, both sexes	56	1	-	-	-	-	-	-	-	-	-	1
Male	22	1	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	1	-	-	-	-	-	-	-	-	-	1
Male	22	1	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	170	2	-	-	-	-	-	2	1	1	4	1
Male	88	2	-	-	-	-	-	1	-	1	2	1
Female	82	-	-	-	-	-	-	1	1	-	2	-
White, both sexes	122	2	-	-	-	-	-	-	-	-	1	1
Male	63	2	-	-	-	-	-	-	-	-	1	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	46	-	-	-	-	-	-	2	-	1	3	-
Male	25	-	-	-	-	-	-	1	-	1	1	-
Female	21	-	-	-	-	-	-	1	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Massachusetts												
All races, both sexes	293	-	-	-	-	-	-	-	-	1	2	7
Male	132	-	-	-	-	-	-	-	-	1	2	4
Female	161	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	281	-	-	-	-	-	-	-	-	1	2	5
Male	124	-	-	-	-	-	-	-	-	1	2	3
Female	157	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	300	2	1	-	-	-	2	1	1	2	2	4
Male	134	1	1	-	-	-	2	1	1	1	2	2
Female	166	1	-	-	-	-	-	-	-	1	-	2
White, both sexes	268	-	1	-	-	-	2	-	1	2	2	2
Male	116	-	1	-	-	-	2	-	1	1	2	-
Female	152	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	31	2	-	-	-	-	-	1	-	-	-	2
Male	18	1	-	-	-	-	-	1	-	-	-	2
Female	13	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Minnesota												
All races, both sexes	147	-	-	-	-	1	-	-	-	-	1	3
Male	70	-	-	-	-	1	-	-	-	-	1	1
Female	77	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	139	-	-	-	-	1	-	-	-	-	-	3
Male	63	-	-	-	-	1	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	124	1	2	-	-	-	1	1	2	-	-	1
Male	51	-	2	-	-	-	1	-	-	-	-	-
Female	73	1	-	-	-	-	-	1	2	-	-	1
White, both sexes	75	-	1	-	-	-	-	-	-	-	-	-
Male	30	-	1	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	49	1	1	-	-	-	1	1	2	-	-	1
Male	21	-	1	-	-	-	1	-	-	-	-	-
Female	28	1	-	-	-	-	-	1	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Missouri												
All races, both sexes	218	1	-	-	-	-	-	-	-	2	1	3
Male	96	-	-	-	-	-	-	-	-	1	1	2
Female	122	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	201	-	-	-	-	-	-	-	-	2	1	2
Male	89	-	-	-	-	-	-	-	-	1	1	1
Female	112	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	15	1	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	37	-	-	-	-	-	-	-	-	1	1	2
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	36	-	-	-	-	-	-	-	-	1	1	2
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Nebraska												
All races, both sexes	60	-	-	1	-	-	-	-	-	-	-	-
Male	27	-	-	1	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	1	-	-	-	-	-	-	-	-
Male	25	-	-	1	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	75	-	1	-	-	-	-	-	-	-	2	-
Male	43	-	1	-	-	-	-	-	-	-	2	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	1	-	-	-	-	-	-	-	2	-
Male	41	-	1	-	-	-	-	-	-	-	2	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
New Hampshire												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	264	-	2	-	-	1	-	-	-	-	2	8
Male	118	-	-	-	-	1	-	-	-	-	1	4
Female	146	-	2	-	-	-	-	-	-	-	1	4
White, both sexes	232	-	1	-	-	1	-	-	-	-	2	6
Male	107	-	-	-	-	1	-	-	-	-	1	3
Female	125	-	1	-	-	-	-	-	-	-	1	3
Black, both sexes	28	-	1	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
New Mexico												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	437	-	1	-	-	-	1	-	-	1	5	7
Male	185	-	1	-	-	-	1	-	-	1	3	5
Female	252	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	377	-	1	-	-	-	1	-	-	1	5	7
Male	162	-	1	-	-	-	1	-	-	1	3	5
Female	215	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
North Carolina												
All races, both sexes	259	6	1	-	-	-	-	2	1	2	1	3
Male	118	1	-	-	-	-	-	2	-	2	1	1
Female	141	5	1	-	-	-	-	-	1	-	-	2
White, both sexes	185	4	-	-	-	-	-	2	1	1	1	3
Male	85	1	-	-	-	-	-	2	-	1	1	1
Female	100	3	-	-	-	-	-	-	1	-	-	2
Black, both sexes	68	2	1	-	-	-	-	-	-	1	-	-
Male	33	-	-	-	-	-	-	-	-	1	-	-
Female	35	2	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Ohio												
All races, both sexes	360	2	1	-	-	-	-	1	1	2	-	7
Male	151	1	1	-	-	-	-	-	-	2	-	5
Female	209	1	-	-	-	-	-	1	1	-	-	2
White, both sexes	324	2	-	-	-	-	-	1	1	2	-	5
Male	134	1	-	-	-	-	-	-	-	2	-	3
Female	190	1	-	-	-	-	-	1	1	-	-	2
Black, both sexes	36	-	1	-	-	-	-	-	-	-	-	2
Male	17	-	1	-	-	-	-	-	-	-	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	118	1	-	-	-	-	-	-	-	1	2	-
Male	58	1	-	-	-	-	-	-	-	1	1	-
Female	60	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	110	1	-	-	-	-	-	-	-	1	2	-
Male	54	1	-	-	-	-	-	-	-	1	1	-
Female	56	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Oregon												
All races, both sexes	112	2	1	-	-	-	-	1	-	1	1	2
Male	43	1	1	-	-	-	-	-	-	1	1	2
Female	69	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	109	2	1	-	-	-	-	1	-	1	1	2
Male	40	1	1	-	-	-	-	-	-	1	1	2
Female	69	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	564	1	1	-	-	-	2	1	1	4	4	12
Male	269	1	1	-	-	-	1	-	1	4	3	7
Female	295	-	-	-	-	-	1	1	-	-	1	5
White, both sexes	515	1	1	-	-	-	-	1	1	3	4	11
Male	245	1	1	-	-	-	-	-	1	3	3	7
Female	270	-	-	-	-	-	-	1	-	-	1	4
Black, both sexes	47	-	-	-	-	-	2	-	-	1	-	1
Male	23	-	-	-	-	-	1	-	-	1	-	-
Female	24	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Rhode Island												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	161	-	-	1	-	-	-	-	1	-	4	6
Male	77	-	-	1	-	-	-	-	1	-	1	5
Female	84	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	109	-	-	-	-	-	-	-	-	-	2	3
Male	54	-	-	-	-	-	-	-	-	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	51	-	-	1	-	-	-	-	1	-	2	3
Male	23	-	-	1	-	-	-	-	1	-	1	3
Female	28	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
South Dakota												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	175	-	-	-	-	-	-	-	-	1	5	4
Male	73	-	-	-	-	-	-	-	-	-	1	3
Female	102	-	-	-	-	-	-	-	-	1	4	1
White, both sexes	149	-	-	-	-	-	-	-	-	1	4	2
Male	60	-	-	-	-	-	-	-	-	-	1	2
Female	89	-	-	-	-	-	-	-	-	1	3	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Texas												
All races, both sexes	403	6	-	-	-	1	-	-	1	2	5	6
Male	174	4	-	-	-	1	-	-	1	1	1	6
Female	229	2	-	-	-	-	-	-	-	1	4	-
White, both sexes	357	5	-	-	-	1	-	-	1	2	4	4
Male	152	3	-	-	-	1	-	-	1	1	1	4
Female	205	2	-	-	-	-	-	-	-	1	3	-
Black, both sexes	42	1	-	-	-	-	-	-	-	-	1	2
Male	21	1	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Vermont												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	228	-	1	-	-	-	-	1	-	4	1	6
Male	109	-	1	-	-	-	-	1	-	2	1	4
Female	119	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	177	-	-	-	-	-	-	1	-	4	1	4
Male	86	-	-	-	-	-	-	1	-	2	1	2
Female	91	-	-	-	-	-	-	-	-	2	-	2
Black, both sexes	50	-	1	-	-	-	-	-	-	-	-	2
Male	23	-	1	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Washington												
All races, both sexes	157	-	-	-	-	1	-	-	-	-	-	2
Male	72	-	-	-	-	-	-	-	-	-	-	1
Female	85	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	150	-	-	-	-	1	-	-	-	-	-	2
Male	68	-	-	-	-	-	-	-	-	-	-	1
Female	82	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	-	3
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	34	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of digestive system (K90,K92)												
Wisconsin												
All races, both sexes	238	-	-	-	-	-	-	-	-	1	-	5
Male	103	-	-	-	-	-	-	-	-	1	-	4
Female	135	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	225	-	-	-	-	-	-	-	-	1	-	5
Male	99	-	-	-	-	-	-	-	-	1	-	4
Female	126	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
United States												
All races, both sexes	235	323	343	478	638	946	1,348	1,496	1,191	538	97	-
Male	161	213	203	284	347	507	607	562	362	129	18	-
Female	74	110	140	194	291	439	741	934	829	409	79	-
White, both sexes	174	260	270	378	527	802	1,217	1,383	1,110	489	85	-
Male	126	171	160	220	291	436	547	525	349	111	14	-
Female	48	89	110	158	236	366	670	858	761	378	71	-
Black, both sexes	57	56	64	87	97	124	113	92	69	40	9	-
Male	34	37	36	56	48	59	50	30	8	11	2	-
Female	23	19	28	31	49	65	63	62	61	29	7	-
American Indian 1/, both sexes	2	5	4	1	4	2	7	4	3	1	1	-
Male	-	4	3	1	2	2	2	-	-	1	1	-
Female	2	1	1	-	2	-	5	4	3	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	5	12	10	18	11	17	9	8	2	-
Male	1	1	4	7	6	10	8	7	5	6	1	-
Female	1	1	1	5	4	8	3	10	4	2	1	-
Alabama												
All races, both sexes	7	10	5	17	21	20	32	26	17	8	-	-
Male	4	8	3	9	10	9	14	8	5	-	-	-
Female	3	2	2	8	11	11	18	18	12	8	-	-
White, both sexes	2	9	3	10	15	16	27	20	14	6	-	-
Male	1	8	2	4	8	8	11	8	5	-	-	-
Female	1	1	1	6	7	8	16	12	9	6	-	-
Black, both sexes	5	1	2	7	6	4	5	6	3	2	-	-
Male	3	-	1	5	2	1	3	-	-	-	-	-
Female	2	1	1	2	4	3	2	6	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Alaska												
All races, both sexes	-	-	1	1	-	3	1	2	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	1	1	-	2	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	2	8	3	12	17	24	28	19	8	1	-
Male	2	1	5	3	6	8	15	8	7	3	-	-
Female	1	1	3	-	6	9	9	20	12	5	1	-
White, both sexes	2	2	7	3	10	15	20	27	18	7	1	-
Male	2	1	5	3	5	7	13	8	7	2	-	-
Female	-	1	2	-	5	8	7	19	11	5	1	-
Black, both sexes	-	-	1	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	3	1	1	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	1	-	-	-	-	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Arkansas												
All races, both sexes	4	7	7	9	9	16	16	22	18	8	3	-
Male	3	2	5	7	3	7	9	9	5	1	-	-
Female	1	5	2	2	6	9	7	13	13	7	3	-
White, both sexes	4	5	5	7	8	13	16	21	16	6	2	-
Male	3	2	3	5	3	5	9	9	5	-	-	-
Female	1	3	2	2	5	8	7	12	11	6	2	-
Black, both sexes	-	2	2	2	1	3	-	1	2	2	1	-
Male	-	-	2	2	-	2	-	-	-	1	-	-
Female	-	2	-	-	1	1	-	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	7	12	18	24	32	57	64	47	24	4	-
Male	-	4	10	11	17	19	30	28	18	7	1	-
Female	3	3	2	7	7	13	27	36	29	17	3	-
White, both sexes	3	4	11	12	22	25	55	61	44	22	4	-
Male	-	2	9	6	15	16	29	27	17	6	1	-
Female	3	2	2	6	7	9	26	34	27	16	3	-
Black, both sexes	-	2	1	1	1	4	2	-	1	-	-	-
Male	-	1	1	1	1	2	1	-	-	-	-	-
Female	-	1	-	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	4	1	3	-	3	1	2	-	-
Male	-	1	-	3	1	1	-	1	1	1	-	-
Female	-	-	-	1	-	2	-	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Colorado												
All races, both sexes	2	2	6	4	5	19	15	28	17	13	-	-
Male	1	1	2	2	3	15	8	12	5	4	-	-
Female	1	1	4	2	2	4	7	16	12	9	-	-
White, both sexes	2	2	4	4	5	19	15	28	16	13	-	-
Male	1	1	2	2	3	15	8	12	5	4	-	-
Female	1	1	2	2	2	4	7	16	11	9	-	-
Black, both sexes	-	-	2	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	7	3	2	6	20	24	33	15	22	3	-
Male	4	5	2	-	3	13	8	16	5	4	-	-
Female	-	2	1	2	3	7	16	17	10	18	3	-
White, both sexes	4	7	3	2	6	19	23	33	15	22	3	-
Male	4	5	2	-	3	12	7	16	5	4	-	-
Female	-	2	1	2	3	7	16	17	10	18	3	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Delaware												
All races, both sexes	-	1	4	2	2	3	5	8	2	1	-	-
Male	-	1	4	1	-	-	1	5	-	-	-	-
Female	-	-	-	1	2	3	4	3	2	1	-	-
White, both sexes	-	1	4	1	1	2	2	6	2	1	-	-
Male	-	1	4	-	-	-	1	4	-	-	-	-
Female	-	-	-	1	1	2	1	2	2	1	-	-
Black, both sexes	-	-	-	1	1	1	3	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	3	-	-	1	1	2	2	1	2	-	-
Male	-	2	-	-	1	-	1	2	-	1	-	-
Female	-	1	-	-	-	1	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	3	-	-	1	1	2	2	-	2	-	-
Male	-	2	-	-	1	-	1	2	-	1	-	-
Female	-	1	-	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Florida												
All races, both sexes	12	16	16	18	26	48	55	85	44	21	7	-
Male	9	11	9	13	9	19	25	37	16	7	3	-
Female	3	5	7	5	17	29	30	48	28	14	4	-
White, both sexes	9	16	11	16	22	42	53	78	43	21	6	-
Male	7	11	7	11	8	17	25	32	16	7	2	-
Female	2	5	4	5	14	25	28	46	27	14	4	-
Black, both sexes	3	-	5	2	4	5	2	6	1	-	1	-
Male	2	-	2	2	1	2	-	4	-	-	1	-
Female	1	-	3	-	3	3	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Georgia												
All races, both sexes	17	18	11	19	26	31	37	35	23	8	1	-
Male	7	12	4	11	15	15	16	15	3	1	-	-
Female	10	6	7	8	11	16	21	20	20	7	1	-
White, both sexes	7	12	9	9	19	22	28	29	17	7	1	-
Male	2	8	3	4	11	12	13	15	3	1	-	-
Female	5	4	6	5	8	10	15	14	14	6	1	-
Black, both sexes	10	6	2	10	7	8	9	6	6	1	-	-
Male	5	4	1	7	4	3	3	-	-	-	-	-
Female	5	2	1	3	3	5	6	6	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Hawaii												
All races, both sexes	-	3	3	4	5	3	9	6	7	4	1	-
Male	-	2	2	3	3	2	6	1	4	2	1	-
Female	-	1	1	1	2	1	3	5	3	2	-	-
White, both sexes	-	3	1	1	-	2	2	2	-	2	-	-
Male	-	2	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	5	1	7	4	6	2	1	-
Male	-	-	1	2	3	1	5	-	4	2	1	-
Female	-	-	1	1	2	-	2	4	2	-	-	-
Idaho												
All races, both sexes	1	1	1	2	3	-	8	9	7	4	-	-
Male	1	1	-	-	1	-	5	1	2	1	-	-
Female	-	-	1	2	2	-	3	8	5	3	-	-
White, both sexes	1	1	1	2	2	-	8	9	7	4	-	-
Male	1	1	-	-	-	-	5	1	2	1	-	-
Female	-	-	1	2	2	-	3	8	5	3	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Illinois												
All races, both sexes	9	19	12	23	29	49	56	63	53	30	3	-
Male	6	15	5	16	16	28	26	26	12	4	1	-
Female	3	4	7	7	13	21	30	37	41	26	2	-
White, both sexes	8	14	10	19	25	43	53	57	49	27	1	-
Male	5	11	4	12	13	25	26	21	12	3	-	-
Female	3	3	6	7	12	18	27	36	37	24	1	-
Black, both sexes	1	5	2	3	2	4	3	4	4	3	1	-
Male	1	4	1	3	2	1	-	3	-	1	1	-
Female	-	1	1	-	-	3	3	1	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	2	-	2	-	-	1	-
Male	-	-	-	1	1	2	-	2	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
Indiana												
All races, both sexes	7	11	7	14	14	25	39	28	31	8	2	-
Male	5	8	5	9	6	14	21	10	8	3	1	-
Female	2	3	2	5	8	11	18	18	23	5	1	-
White, both sexes	4	11	6	14	14	21	38	26	30	8	1	-
Male	2	8	4	9	6	12	21	10	8	3	1	-
Female	2	3	2	5	8	9	17	16	22	5	-	-
Black, both sexes	3	-	1	-	-	2	1	2	1	-	1	-
Male	3	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Iowa												
All races, both sexes	1	1	-	4	3	9	15	23	22	5	1	-
Male	1	1	-	4	-	4	8	7	4	1	-	-
Female	-	-	-	-	3	5	7	16	18	4	1	-
White, both sexes	1	1	-	4	3	9	15	22	22	5	1	-
Male	1	1	-	4	-	4	8	6	4	1	-	-
Female	-	-	-	-	3	5	7	16	18	4	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	2	4	3	1	7	15	24	16	7	1	-
Male	3	-	3	1	1	5	5	8	5	4	1	-
Female	-	2	1	2	-	2	10	16	11	3	-	-
White, both sexes	3	2	4	3	-	6	15	23	15	7	1	-
Male	3	-	3	1	-	4	5	8	5	4	1	-
Female	-	2	1	2	-	2	10	15	10	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Kentucky												
All races, both sexes	2	6	12	10	8	14	15	21	20	5	1	-
Male	1	6	3	4	5	7	6	14	6	1	-	-
Female	1	-	9	6	3	7	9	7	14	4	1	-
White, both sexes	1	6	12	10	7	13	15	19	19	5	1	-
Male	-	6	3	4	4	6	6	13	6	1	-	-
Female	1	-	9	6	3	7	9	6	13	4	1	-
Black, both sexes	1	-	-	-	1	1	-	2	1	-	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	8	7	18	18	16	24	18	13	7	-	-
Male	6	6	5	10	11	8	11	4	4	2	-	-
Female	2	2	2	8	7	8	13	14	9	5	-	-
White, both sexes	6	5	4	12	14	12	20	16	10	4	-	-
Male	4	3	3	7	8	6	8	3	3	1	-	-
Female	2	2	1	5	6	6	12	13	7	3	-	-
Black, both sexes	2	3	3	6	4	4	4	2	3	3	-	-
Male	2	3	2	3	3	2	3	1	1	1	-	-
Female	-	-	1	3	1	2	1	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Maine												
All races, both sexes	1	3	2	6	4	5	10	13	8	2	-	-
Male	1	2	1	2	1	3	5	4	-	1	-	-
Female	-	1	1	4	3	2	5	9	8	1	-	-
White, both sexes	1	3	2	6	4	5	10	13	8	2	-	-
Male	1	2	1	2	1	3	5	4	-	1	-	-
Female	-	1	1	4	3	2	5	9	8	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	7	14	14	8	24	28	31	20	8	3	-
Male	2	5	9	11	4	16	14	11	9	-	-	-
Female	-	2	5	3	4	8	14	20	11	8	3	-
White, both sexes	1	4	9	9	6	17	22	26	16	6	2	-
Male	1	3	5	7	3	11	11	10	8	-	-	-
Female	-	1	4	2	3	6	11	16	8	6	2	-
Black, both sexes	1	3	5	5	2	6	6	5	4	2	1	-
Male	1	2	4	4	1	5	3	1	1	-	-	-
Female	-	1	1	1	1	1	3	4	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Massachusetts												
All races, both sexes	9	5	11	17	20	38	51	56	48	26	2	-
Male	7	2	6	10	11	24	21	20	16	7	1	-
Female	2	3	5	7	9	14	30	36	32	19	1	-
White, both sexes	8	5	10	15	18	36	51	56	48	24	2	-
Male	6	2	6	8	11	22	21	20	16	5	1	-
Female	2	3	4	7	7	14	30	36	32	19	1	-
Black, both sexes	1	-	1	2	1	2	-	-	-	-	-	-
Male	1	-	-	2	-	2	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5	10	13	12	21	28	55	60	52	23	6	-
Male	4	5	5	6	13	15	23	30	17	4	1	-
Female	1	5	8	6	8	13	32	30	35	19	5	-
White, both sexes	4	9	12	9	14	24	50	59	50	21	6	-
Male	3	4	4	3	10	13	21	30	17	3	1	-
Female	1	5	8	6	4	11	29	29	33	18	5	-
Black, both sexes	1	1	1	3	7	4	4	1	2	2	-	-
Male	1	1	1	3	3	2	2	-	-	1	-	-
Female	-	-	-	-	4	2	2	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Minnesota												
All races, both sexes	2	3	8	8	6	10	27	27	35	12	4	-
Male	2	2	7	6	3	5	15	14	9	4	-	-
Female	-	1	1	2	3	5	12	13	26	8	4	-
White, both sexes	2	2	4	8	6	9	26	27	35	12	4	-
Male	2	1	3	6	3	5	14	14	9	4	-	-
Female	-	1	1	2	3	4	12	13	26	8	4	-
Black, both sexes	-	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	-	-	-	-	-	-	-	-	-
Male	-	1	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	7	6	10	13	20	19	18	13	7	-	-
Male	2	5	3	3	5	10	9	6	2	3	-	-
Female	1	2	3	7	8	10	10	12	11	4	-	-
White, both sexes	1	5	3	4	8	14	13	14	8	4	-	-
Male	1	4	1	1	2	8	6	4	1	1	-	-
Female	-	1	2	3	6	6	7	10	7	3	-	-
Black, both sexes	2	2	3	6	5	6	6	4	5	3	-	-
Male	1	1	2	2	3	2	3	2	1	2	-	-
Female	1	1	1	4	2	4	3	2	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Missouri												
All races, both sexes	7	5	8	12	20	36	29	37	34	19	4	-
Male	5	2	2	4	15	22	13	13	10	6	-	-
Female	2	3	6	8	5	14	16	24	24	13	4	-
White, both sexes	7	4	5	10	18	32	28	36	33	19	4	-
Male	5	2	2	4	13	19	13	12	10	6	-	-
Female	2	2	3	6	5	13	15	24	23	13	4	-
Black, both sexes	-	1	3	2	1	4	1	1	1	-	-	-
Male	-	-	-	-	1	3	-	1	-	-	-	-
Female	-	1	3	2	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	1	1	3	2	2	4	6	7	4	-	-
Male	3	-	1	2	1	1	-	3	3	-	-	-
Female	-	1	-	1	1	1	4	3	4	4	-	-
White, both sexes	3	1	1	3	2	2	3	6	7	4	-	-
Male	3	-	1	2	1	1	-	3	3	-	-	-
Female	-	1	-	1	1	1	3	3	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Nebraska												
All races, both sexes	3	1	4	-	4	7	7	18	9	6	-	-
Male	3	1	2	-	2	3	5	7	2	1	-	-
Female	-	-	2	-	2	4	2	11	7	5	-	-
White, both sexes	3	1	4	-	4	5	7	18	9	6	-	-
Male	3	1	2	-	2	1	5	7	2	1	-	-
Female	-	-	2	-	2	4	2	11	7	5	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	6	7	8	8	10	12	10	5	1	1	-
Male	2	4	5	7	4	4	8	3	1	1	1	-
Female	2	2	2	1	4	6	4	7	4	-	-	-
White, both sexes	3	5	7	7	7	10	12	10	5	1	1	-
Male	2	3	5	6	4	4	8	3	1	1	1	-
Female	1	2	2	1	3	6	4	7	4	-	-	-
Black, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
New Hampshire												
All races, both sexes	1	2	-	3	4	3	2	10	6	5	-	-
Male	1	2	-	2	1	1	1	6	1	-	-	-
Female	-	-	-	1	3	2	1	4	5	5	-	-
White, both sexes	1	2	-	3	4	3	2	10	6	5	-	-
Male	1	2	-	2	1	1	1	6	1	-	-	-
Female	-	-	-	1	3	2	1	4	5	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	4	8	18	28	28	33	58	43	17	3	-
Male	8	4	5	12	15	13	14	20	16	5	-	-
Female	3	-	3	6	13	15	19	38	27	12	3	-
White, both sexes	7	4	7	18	24	20	28	54	41	16	3	-
Male	5	4	4	12	14	11	11	20	16	5	-	-
Female	2	-	3	6	10	9	17	34	25	11	3	-
Black, both sexes	4	-	1	-	4	7	4	3	1	1	-	-
Male	3	-	1	-	1	1	2	-	-	-	-	-
Female	1	-	-	-	3	6	2	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
New Mexico												
All races, both sexes	6	3	2	2	7	4	12	6	4	2	-	-
Male	5	3	2	2	4	-	6	1	2	1	-	-
Female	1	-	-	-	3	4	6	5	2	1	-	-
White, both sexes	5	2	2	1	6	4	12	5	4	2	-	-
Male	5	2	2	1	4	-	6	1	2	1	-	-
Female	-	-	-	-	2	4	6	4	2	1	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	10	14	14	25	33	51	63	89	80	36	7	-
Male	8	7	8	10	16	28	27	36	30	3	1	-
Female	2	7	6	15	17	23	36	53	50	33	6	-
White, both sexes	7	12	10	18	21	42	59	78	75	34	6	-
Male	7	7	6	8	10	24	25	31	29	3	1	-
Female	-	5	4	10	11	18	34	47	46	31	5	-
Black, both sexes	3	2	3	7	12	6	4	7	5	2	1	-
Male	1	-	1	2	6	1	2	2	1	-	-	-
Female	2	2	2	5	6	5	2	5	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	3	-	4	-	-	-	-
Male	-	-	1	-	-	3	-	3	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
North Carolina												
All races, both sexes	5	14	17	19	24	35	40	42	36	10	1	-
Male	1	9	12	16	14	18	18	8	12	3	-	-
Female	4	5	5	3	10	17	22	34	24	7	1	-
White, both sexes	3	8	11	10	17	25	28	29	33	8	1	-
Male	1	5	8	10	12	12	12	5	12	2	-	-
Female	2	3	3	-	5	13	16	24	21	6	1	-
Black, both sexes	1	5	6	9	7	9	10	12	3	2	-	-
Male	-	4	4	6	2	6	6	3	-	1	-	-
Female	1	1	2	3	5	3	4	9	3	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	1	1	5	2	-	-
Male	-	-	-	-	-	1	1	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	3	2	-	-
White, both sexes	-	-	-	-	-	1	1	1	5	2	-	-
Male	-	-	-	-	-	1	1	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Ohio												
All races, both sexes	13	15	15	18	34	39	68	62	57	23	2	-
Male	6	11	8	11	17	19	26	21	17	6	-	-
Female	7	4	7	7	17	20	42	41	40	17	2	-
White, both sexes	9	15	13	13	31	32	66	60	53	19	2	-
Male	5	11	7	7	16	14	24	21	17	6	-	-
Female	4	4	6	6	15	18	42	39	36	13	2	-
Black, both sexes	4	-	2	5	3	7	2	2	4	4	-	-
Male	1	-	1	4	1	5	2	-	-	-	-	-
Female	3	-	1	1	2	2	-	2	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	9	8	8	11	10	28	18	15	3	2	-
Male	2	8	3	5	5	7	9	10	5	1	-	-
Female	-	1	5	3	6	3	19	8	10	2	2	-
White, both sexes	2	7	7	8	10	10	27	17	13	3	2	-
Male	2	6	2	5	5	7	9	10	4	1	-	-
Female	-	1	5	3	5	3	18	7	9	2	2	-
Black, both sexes	-	2	-	-	-	-	1	-	2	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Oregon												
All races, both sexes	2	2	7	6	3	11	19	22	18	12	2	-
Male	-	-	6	2	2	3	9	3	11	1	-	-
Female	2	2	1	4	1	8	10	19	7	11	2	-
White, both sexes	2	2	5	6	3	11	18	22	18	12	2	-
Male	-	-	4	2	2	3	8	3	11	1	-	-
Female	2	2	1	4	1	8	10	19	7	11	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	12	16	20	28	39	81	112	100	87	30	13	-
Male	9	11	17	20	25	43	51	34	33	6	2	-
Female	3	5	3	8	14	38	61	66	54	24	11	-
White, both sexes	8	14	18	24	37	73	99	98	81	29	12	-
Male	7	10	16	18	23	40	45	33	30	5	2	-
Female	1	4	2	6	14	33	54	65	51	24	10	-
Black, both sexes	3	2	2	3	2	8	13	2	6	1	1	-
Male	1	1	1	2	2	3	6	1	3	1	-	-
Female	2	1	1	1	-	5	7	1	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Rhode Island												
All races, both sexes	-	1	-	4	5	7	6	12	8	4	2	-
Male	-	1	-	2	2	4	3	2	1	-	1	-
Female	-	-	-	2	3	3	3	10	7	4	1	-
White, both sexes	-	1	-	4	5	6	6	12	8	4	2	-
Male	-	1	-	2	2	3	3	2	1	-	1	-
Female	-	-	-	2	3	3	3	10	7	4	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	8	8	10	11	14	17	29	24	19	7	2	-
Male	8	6	3	7	10	7	14	8	4	2	-	-
Female	-	2	7	4	4	10	15	16	15	5	2	-
White, both sexes	6	4	6	7	10	6	23	21	14	6	1	-
Male	6	3	2	5	7	4	11	8	4	2	-	-
Female	-	1	4	2	3	2	12	13	10	4	1	-
Black, both sexes	2	4	3	4	4	11	6	3	5	1	1	-
Male	2	3	1	2	3	3	3	-	-	-	-	-
Female	-	1	2	2	1	8	3	3	5	1	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
South Dakota												
All races, both sexes	1	-	-	-	1	4	5	4	3	6	-	-
Male	1	-	-	-	-	2	3	3	-	2	-	-
Female	-	-	-	-	1	2	2	1	3	4	-	-
White, both sexes	1	-	-	-	1	4	5	4	3	6	-	-
Male	1	-	-	-	-	2	3	3	-	2	-	-
Female	-	-	-	-	1	2	2	1	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	11	12	10	13	19	25	34	26	12	1	-
Male	1	4	9	6	9	12	10	9	7	2	-	-
Female	1	7	3	4	4	7	15	25	19	10	1	-
White, both sexes	1	9	10	7	9	18	24	30	26	8	-	-
Male	-	2	7	4	7	11	10	8	7	1	-	-
Female	1	7	3	3	2	7	14	22	19	7	-	-
Black, both sexes	1	2	2	3	4	1	1	4	-	4	1	-
Male	1	2	2	2	2	1	-	1	-	1	-	-
Female	-	-	-	1	2	-	1	3	-	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Texas												
All races, both sexes	10	18	19	24	37	49	74	63	61	22	5	-
Male	6	9	11	9	24	31	30	16	15	9	-	-
Female	4	9	8	15	13	18	44	47	46	13	5	-
White, both sexes	8	15	16	21	29	44	65	60	59	18	5	-
Male	5	8	9	8	19	27	27	16	15	7	-	-
Female	3	7	7	13	10	17	38	44	44	11	5	-
Black, both sexes	2	2	3	2	8	5	9	3	2	2	-	-
Male	1	1	2	1	5	4	3	-	-	1	-	-
Female	1	1	1	1	3	1	6	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	1	-	-	-	-	-	1	-	-
Utah												
All races, both sexes	-	1	1	-	6	3	13	9	10	1	-	-
Male	-	-	-	-	3	2	6	3	-	-	-	-
Female	-	1	1	-	3	1	7	6	10	1	-	-
White, both sexes	-	1	1	-	6	2	12	9	10	1	-	-
Male	-	-	-	-	3	1	5	3	-	-	-	-
Female	-	1	1	-	3	1	7	6	10	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Vermont												
All races, both sexes	-	-	-	-	1	1	2	4	6	3	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	-	1	2	5	3	-	-
White, both sexes	-	-	-	-	1	1	2	4	6	3	-	-
Male	-	-	-	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	1	-	1	2	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	12	15	9	13	22	22	48	38	25	9	2	-
Male	9	11	5	7	10	12	17	19	7	2	1	-
Female	3	4	4	6	12	10	31	19	18	7	1	-
White, both sexes	8	10	4	9	17	19	37	30	23	8	2	-
Male	7	7	2	6	8	12	13	15	7	2	1	-
Female	1	3	2	3	9	7	24	15	16	6	1	-
Black, both sexes	4	5	5	3	5	3	11	8	2	1	-	-
Male	2	4	3	1	2	-	4	4	-	-	-	-
Female	2	1	2	2	3	3	7	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Washington												
All races, both sexes	4	5	1	9	11	18	27	26	35	14	4	-
Male	2	4	1	5	3	9	16	12	12	6	1	-
Female	2	1	-	4	8	9	11	14	23	8	3	-
White, both sexes	4	4	1	9	11	17	27	24	33	14	3	-
Male	2	3	1	5	3	8	16	11	12	6	-	-
Female	2	1	-	4	8	9	11	13	21	8	3	-
Black, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	2	1	1	8	3	9	11	11	9	5	-	-
Male	2	1	-	5	3	4	4	4	2	1	-	-
Female	-	-	1	3	-	5	7	7	7	4	-	-
White, both sexes	2	1	1	8	3	9	11	10	9	4	-	-
Male	2	1	-	5	3	4	4	3	2	1	-	-
Female	-	-	1	3	-	5	7	7	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Wisconsin												
All races, both sexes	12	12	5	10	23	21	37	55	33	21	3	-
Male	8	8	3	6	15	12	12	21	6	6	1	-
Female	4	4	2	4	8	9	25	34	27	15	2	-
White, both sexes	10	9	4	10	22	20	35	55	31	20	3	-
Male	7	6	3	6	15	12	11	21	6	6	1	-
Female	3	3	1	4	7	8	24	34	25	14	2	-
Black, both sexes	2	1	1	-	1	1	2	-	1	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	1	-	1	1	1	-	1	1	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	-	-	1	1	-	-	2	7	2	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	5	2	1	-	-
White, both sexes	-	-	1	1	-	-	2	7	2	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
United States												
All races, both sexes	4,234	-	3	3	1	2	11	20	26	45	77	125
Male	1,628	-	1	2	-	-	5	6	15	20	42	55
Female	2,606	-	2	1	1	2	6	14	11	25	35	70
White, both sexes	3,239	-	1	3	1	1	6	9	15	29	51	90
Male	1,272	-	1	2	-	-	2	3	8	15	28	45
Female	1,967	-	-	1	1	1	4	6	7	14	23	45
Black, both sexes	894	-	1	-	-	1	5	10	9	14	20	31
Male	314	-	-	-	-	-	3	3	5	4	10	10
Female	580	-	1	-	-	1	2	7	4	10	10	21
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	1	-	2	3
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	77	-	1	-	-	-	-	1	1	2	4	1
Male	37	-	-	-	-	-	-	-	1	1	3	-
Female	40	-	1	-	-	-	-	1	-	1	1	1
Alabama												
All races, both sexes	80	-	-	-	-	-	1	-	1	-	2	4
Male	32	-	-	-	-	-	-	-	1	-	-	1
Female	48	-	-	-	-	-	1	-	-	-	2	3
White, both sexes	46	-	-	-	-	-	1	-	1	-	1	2
Male	21	-	-	-	-	-	-	-	1	-	-	1
Female	25	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	75	-	-	-	-	-	-	-	-	1	2	1
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	44	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	66	-	-	-	-	-	-	-	-	1	2	1
Male	29	-	-	-	-	-	-	-	-	-	2	1
Female	37	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Arkansas												
All races, both sexes	50	-	-	-	-	-	-	-	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	36	-	-	-	-	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	290	-	-	-	1	-	-	3	2	3	10	20
Male	126	-	-	-	-	-	-	1	1	3	6	9
Female	164	-	-	-	1	-	-	2	1	-	4	11
White, both sexes	232	-	-	-	1	-	-	1	1	3	6	16
Male	105	-	-	-	-	-	-	1	1	3	3	7
Female	127	-	-	-	1	-	-	-	-	-	3	9
Black, both sexes	34	-	-	-	-	-	-	2	1	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	24	-	-	-	-	-	-	2	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	3	1
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	12	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Colorado												
All races, both sexes	55	-	-	-	-	-	1	-	-	-	2	2
Male	18	-	-	-	-	-	-	-	-	-	2	-
Female	37	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	51	-	-	-	-	-	1	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	62	-	1	-	-	-	-	-	-	-	-	-
Male	25	-	1	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	-	1	-	-	-	-	-	-	-	-	-
Male	21	-	1	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Florida												
All races, both sexes	216	-	1	-	-	-	-	1	2	1	5	4
Male	88	-	-	-	-	-	-	-	1	-	3	2
Female	128	-	1	-	-	-	-	1	1	1	2	2
White, both sexes	167	-	-	-	-	-	-	1	1	-	3	3
Male	75	-	-	-	-	-	-	-	-	-	2	2
Female	92	-	-	-	-	-	-	1	1	-	1	1
Black, both sexes	49	-	1	-	-	-	-	-	1	1	2	1
Male	13	-	-	-	-	-	-	-	1	-	1	-
Female	36	-	1	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	149	-	-	-	-	-	1	-	1	1	-	4
Male	51	-	-	-	-	-	1	-	-	-	-	2
Female	98	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	91	-	-	-	-	-	-	-	-	1	-	2
Male	37	-	-	-	-	-	-	-	-	-	-	2
Female	54	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	57	-	-	-	-	-	1	-	1	-	-	2
Male	14	-	-	-	-	-	1	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Hawaii												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Illinois												
All races, both sexes	206	-	-	-	-	-	1	-	-	5	4	4
Male	69	-	-	-	-	-	1	-	-	-	1	2
Female	137	-	-	-	-	-	-	-	-	5	3	2
White, both sexes	155	-	-	-	-	-	-	-	-	2	3	3
Male	52	-	-	-	-	-	-	-	-	-	1	1
Female	103	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	48	-	-	-	-	-	1	-	-	3	1	1
Male	17	-	-	-	-	-	1	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	105	-	-	1	-	-	-	-	1	-	4	4
Male	43	-	-	-	-	-	-	-	-	-	1	1
Female	62	-	-	1	-	-	-	-	1	-	3	3
White, both sexes	88	-	-	1	-	-	-	-	1	-	4	4
Male	38	-	-	-	-	-	-	-	-	-	1	1
Female	50	-	-	1	-	-	-	-	1	-	3	3
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Iowa												
All races, both sexes	37	-	-	-	-	-	-	-	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Kansas												
All races, both sexes	37	-	-	1	-	-	-	1	-	-	-	2
Male	16	-	-	1	-	-	-	1	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	34	-	-	1	-	-	-	-	-	-	-	1
Male	14	-	-	1	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Kentucky												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	2	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	110	-	-	-	-	-	2	1	2	2	2	2
Male	46	-	-	-	-	-	-	-	1	2	2	1
Female	64	-	-	-	-	-	2	1	1	-	-	1
White, both sexes	63	-	-	-	-	-	1	-	1	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	1	-
Female	39	-	-	-	-	-	1	-	1	-	-	1
Black, both sexes	47	-	-	-	-	-	1	1	1	1	1	1
Male	22	-	-	-	-	-	-	-	1	1	1	1
Female	25	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Maine												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	88	-	-	-	-	-	-	-	-	-	3	2
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	2	1
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Massachusetts												
All races, both sexes	84	-	-	-	-	-	-	1	-	1	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	76	-	-	-	-	-	-	1	-	-	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	184	-	-	-	-	1	-	-	1	4	4	4
Male	76	-	-	-	-	-	-	-	-	1	4	3
Female	108	-	-	-	-	1	-	-	1	3	-	1
White, both sexes	136	-	-	-	-	-	-	-	-	1	4	4
Male	56	-	-	-	-	-	-	-	-	-	4	3
Female	80	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	46	-	-	-	-	1	-	-	1	3	-	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	1	-	-	1	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Minnesota												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	28	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	1	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	28	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	69	-	-	-	-	-	-	2	-	-	4	2
Male	28	-	-	-	-	-	-	1	-	-	3	2
Female	41	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	40	-	-	-	-	-	-	1	-	-	3	-
Male	15	-	-	-	-	-	-	1	-	-	3	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Missouri												
All races, both sexes	76	-	-	-	-	-	-	1	2	1	2	6
Male	33	-	-	-	-	-	-	-	1	1	1	3
Female	43	-	-	-	-	-	-	1	1	-	1	3
White, both sexes	60	-	-	-	-	-	-	-	-	1	1	4
Male	26	-	-	-	-	-	-	-	-	1	1	2
Female	34	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	15	-	-	-	-	-	-	1	1	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Nebraska												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	1	-	1	-	3	-
Male	17	-	-	-	-	-	1	-	-	-	2	-
Female	20	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	30	-	-	-	-	-	1	-	1	-	2	-
Male	14	-	-	-	-	-	1	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
New Hampshire												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	175	-	-	-	-	-	1	-	-	2	1	5
Male	67	-	-	-	-	-	-	-	-	1	1	2
Female	108	-	-	-	-	-	1	-	-	1	-	3
White, both sexes	135	-	-	-	-	-	-	-	-	-	1	1
Male	52	-	-	-	-	-	-	-	-	-	1	-
Female	83	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	39	-	-	-	-	-	1	-	-	2	-	4
Male	15	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	1	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
New Mexico												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	262	-	-	-	-	-	-	1	1	2	1	1
Male	104	-	-	-	-	-	-	-	1	-	-	1
Female	158	-	-	-	-	-	-	1	-	2	1	-
White, both sexes	218	-	-	-	-	-	-	-	-	2	1	1
Male	87	-	-	-	-	-	-	-	-	-	-	1
Female	131	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	42	-	-	-	-	-	-	1	1	-	-	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
North Carolina												
All races, both sexes	168	-	-	1	-	-	-	2	1	2	3	4
Male	71	-	-	1	-	-	-	1	1	2	1	-
Female	97	-	-	-	-	-	-	1	-	-	2	4
White, both sexes	114	-	-	1	-	-	-	-	1	2	2	1
Male	54	-	-	1	-	-	-	-	1	2	1	-
Female	60	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	52	-	-	-	-	-	-	2	-	-	1	3
Male	17	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Ohio												
All races, both sexes	169	-	-	-	-	-	1	1	1	1	2	6
Male	56	-	-	-	-	-	1	-	1	-	2	3
Female	113	-	-	-	-	-	-	1	-	1	-	3
White, both sexes	138	-	-	-	-	-	-	-	-	1	1	4
Male	46	-	-	-	-	-	-	-	-	-	1	2
Female	92	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	31	-	-	-	-	-	1	1	1	-	1	2
Male	10	-	-	-	-	-	1	-	1	-	1	1
Female	21	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	65	-	-	-	-	-	-	-	-	1	5	3
Male	18	-	-	-	-	-	-	-	-	1	1	-
Female	47	-	-	-	-	-	-	-	-	-	4	3
White, both sexes	54	-	-	-	-	-	-	-	-	1	3	2
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	38	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Oregon												
All races, both sexes	52	-	-	-	-	-	1	-	2	1	1	1
Male	19	-	-	-	-	-	1	-	1	1	-	-
Female	33	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	51	-	-	-	-	-	1	-	2	1	-	1
Male	19	-	-	-	-	-	1	-	1	1	-	-
Female	32	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	207	-	-	-	-	-	-	1	1	1	1	6
Male	79	-	-	-	-	-	-	1	1	1	-	4
Female	128	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	176	-	-	-	-	-	-	1	-	-	-	5
Male	66	-	-	-	-	-	-	1	-	-	-	4
Female	110	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	30	-	-	-	-	-	-	-	1	1	1	1
Male	13	-	-	-	-	-	-	-	1	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Rhode Island												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	75	-	-	-	-	-	-	-	-	3	-	1
Male	22	-	-	-	-	-	-	-	-	2	-	-
Female	53	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	2	-	1
Male	16	-	-	-	-	-	-	-	-	2	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	33	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
South Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	96	-	-	-	-	-	-	-	1	2	1	2
Male	37	-	-	-	-	-	-	-	1	1	-	1
Female	59	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	72	-	-	-	-	-	-	-	1	2	1	1
Male	28	-	-	-	-	-	-	-	1	1	-	1
Female	44	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Texas												
All races, both sexes	305	-	-	-	-	1	1	3	3	4	7	13
Male	131	-	-	-	-	-	-	1	3	1	4	4
Female	174	-	-	-	-	1	1	2	-	3	3	9
White, both sexes	221	-	-	-	-	1	1	3	3	3	7	10
Male	92	-	-	-	-	-	-	1	3	1	4	3
Female	129	-	-	-	-	1	1	2	-	2	3	7
Black, both sexes	80	-	-	-	-	-	-	-	-	1	-	3
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	21	-	1	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Vermont												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	130	-	-	-	-	-	-	-	1	-	2	2
Male	50	-	-	-	-	-	-	-	-	-	2	1
Female	80	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	76	-	-	-	-	-	-	-	1	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	54	-	-	-	-	-	-	-	-	-	2	1
Male	23	-	-	-	-	-	-	-	-	-	2	-
Female	31	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Washington												
All races, both sexes	72	-	-	-	-	-	-	2	-	1	1	4
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	2	-	1	1	3
White, both sexes	67	-	-	-	-	-	-	2	-	1	1	3
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	2	-	1	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Wisconsin												
All races, both sexes	71	-	-	-	-	-	-	-	1	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	65	-	-	-	-	-	-	-	1	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
United States												
All races, both sexes	156	203	211	256	359	533	714	674	522	241	52	-
Male	87	109	100	121	157	222	264	227	133	50	12	-
Female	69	94	111	135	202	311	450	447	389	191	40	-
White, both sexes	103	142	143	193	264	400	573	554	430	197	34	-
Male	60	79	71	95	113	173	215	196	117	42	7	-
Female	43	63	72	98	151	227	358	358	313	155	27	-
Black, both sexes	46	57	63	58	85	119	125	108	83	42	17	-
Male	23	29	27	23	38	46	42	27	11	8	5	-
Female	23	28	36	35	47	73	83	81	72	34	12	-
American Indian 1/, both sexes	3	1	1	-	3	1	4	1	2	1	1	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	2	1	1	-	2	-	4	1	2	1	1	-
Asian or Pacific Islander 2/, both sexes	4	3	4	5	7	13	12	11	7	1	-	-
Male	3	1	2	3	5	2	7	4	5	-	-	-
Female	1	2	2	2	2	11	5	7	2	1	-	-
Alabama												
All races, both sexes	2	1	3	5	8	10	16	8	11	5	3	-
Male	1	1	2	3	4	3	9	2	3	-	2	-
Female	1	-	1	2	4	7	7	6	8	5	1	-
White, both sexes	-	1	2	2	5	5	10	5	7	2	2	-
Male	-	1	1	2	2	2	4	2	3	-	2	-
Female	-	-	1	-	3	3	6	3	4	2	-	-
Black, both sexes	2	-	1	3	2	5	6	3	4	3	1	-
Male	1	-	1	1	1	1	5	-	-	-	-	-
Female	1	-	-	2	1	4	1	3	4	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Alaska												
All races, both sexes	1	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	5	5	7	6	11	9	10	10	2	1	-
Male	3	1	2	4	2	5	5	4	2	-	-	-
Female	2	4	3	3	4	6	4	6	8	2	1	-
White, both sexes	3	4	5	7	4	11	8	10	9	1	-	-
Male	2	1	2	4	1	5	5	4	2	-	-	-
Female	1	3	3	3	3	6	3	6	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	1	-	1	-	1	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	1	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Arkansas												
All races, both sexes	1	5	3	-	4	9	8	11	4	1	1	-
Male	-	2	2	-	-	3	2	1	-	-	-	-
Female	1	3	1	-	4	6	6	10	4	1	1	-
White, both sexes	1	3	1	-	2	7	4	10	4	1	-	-
Male	-	1	1	-	-	2	1	-	-	-	-	-
Female	1	2	-	-	2	5	3	10	4	1	-	-
Black, both sexes	-	2	2	-	2	2	4	1	-	-	1	-
Male	-	1	1	-	-	1	1	1	-	-	-	-
Female	-	1	1	-	2	1	3	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	18	19	13	18	23	32	43	40	30	14	1	-
Male	9	9	7	11	14	13	14	13	13	3	-	-
Female	9	10	6	7	9	19	29	27	17	11	1	-
White, both sexes	14	13	12	15	17	25	35	34	25	14	-	-
Male	7	7	7	9	10	12	14	10	11	3	-	-
Female	7	6	5	6	7	13	21	24	14	11	-	-
Black, both sexes	3	5	1	1	3	3	6	3	1	-	1	-
Male	2	1	-	1	2	-	-	1	-	-	-	-
Female	1	4	1	-	1	3	6	2	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	3	4	2	3	4	-	-	-
Male	-	1	-	1	2	1	-	2	2	-	-	-
Female	-	-	-	1	1	3	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Colorado												
All races, both sexes	4	2	2	-	5	9	8	6	6	7	1	-
Male	-	2	-	-	2	2	2	5	1	1	1	-
Female	4	-	2	-	3	7	6	1	5	6	-	-
White, both sexes	4	2	2	-	4	9	7	6	6	7	1	-
Male	-	2	-	-	2	2	1	5	1	1	1	-
Female	4	-	2	-	2	7	6	1	5	6	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	4	3	3	7	4	12	10	13	4	1	-
Male	-	4	1	3	1	2	5	3	2	3	-	-
Female	-	-	2	-	6	2	7	7	11	1	1	-
White, both sexes	-	3	2	2	7	3	10	10	13	4	1	-
Male	-	3	1	2	1	1	4	3	2	3	-	-
Female	-	-	1	-	6	2	6	7	11	1	1	-
Black, both sexes	-	1	1	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Delaware												
All races, both sexes	-	-	1	4	-	1	2	4	1	-	-	-
Male	-	-	1	2	-	-	-	2	-	-	-	-
Female	-	-	-	2	-	1	2	2	1	-	-	-
White, both sexes	-	-	1	4	-	1	2	4	1	-	-	-
Male	-	-	1	2	-	-	-	2	-	-	-	-
Female	-	-	-	2	-	1	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	3	2	1	2	3	-	5	2	-	-
Male	-	1	-	-	-	2	-	-	1	2	-	-
Female	1	-	3	2	1	-	3	-	4	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	3	2	1	2	3	-	5	2	-	-
Male	-	1	-	-	-	2	-	-	1	2	-	-
Female	1	-	3	2	1	-	3	-	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Florida												
All races, both sexes	10	12	13	12	16	23	48	28	26	12	2	-
Male	6	6	7	6	8	10	15	12	8	4	-	-
Female	4	6	6	6	8	13	33	16	18	8	2	-
White, both sexes	5	8	8	9	13	20	39	25	20	12	-	-
Male	4	5	4	6	5	10	14	12	7	4	-	-
Female	1	3	4	3	8	10	25	13	13	8	-	-
Black, both sexes	5	4	5	3	3	3	9	3	6	-	2	-
Male	2	1	3	-	3	-	1	-	1	-	-	-
Female	3	3	2	3	-	3	8	3	5	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	13	9	7	8	21	12	43	18	5	4	-
Male	1	9	7	1	1	9	3	15	-	1	1	-
Female	1	4	2	6	7	12	9	28	18	4	3	-
White, both sexes	-	6	6	4	5	15	9	28	11	2	2	-
Male	-	6	4	-	-	9	2	14	-	-	-	-
Female	-	-	2	4	5	6	7	14	11	2	2	-
Black, both sexes	2	7	3	3	3	6	3	14	7	3	2	-
Male	1	3	3	1	1	-	1	1	-	1	1	-
Female	1	4	-	2	2	6	2	13	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Hawaii												
All races, both sexes	3	2	3	-	2	5	8	3	1	-	1	-
Male	2	1	2	-	-	-	4	2	1	-	-	-
Female	1	1	1	-	2	5	4	1	-	-	1	-
White, both sexes	-	2	-	-	1	1	2	-	-	-	1	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	2	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	3	-	1	4	6	3	1	-	-	-
Male	2	-	2	-	-	-	4	2	1	-	-	-
Female	1	-	1	-	1	4	2	1	-	-	-	-
Idaho												
All races, both sexes	1	2	-	1	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	2	-	1	-	-	2	1	-	-	-	-
White, both sexes	1	2	-	1	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	2	-	1	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Illinois												
All races, both sexes	8	15	10	6	24	20	34	32	31	10	2	-
Male	4	4	2	4	8	8	12	14	9	-	-	-
Female	4	11	8	2	16	12	22	18	22	10	2	-
White, both sexes	5	9	5	6	17	13	26	29	29	6	2	-
Male	3	1	1	4	5	4	10	13	9	-	-	-
Female	2	8	4	2	12	9	16	16	20	6	2	-
Black, both sexes	3	5	5	-	7	6	8	3	2	3	-	-
Male	1	3	1	-	3	4	2	1	-	-	-	-
Female	2	2	4	-	4	2	6	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-
Indiana												
All races, both sexes	3	5	6	15	9	12	14	16	8	7	-	-
Male	2	2	4	5	4	3	8	6	6	1	-	-
Female	1	3	2	10	5	9	6	10	2	6	-	-
White, both sexes	2	4	4	13	6	9	12	14	8	6	-	-
Male	2	1	2	5	2	3	8	6	6	1	-	-
Female	-	3	2	8	4	6	4	8	2	5	-	-
Black, both sexes	1	1	2	2	3	3	2	2	-	1	-	-
Male	-	1	2	-	2	-	-	-	-	-	-	-
Female	1	-	-	2	1	3	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Iowa												
All races, both sexes	1	-	2	1	6	4	5	8	6	2	1	-
Male	1	-	2	-	3	1	3	2	4	1	-	-
Female	-	-	-	1	3	3	2	6	2	1	1	-
White, both sexes	1	-	2	1	6	4	5	8	6	2	1	-
Male	1	-	2	-	3	1	3	2	4	1	-	-
Female	-	-	-	1	3	3	2	6	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	4	2	-	3	3	4	8	3	4	-	-
Male	2	3	-	-	-	1	1	3	1	2	-	-
Female	-	1	2	-	3	2	3	5	2	2	-	-
White, both sexes	2	4	2	-	3	3	4	7	3	4	-	-
Male	2	3	-	-	-	1	1	3	1	2	-	-
Female	-	1	2	-	3	2	3	4	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Kentucky												
All races, both sexes	1	1	2	4	4	7	10	16	3	-	2	-
Male	-	-	1	3	2	3	3	3	-	-	1	-
Female	1	1	1	1	2	4	7	13	3	-	1	-
White, both sexes	1	1	2	3	4	4	8	14	3	-	2	-
Male	-	-	1	2	2	2	1	3	-	-	1	-
Female	1	1	1	1	2	2	7	11	3	-	1	-
Black, both sexes	-	-	-	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Louisiana												
All races, both sexes	5	5	8	8	8	12	19	12	13	8	1	-
Male	2	4	4	4	4	5	8	2	5	1	1	-
Female	3	1	4	4	4	7	11	10	8	7	-	-
White, both sexes	3	2	4	3	3	8	12	8	9	6	-	-
Male	1	2	2	2	1	4	5	2	3	-	-	-
Female	2	-	2	1	2	4	7	6	6	6	-	-
Black, both sexes	2	3	4	5	5	4	7	4	4	2	1	-
Male	1	2	2	2	3	1	3	-	2	1	1	-
Female	1	1	2	3	2	3	4	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Maine												
All races, both sexes	-	1	2	-	1	3	5	2	-	3	-	-
Male	-	1	1	-	1	1	-	-	-	1	-	-
Female	-	-	1	-	-	2	5	2	-	2	-	-
White, both sexes	-	1	2	-	1	3	5	2	-	3	-	-
Male	-	1	1	-	1	1	-	-	-	1	-	-
Female	-	-	1	-	-	2	5	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	2	7	8	12	9	12	15	7	4	-	-
Male	3	1	3	1	5	4	3	5	2	1	-	-
Female	4	1	4	7	7	5	9	10	5	3	-	-
White, both sexes	3	1	3	4	8	4	6	10	5	2	-	-
Male	3	-	2	-	4	2	3	3	2	1	-	-
Female	-	1	1	4	4	2	3	7	3	1	-	-
Black, both sexes	4	1	4	4	4	5	6	5	2	2	-	-
Male	-	1	1	1	1	2	-	2	-	-	-	-
Female	4	-	3	3	3	3	6	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Massachusetts												
All races, both sexes	-	5	1	1	2	7	21	19	10	13	1	-
Male	-	3	-	1	-	-	5	6	1	5	-	-
Female	-	2	1	-	2	7	16	13	9	8	1	-
White, both sexes	-	5	1	1	2	6	16	19	9	13	1	-
Male	-	3	-	1	-	-	5	6	1	5	-	-
Female	-	2	1	-	2	6	11	13	8	8	1	-
Black, both sexes	-	-	-	-	-	1	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	2	9	13	13	15	27	37	26	21	5	2	-
Male	1	6	9	6	6	12	14	9	4	1	-	-
Female	1	3	4	7	9	15	23	17	17	4	2	-
White, both sexes	1	6	9	10	9	17	31	20	17	5	2	-
Male	-	3	7	4	4	7	11	8	4	1	-	-
Female	1	3	2	6	5	10	20	12	13	4	2	-
Black, both sexes	1	3	4	3	5	10	6	5	4	-	-	-
Male	1	3	2	2	1	5	3	1	-	-	-	-
Female	-	-	2	1	4	5	3	4	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Minnesota												
All races, both sexes	1	-	1	2	6	6	8	5	6	5	1	-
Male	1	-	-	-	2	3	3	1	-	3	1	-
Female	-	-	1	2	4	3	5	4	6	2	-	-
White, both sexes	1	-	1	2	6	5	8	5	6	5	1	-
Male	1	-	-	-	2	2	3	1	-	3	1	-
Female	-	-	1	2	4	3	5	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	5	1	3	5	10	12	10	8	3	1	-
Male	2	4	1	1	3	3	3	2	2	-	1	-
Female	1	1	-	2	2	7	9	8	6	3	-	-
White, both sexes	1	2	1	2	3	1	6	5	2	1	-	-
Male	1	2	1	-	2	1	-	2	1	-	-	-
Female	-	-	-	2	1	-	6	3	1	1	-	-
Black, both sexes	2	3	-	1	2	9	5	5	6	2	1	-
Male	1	2	-	1	1	2	2	-	1	-	1	-
Female	1	1	-	-	1	7	3	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Missouri												
All races, both sexes	4	-	3	8	4	9	18	7	7	3	1	-
Male	2	-	1	6	2	2	6	5	2	1	-	-
Female	2	-	2	2	2	7	12	2	5	2	1	-
White, both sexes	2	-	2	5	3	8	18	7	6	3	-	-
Male	1	-	-	4	1	2	6	5	2	1	-	-
Female	1	-	2	1	2	6	12	2	4	2	-	-
Black, both sexes	2	-	1	3	1	1	-	-	1	-	1	-
Male	1	-	1	2	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	1	-	2	3	4	3	-	1	-	-
Male	-	-	1	-	2	1	1	1	-	-	-	-
Female	1	-	-	-	-	2	3	2	-	1	-	-
White, both sexes	1	-	1	-	2	3	4	3	-	1	-	-
Male	-	-	1	-	2	1	1	1	-	-	-	-
Female	1	-	-	-	-	2	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Nebraska												
All races, both sexes	-	-	1	1	2	5	2	6	5	1	-	-
Male	-	-	1	-	1	1	-	1	3	-	-	-
Female	-	-	-	1	1	4	2	5	2	1	-	-
White, both sexes	-	-	1	-	2	5	2	6	4	1	-	-
Male	-	-	1	-	1	1	-	1	3	-	-	-
Female	-	-	-	-	1	4	2	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	6	3	2	3	4	6	7	-	-	-
Male	1	-	2	1	1	-	1	4	4	-	-	-
Female	-	-	4	2	1	3	3	2	3	-	-	-
White, both sexes	1	-	3	2	2	3	3	6	6	-	-	-
Male	1	-	2	-	1	-	1	4	3	-	-	-
Female	-	-	1	2	1	3	2	2	3	-	-	-
Black, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
New Hampshire												
All races, both sexes	-	1	2	1	-	2	5	3	-	1	-	-
Male	-	1	1	-	-	1	2	1	-	-	-	-
Female	-	-	1	1	-	1	3	2	-	1	-	-
White, both sexes	-	1	2	1	-	2	5	3	-	1	-	-
Male	-	1	1	-	-	1	2	1	-	-	-	-
Female	-	-	1	1	-	1	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	4	1	10	14	27	29	36	20	14	1	-
Male	4	1	1	4	9	12	12	12	7	1	-	-
Female	6	3	-	6	5	15	17	24	13	13	1	-
White, both sexes	7	2	1	9	10	20	21	32	18	12	1	-
Male	3	1	1	4	5	10	9	11	6	1	-	-
Female	4	1	-	5	5	10	12	21	12	11	1	-
Black, both sexes	3	2	-	1	4	7	8	3	2	2	-	-
Male	1	-	-	-	4	2	3	1	1	-	-	-
Female	2	2	-	1	-	5	5	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
New Mexico												
All races, both sexes	1	-	1	4	3	2	3	1	-	1	-	-
Male	-	-	-	2	3	1	1	-	-	-	-	-
Female	1	-	1	2	-	1	2	1	-	1	-	-
White, both sexes	1	-	1	4	2	2	2	1	-	1	-	-
Male	-	-	-	2	2	1	1	-	-	-	-	-
Female	1	-	1	2	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	15	7	19	19	36	45	50	42	15	4	-
Male	3	7	2	6	10	18	21	18	13	4	-	-
Female	1	8	5	13	9	18	24	32	29	11	4	-
White, both sexes	2	11	5	15	15	31	42	42	38	10	3	-
Male	2	5	1	5	7	16	20	17	11	2	-	-
Female	-	6	4	10	8	15	22	25	27	8	3	-
Black, both sexes	2	4	2	4	4	4	3	8	3	5	1	-
Male	1	2	1	1	3	2	1	1	1	2	-	-
Female	1	2	1	3	1	2	2	7	2	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
North Carolina												
All races, both sexes	9	6	11	12	14	25	26	22	20	9	1	-
Male	6	4	6	4	6	15	8	11	3	2	-	-
Female	3	2	5	8	8	10	18	11	17	7	1	-
White, both sexes	8	3	8	8	10	19	19	13	12	7	-	-
Male	5	2	5	3	5	12	5	7	3	2	-	-
Female	3	1	3	5	5	7	14	6	9	5	-	-
Black, both sexes	1	2	3	4	3	6	7	9	8	2	1	-
Male	1	2	1	1	1	3	3	4	-	-	-	-
Female	-	-	2	3	2	3	4	5	8	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	2	1	1	-	1	1	3	1	-	-
Male	-	1	2	1	1	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
White, both sexes	-	1	2	1	1	-	1	1	3	1	-	-
Male	-	1	2	1	1	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Ohio												
All races, both sexes	4	9	8	6	15	22	21	33	30	9	-	-
Male	2	6	3	2	4	9	10	7	5	1	-	-
Female	2	3	5	4	11	13	11	26	25	8	-	-
White, both sexes	1	7	7	5	13	20	20	27	24	8	-	-
Male	-	4	3	2	4	8	10	6	5	1	-	-
Female	1	3	4	3	9	12	10	21	19	7	-	-
Black, both sexes	3	2	1	1	2	2	1	6	6	1	-	-
Male	2	2	-	-	-	1	-	1	-	-	-	-
Female	1	-	1	1	2	1	1	5	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	4	3	3	4	8	13	10	6	2	-	-
Male	1	3	-	1	1	1	2	4	3	-	-	-
Female	2	1	3	2	3	7	11	6	3	2	-	-
White, both sexes	3	4	2	2	2	7	13	9	5	1	-	-
Male	1	3	-	1	1	1	2	3	2	-	-	-
Female	2	1	2	1	1	6	11	6	3	1	-	-
Black, both sexes	-	-	1	1	2	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	1	2	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Oregon												
All races, both sexes	-	-	-	5	3	7	10	8	8	3	2	-
Male	-	-	-	1	1	4	3	4	1	1	1	-
Female	-	-	-	4	2	3	7	4	7	2	1	-
White, both sexes	-	-	-	5	3	7	10	8	8	3	2	-
Male	-	-	-	1	1	4	3	4	1	1	1	-
Female	-	-	-	4	2	3	7	4	7	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	7	7	8	11	11	39	33	39	26	14	2	-
Male	6	4	3	5	7	20	15	8	2	2	-	-
Female	1	3	5	6	4	19	18	31	24	12	2	-
White, both sexes	5	7	6	10	9	32	31	34	23	12	1	-
Male	4	4	2	5	5	16	14	7	2	2	-	-
Female	1	3	4	5	4	16	17	27	21	10	1	-
Black, both sexes	2	-	2	1	2	6	2	5	3	2	1	-
Male	2	-	1	-	2	4	1	1	-	-	-	-
Female	-	-	1	1	-	2	1	4	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Rhode Island												
All races, both sexes	-	-	-	2	1	2	2	1	2	1	1	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	2	1	-	2	1	1	-
White, both sexes	-	-	-	2	1	2	2	1	2	1	1	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	2	1	-	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	2	6	9	4	11	14	9	8	5	1	-
Male	2	1	3	4	1	2	4	1	1	1	-	-
Female	-	1	3	5	3	9	10	8	7	4	1	-
White, both sexes	2	1	3	5	1	3	8	4	8	3	1	-
Male	2	1	1	3	-	1	3	1	1	1	-	-
Female	-	-	2	2	1	2	5	3	7	2	1	-
Black, both sexes	-	1	3	4	3	8	6	5	-	2	-	-
Male	-	-	2	1	1	1	1	-	-	-	-	-
Female	-	1	1	3	2	7	5	5	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
South Dakota												
All races, both sexes	-	-	1	-	1	1	-	4	2	2	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	2	2	-	-
White, both sexes	-	-	1	-	1	1	-	4	2	2	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	5	5	8	12	10	12	15	15	4	-	-
Male	2	3	3	5	3	5	4	6	2	1	-	-
Female	2	2	2	3	9	5	8	9	13	3	-	-
White, both sexes	2	2	3	7	10	7	9	11	12	4	-	-
Male	1	2	2	4	3	4	3	4	1	1	-	-
Female	1	-	1	3	7	3	6	7	11	3	-	-
Black, both sexes	2	3	2	1	2	3	3	4	3	-	-	-
Male	1	1	1	1	-	1	1	2	1	-	-	-
Female	1	2	1	-	2	2	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Texas												
All races, both sexes	17	16	11	15	36	33	56	40	33	12	4	-
Male	14	8	5	11	16	13	25	12	10	2	2	-
Female	3	8	6	4	20	20	31	28	23	10	2	-
White, both sexes	11	13	7	9	25	25	42	28	26	7	-	-
Male	9	7	3	7	11	9	19	6	8	1	-	-
Female	2	6	4	2	14	16	23	22	18	6	-	-
Black, both sexes	5	3	4	5	11	8	13	11	7	5	4	-
Male	4	1	2	3	5	4	6	6	2	1	2	-
Female	1	2	2	2	6	4	7	5	5	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	1	2	-	1	2	2	3	3	1	2	-	-
Male	1	1	-	1	1	1	-	1	-	-	-	-
Female	-	1	-	-	1	1	3	2	1	2	-	-
White, both sexes	1	2	-	-	2	2	3	3	1	2	-	-
Male	1	1	-	-	1	1	-	1	-	-	-	-
Female	-	1	-	-	1	1	3	2	1	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Vermont												
All races, both sexes	-	-	1	-	1	-	2	4	1	1	-	-
Male	-	-	1	-	1	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-
White, both sexes	-	-	1	-	1	-	2	4	1	1	-	-
Male	-	-	1	-	1	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	5	10	6	15	17	24	12	18	10	5	-
Male	1	2	3	4	7	10	8	5	4	2	1	-
Female	2	3	7	2	8	7	16	7	14	8	4	-
White, both sexes	3	1	3	3	6	9	15	7	15	7	5	-
Male	1	1	1	2	4	5	3	2	4	2	1	-
Female	2	-	2	1	2	4	12	5	11	5	4	-
Black, both sexes	-	4	7	3	9	8	9	5	3	3	-	-
Male	-	1	2	2	3	5	5	3	-	-	-	-
Female	-	3	5	1	6	3	4	2	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Washington												
All races, both sexes	1	6	6	4	5	2	10	11	12	6	1	-
Male	1	3	3	2	3	-	6	4	2	-	-	-
Female	-	3	3	2	2	2	4	7	10	6	1	-
White, both sexes	1	6	5	4	5	2	10	9	11	6	1	-
Male	1	3	3	2	3	-	6	4	2	-	-	-
Female	-	3	2	2	2	2	4	5	9	6	1	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	2	-	3	3	4	5	8	5	4	2	-	-
Male	-	-	-	2	1	3	5	1	1	-	-	-
Female	2	-	3	1	3	2	3	4	3	2	-	-
White, both sexes	2	-	3	3	3	3	8	5	3	2	-	-
Male	-	-	-	2	1	1	5	1	1	-	-	-
Female	2	-	3	1	2	2	3	4	2	2	-	-
Black, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Wisconsin												
All races, both sexes	1	2	1	4	7	12	13	10	10	6	2	-
Male	1	-	-	4	4	7	4	4	3	2	-	-
Female	-	2	1	-	3	5	9	6	7	4	2	-
White, both sexes	1	2	1	4	7	11	12	10	8	5	2	-
Male	1	-	-	4	4	6	3	4	3	1	-	-
Female	-	2	1	-	3	5	9	6	5	4	2	-
Black, both sexes	-	-	-	-	-	1	1	-	2	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	3	2	1	-	1	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	1	-
White, both sexes	-	-	-	-	-	-	3	2	1	-	1	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
United States												
All races, both sexes	1,702	-	1	-	-	-	4	8	9	20	40	68
Male	745	-	-	-	-	-	3	2	9	11	25	31
Female	957	-	1	-	-	-	1	6	-	9	15	37
White, both sexes	1,461	-	-	-	-	-	2	6	4	17	32	52
Male	641	-	-	-	-	-	1	1	4	9	19	28
Female	820	-	-	-	-	-	1	5	-	8	13	24
Black, both sexes	205	-	-	-	-	-	2	2	4	2	4	16
Male	83	-	-	-	-	-	2	1	4	1	2	3
Female	122	-	-	-	-	-	-	1	-	1	2	13
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	1	-	-	-	-	-	-	1	3	-
Male	17	-	-	-	-	-	-	-	-	1	3	-
Female	11	-	1	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	33	-	-	-	-	-	-	-	-	1	2	1
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	21	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	1	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Arkansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	154	-	-	-	-	-	-	1	1	3	8	12
Male	76	-	-	-	-	-	-	1	1	3	5	7
Female	78	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	133	-	-	-	-	-	-	1	1	3	6	9
Male	67	-	-	-	-	-	-	1	1	3	3	5
Female	66	-	-	-	-	-	-	-	-	-	3	4
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Colorado												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Florida												
All races, both sexes	95	-	-	-	-	-	-	-	1	-	2	4
Male	44	-	-	-	-	-	-	-	1	-	1	2
Female	51	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	78	-	-	-	-	-	-	-	-	-	2	3
Male	40	-	-	-	-	-	-	-	-	-	1	2
Female	38	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	17	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	40	-	-	-	-	-	1	-	-	-	-	1
Male	19	-	-	-	-	-	1	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Hawaii												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Illinois												
All races, both sexes	90	-	-	-	-	-	-	-	-	1	3	3
Male	38	-	-	-	-	-	-	-	-	-	1	1
Female	52	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	79	-	-	-	-	-	-	-	-	1	2	3
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	45	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Kentucky												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	32	-	-	-	-	-	-	-	1	1	1	1
Male	18	-	-	-	-	-	-	-	1	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Maine												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Massachusetts												
All races, both sexes	35	-	-	-	-	-	-	1	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	35	-	-	-	-	-	-	1	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	70	-	-	-	-	-	-	-	-	3	3	1
Male	32	-	-	-	-	-	-	-	-	1	3	1
Female	38	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	1	3	1
Male	25	-	-	-	-	-	-	-	-	-	3	1
Female	33	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	2	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Minnesota												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Missouri												
All races, both sexes	27	-	-	-	-	-	-	-	1	-	-	1
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Nebraska												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	2	-
Male	8	-	-	-	-	-	1	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	1	-	-	-	2	-
Male	8	-	-	-	-	-	1	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
New Hampshire												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
New Mexico												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	102	-	-	-	-	-	-	-	-	1	1	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	89	-	-	-	-	-	-	-	-	1	1	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
North Carolina												
All races, both sexes	64	-	-	-	-	-	-	1	1	1	2	2
Male	34	-	-	-	-	-	-	1	1	1	1	-
Female	30	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	47	-	-	-	-	-	-	-	1	1	2	-
Male	26	-	-	-	-	-	-	-	1	1	1	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	16	-	-	-	-	-	-	1	-	-	-	2
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Ohio												
All races, both sexes	72	-	-	-	-	-	1	1	1	-	1	4
Male	30	-	-	-	-	-	1	-	1	-	1	2
Female	42	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	62	-	-	-	-	-	-	-	-	-	1	4
Male	26	-	-	-	-	-	-	-	-	-	1	2
Female	36	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	10	-	-	-	-	-	1	1	1	-	-	-
Male	4	-	-	-	-	-	1	-	1	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Oregon												
All races, both sexes	27	-	-	-	-	-	-	-	1	1	-	-
Male	12	-	-	-	-	-	-	-	1	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	1	1	-	-
Male	12	-	-	-	-	-	-	-	1	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	81	-	-	-	-	-	-	-	1	-	-	3
Male	36	-	-	-	-	-	-	-	1	-	-	3
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	3
Male	31	-	-	-	-	-	-	-	-	-	-	3
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Rhode Island												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
South Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Texas												
All races, both sexes	137	-	-	-	-	-	1	2	1	1	4	9
Male	67	-	-	-	-	-	-	-	1	-	2	2
Female	70	-	-	-	-	-	1	2	-	1	2	7
White, both sexes	108	-	-	-	-	-	1	2	1	1	4	7
Male	55	-	-	-	-	-	-	-	1	-	2	2
Female	53	-	-	-	-	-	1	2	-	1	2	5
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	13	-	1	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Washington												
All races, both sexes	36	-	-	-	-	-	-	2	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	2	-	1	-	1
White, both sexes	35	-	-	-	-	-	-	2	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	2	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Wisconsin												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
United States												
All races, both sexes	90	108	89	112	159	221	257	246	177	81	12	-
Male	56	59	52	52	75	100	103	106	47	12	2	-
Female	34	49	37	60	84	121	154	140	130	69	10	-
White, both sexes	66	87	64	96	141	191	233	222	160	79	9	-
Male	41	49	43	45	65	88	92	97	45	12	2	-
Female	25	38	21	51	76	103	141	125	115	67	7	-
Black, both sexes	18	19	23	14	17	26	20	20	14	1	3	-
Male	12	10	8	5	9	10	9	6	1	-	-	-
Female	6	9	15	9	8	16	11	14	13	1	3	-
American Indian 1/, both sexes	3	1	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	2	2	1	3	3	4	3	1	-	-
Male	2	-	1	2	1	1	2	3	1	-	-	-
Female	1	1	1	-	-	2	1	1	2	1	-	-
Alabama												
All races, both sexes	1	-	1	2	3	2	6	3	4	1	1	-
Male	1	-	1	2	2	1	3	2	2	-	1	-
Female	-	-	-	-	1	1	3	1	2	1	-	-
White, both sexes	-	-	1	2	3	2	5	3	3	1	1	-
Male	-	-	1	2	2	1	2	2	2	-	1	-
Female	-	-	-	-	1	1	3	1	1	1	-	-
Black, both sexes	1	-	-	-	-	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Alaska												
All races, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	3	2	5	2	4	5	4	2	-	-	-
Male	1	1	1	2	-	1	2	1	-	-	-	-
Female	1	2	1	3	2	3	3	3	2	-	-	-
White, both sexes	-	2	2	5	2	4	5	4	2	-	-	-
Male	-	1	1	2	-	1	2	1	-	-	-	-
Female	-	1	1	3	2	3	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Arkansas												
All races, both sexes	-	2	2	-	-	5	3	5	1	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	1	1	-	-	3	3	5	1	-	-	-
White, both sexes	-	2	1	-	-	5	3	5	1	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	1	-	-	-	3	3	5	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	17	9	7	12	13	20	15	15	14	7	-	-
Male	8	3	4	7	8	8	8	6	5	2	-	-
Female	9	6	3	5	5	12	7	9	9	5	-	-
White, both sexes	14	7	6	10	12	18	14	13	12	7	-	-
Male	7	3	4	5	7	8	8	5	5	2	-	-
Female	7	4	2	5	5	10	6	8	7	5	-	-
Black, both sexes	2	2	1	1	-	-	1	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	2	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	1	2	-	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Colorado												
All races, both sexes	3	-	-	-	2	2	4	3	2	3	-	-
Male	-	-	-	-	1	1	1	2	1	-	-	-
Female	3	-	-	-	1	1	3	1	1	3	-	-
White, both sexes	3	-	-	-	2	2	3	3	2	3	-	-
Male	-	-	-	-	1	1	-	2	1	-	-	-
Female	3	-	-	-	1	1	3	1	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	4	2	2	3	1	4	8	6	2	1	-
Male	-	4	1	2	-	-	2	3	1	1	-	-
Female	-	-	1	-	3	1	2	5	5	1	1	-
White, both sexes	-	3	1	2	3	1	3	8	6	2	1	-
Male	-	3	1	2	-	-	2	3	1	1	-	-
Female	-	-	-	-	3	1	1	5	5	1	1	-
Black, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Delaware												
All races, both sexes	-	-	1	1	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Florida												
All races, both sexes	7	9	9	5	5	14	14	12	9	4	-	-
Male	5	5	5	3	3	5	5	6	3	-	-	-
Female	2	4	4	2	2	9	9	6	6	4	-	-
White, both sexes	5	6	7	4	5	12	12	11	7	4	-	-
Male	4	4	4	3	3	5	5	6	3	-	-	-
Female	1	2	3	1	2	7	7	5	4	4	-	-
Black, both sexes	2	3	2	1	-	2	2	1	2	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	2	1	1	-	2	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	4	3	3	2	9	1	11	2	-	2	-
Male	1	2	3	-	1	5	1	5	-	-	-	-
Female	-	2	-	3	1	4	-	6	2	-	2	-
White, both sexes	-	2	1	3	1	8	1	10	1	-	1	-
Male	-	2	1	-	-	5	1	5	-	-	-	-
Female	-	-	-	3	1	3	-	5	1	-	1	-
Black, both sexes	1	2	2	-	1	1	-	1	1	-	1	-
Male	1	-	2	-	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	1	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Hawaii												
All races, both sexes	2	1	2	-	1	-	3	2	-	-	1	-
Male	1	-	1	-	-	-	1	2	-	-	-	-
Female	1	1	1	-	1	-	2	-	-	-	1	-
White, both sexes	-	1	-	-	1	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	-	-	-	2	2	-	-	-	-
Male	1	-	1	-	-	-	1	2	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	2	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	2	1	-	-	-	-
White, both sexes	-	2	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Illinois												
All races, both sexes	4	5	4	4	11	5	17	12	17	4	-	-
Male	2	1	1	3	5	2	8	8	6	-	-	-
Female	2	4	3	1	6	3	9	4	11	4	-	-
White, both sexes	4	3	3	4	10	4	16	12	15	2	-	-
Male	2	-	1	3	4	1	7	8	6	-	-	-
Female	2	3	2	1	6	3	9	4	9	2	-	-
Black, both sexes	-	2	1	-	1	1	1	-	2	1	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	2	2	1	8	3	6	6	8	3	4	-	-
Male	1	1	1	2	1	2	3	2	3	1	-	-
Female	1	1	-	6	2	4	3	6	-	3	-	-
White, both sexes	1	2	-	7	2	4	6	8	3	4	-	-
Male	1	1	-	2	1	2	3	2	3	1	-	-
Female	-	1	-	5	1	2	3	6	-	3	-	-
Black, both sexes	1	-	1	1	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Iowa												
All races, both sexes	1	-	2	1	5	4	4	4	1	1	-	-
Male	1	-	2	-	2	1	3	-	-	-	-	-
Female	-	-	-	1	3	3	1	4	1	1	-	-
White, both sexes	1	-	2	1	5	4	4	4	1	1	-	-
Male	1	-	2	-	2	1	3	-	-	-	-	-
Female	-	-	-	1	3	3	1	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	4	2	-	1	1	2	3	-	3	-	-
Male	1	3	-	-	-	1	-	3	-	1	-	-
Female	-	1	2	-	1	-	2	-	-	2	-	-
White, both sexes	1	4	2	-	1	1	2	3	-	3	-	-
Male	1	3	-	-	-	1	-	3	-	1	-	-
Female	-	1	2	-	1	-	2	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Kentucky												
All races, both sexes	1	-	1	2	2	1	2	4	1	-	1	-
Male	-	-	1	1	1	-	-	1	-	-	1	-
Female	1	-	-	1	1	1	2	3	1	-	-	-
White, both sexes	1	-	1	2	2	1	2	4	1	-	1	-
Male	-	-	1	1	1	-	-	1	-	-	1	-
Female	1	-	-	1	1	1	2	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	2	1	3	4	7	3	3	2	-	-
Male	1	2	2	1	1	1	4	-	3	-	-	-
Female	-	-	-	-	2	3	3	3	-	2	-	-
White, both sexes	-	2	2	-	2	3	5	3	2	2	-	-
Male	-	2	2	-	1	1	2	-	2	-	-	-
Female	-	-	-	-	1	2	3	3	-	2	-	-
Black, both sexes	1	-	-	1	1	1	2	-	1	-	-	-
Male	1	-	-	1	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Maine												
All races, both sexes	-	-	-	-	-	3	2	1	-	2	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	2	1	-	2	-	-
White, both sexes	-	-	-	-	-	3	2	1	-	2	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	2	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	2	3	1	5	3	4	5	3	-	-	-
Male	3	1	1	-	4	1	1	2	1	-	-	-
Female	1	1	2	1	1	2	3	3	2	-	-	-
White, both sexes	3	1	2	1	5	2	3	5	3	-	-	-
Male	3	-	1	-	4	1	1	2	1	-	-	-
Female	-	1	1	1	1	1	2	3	2	-	-	-
Black, both sexes	1	1	1	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Massachusetts												
All races, both sexes	-	3	1	1	2	2	8	4	4	7	-	-
Male	-	2	-	1	-	-	3	1	-	1	-	-
Female	-	1	1	-	2	2	5	3	4	6	-	-
White, both sexes	-	3	1	1	2	2	8	4	4	7	-	-
Male	-	2	-	1	-	-	3	1	-	1	-	-
Female	-	1	1	-	2	2	5	3	4	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	5	5	5	4	7	17	6	12	2	-	-
Male	-	4	5	2	3	4	5	2	1	1	-	-
Female	-	1	-	3	1	3	12	4	11	1	-	-
White, both sexes	-	3	5	4	4	4	15	6	10	2	-	-
Male	-	2	5	1	3	3	3	2	1	1	-	-
Female	-	1	-	3	1	1	12	4	9	1	-	-
Black, both sexes	-	2	-	1	-	3	2	-	2	-	-	-
Male	-	2	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Minnesota												
All races, both sexes	-	-	-	-	1	4	7	2	4	1	-	-
Male	-	-	-	-	1	3	2	-	-	-	-	-
Female	-	-	-	-	-	1	5	2	4	1	-	-
White, both sexes	-	-	-	-	1	3	7	2	4	1	-	-
Male	-	-	-	-	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	5	2	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	1	1	1	-	1	3	1	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	3	1	-	-	-
White, both sexes	-	1	1	1	1	-	-	2	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Missouri												
All races, both sexes	1	-	1	3	3	5	6	3	2	1	-	-
Male	1	-	1	1	1	2	1	2	1	-	-	-
Female	-	-	-	2	2	3	5	1	1	1	-	-
White, both sexes	1	-	-	2	3	5	6	3	2	1	-	-
Male	1	-	-	1	1	2	1	2	1	-	-	-
Female	-	-	-	1	2	3	5	1	1	1	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	2	1	3	1	-	1	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	-	1	-	-
White, both sexes	1	-	-	-	2	1	3	1	-	1	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Nebraska												
All races, both sexes	-	-	1	-	1	3	-	4	2	1	-	-
Male	-	-	1	-	1	-	-	1	2	-	-	-
Female	-	-	-	-	-	3	-	3	-	1	-	-
White, both sexes	-	-	1	-	1	3	-	4	2	1	-	-
Male	-	-	1	-	1	-	-	1	2	-	-	-
Female	-	-	-	-	-	3	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	2	-	1	1	1	1	3	-	-	-
Male	1	-	1	-	-	-	1	1	2	-	-	-
Female	-	-	1	-	1	1	-	-	1	-	-	-
White, both sexes	1	-	1	-	1	1	1	1	3	-	-	-
Male	1	-	1	-	-	-	1	1	2	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
New Hampshire												
All races, both sexes	-	1	-	1	-	1	4	2	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	1	-	-	2	1	-	-	-	-
White, both sexes	-	1	-	1	-	1	4	2	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	1	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	-	3	7	5	6	11	5	4	-	-
Male	2	1	-	1	3	2	2	6	1	1	-	-
Female	1	1	-	2	4	3	4	5	4	3	-	-
White, both sexes	3	2	-	3	5	4	5	10	5	4	-	-
Male	2	1	-	1	1	1	2	6	1	1	-	-
Female	1	1	-	2	4	3	3	4	4	3	-	-
Black, both sexes	-	-	-	-	2	1	1	1	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
New Mexico												
All races, both sexes	-	-	-	2	1	1	1	1	-	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	1	-	-
White, both sexes	-	-	-	2	1	1	1	1	-	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	10	3	10	12	14	13	20	15	1	1	-
Male	1	6	1	4	5	7	5	11	4	-	-	-
Female	-	4	2	6	7	7	8	9	11	1	1	-
White, both sexes	1	8	2	8	10	13	12	18	13	1	1	-
Male	1	4	1	3	3	7	4	11	3	-	-	-
Female	-	4	1	5	7	6	8	7	10	1	1	-
Black, both sexes	-	2	1	2	2	1	1	2	1	-	-	-
Male	-	2	-	1	2	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
North Carolina												
All races, both sexes	6	3	6	5	4	10	7	8	5	2	1	-
Male	4	1	4	1	2	7	2	7	2	-	-	-
Female	2	2	2	4	2	3	5	1	3	2	1	-
White, both sexes	5	2	4	3	4	8	6	4	5	2	-	-
Male	3	1	3	1	2	6	1	4	2	-	-	-
Female	2	1	1	2	2	2	5	-	3	2	-	-
Black, both sexes	1	-	2	2	-	2	1	4	-	-	1	-
Male	1	-	1	-	-	1	1	3	-	-	-	-
Female	-	-	1	2	-	1	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	-	-	-	1	2	1	-	-
Male	-	-	1	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	1	1	-	-	-	1	2	1	-	-
Male	-	-	1	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Ohio												
All races, both sexes	1	4	4	2	9	11	11	10	7	5	-	-
Male	-	2	1	1	3	5	6	5	1	1	-	-
Female	1	2	3	1	6	6	5	5	6	4	-	-
White, both sexes	-	3	3	2	8	10	11	8	7	5	-	-
Male	-	1	1	1	3	5	6	4	1	1	-	-
Female	-	2	2	1	5	5	5	4	6	4	-	-
Black, both sexes	1	1	1	-	1	1	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	3	-	1	3	3	2	5	2	1	-	-
Male	1	2	-	1	1	-	1	3	1	-	-	-
Female	1	1	-	-	2	3	1	2	1	1	-	-
White, both sexes	2	3	-	1	2	3	2	5	2	1	-	-
Male	1	2	-	1	1	-	1	3	1	-	-	-
Female	1	1	-	-	1	3	1	2	1	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Oregon												
All races, both sexes	-	-	-	3	2	4	7	5	3	1	-	-
Male	-	-	-	-	1	3	2	3	1	-	-	-
Female	-	-	-	3	1	1	5	2	2	1	-	-
White, both sexes	-	-	-	3	2	4	7	5	3	1	-	-
Male	-	-	-	-	1	3	2	3	1	-	-	-
Female	-	-	-	3	1	1	5	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	6	2	3	7	20	8	14	9	4	-	-
Male	3	4	1	2	4	11	2	4	-	1	-	-
Female	1	2	1	1	3	9	6	10	9	3	-	-
White, both sexes	3	6	2	3	6	18	8	12	8	4	-	-
Male	2	4	1	2	3	10	2	3	-	1	-	-
Female	1	2	1	1	3	8	6	9	8	3	-	-
Black, both sexes	1	-	-	-	1	2	-	2	1	-	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Rhode Island												
All races, both sexes	-	-	-	2	-	2	2	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	2	1	-	1	1	-	-
White, both sexes	-	-	-	2	-	2	2	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	2	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	4	2	2	4	3	2	1	-	-
Male	2	-	-	2	-	1	2	-	-	-	-	-
Female	-	-	-	2	2	1	2	3	2	1	-	-
White, both sexes	2	-	-	3	1	2	3	1	2	1	-	-
Male	2	-	-	2	-	1	2	-	-	-	-	-
Female	-	-	-	1	1	1	1	1	2	1	-	-
Black, both sexes	-	-	-	1	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
South Dakota												
All races, both sexes	-	-	1	-	1	1	-	3	2	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	2	2	1	-	-
White, both sexes	-	-	1	-	1	1	-	3	2	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	1	3	4	7	1	3	3	6	1	-	-
Male	1	-	1	1	1	1	1	2	1	-	-	-
Female	1	1	2	3	6	-	2	1	5	1	-	-
White, both sexes	1	-	2	4	5	1	3	1	6	1	-	-
Male	-	-	1	1	1	1	1	1	1	-	-	-
Female	1	-	1	3	4	-	2	-	5	1	-	-
Black, both sexes	1	1	1	-	2	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	2	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Texas												
All races, both sexes	14	11	4	9	14	18	23	13	10	2	1	-
Male	13	7	3	6	8	8	12	3	2	-	-	-
Female	1	4	1	3	6	10	11	10	8	2	1	-
White, both sexes	8	10	2	6	11	14	19	11	9	2	-	-
Male	8	7	2	5	6	6	11	3	2	-	-	-
Female	-	3	-	1	5	8	8	8	7	2	-	-
Black, both sexes	5	1	2	2	3	4	3	1	1	-	1	-
Male	4	-	1	-	2	2	1	-	-	-	-	-
Female	1	1	1	2	1	2	2	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	-	2	-	1	1	2	-	1	-	2	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	2	-	-
White, both sexes	-	2	-	-	1	2	-	1	-	2	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Vermont												
All races, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	3	1	2	6	2	2	4	2	1	-
Male	-	1	2	-	2	4	-	1	1	1	-	-
Female	1	-	1	1	-	2	2	1	3	1	1	-
White, both sexes	1	1	1	1	2	3	2	2	4	2	1	-
Male	-	1	1	-	2	2	-	1	1	1	-	-
Female	1	-	-	1	-	1	2	1	3	1	1	-
Black, both sexes	-	-	2	-	-	3	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Washington												
All races, both sexes	-	3	5	-	2	2	8	4	4	3	-	-
Male	-	1	2	-	1	-	5	2	-	-	-	-
Female	-	2	3	-	1	2	3	2	4	3	-	-
White, both sexes	-	3	4	-	2	2	8	4	4	3	-	-
Male	-	1	2	-	1	-	5	2	-	-	-	-
Female	-	2	2	-	1	2	3	2	4	3	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	2	3	4	1	-	-	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	2	-	-	-	2	1	1	1	-	-	-	-
White, both sexes	2	-	-	-	2	2	4	1	-	-	-	-
Male	-	-	-	-	-	1	3	-	-	-	-	-
Female	2	-	-	-	2	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Wisconsin												
All races, both sexes	1	2	1	3	4	6	5	6	3	2	1	-
Male	1	-	-	3	2	4	1	4	1	1	-	-
Female	-	2	1	-	2	2	4	2	2	1	1	-
White, both sexes	1	2	1	3	4	6	5	6	3	2	1	-
Male	1	-	-	3	2	4	1	4	1	1	-	-
Female	-	2	1	-	2	2	4	2	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
United States												
All races, both sexes	2,532	-	2	3	1	2	7	12	17	25	37	57
Male	883	-	1	2	-	-	2	4	6	9	17	24
Female	1,649	-	1	1	1	2	5	8	11	16	20	33
White, both sexes	1,778	-	1	3	1	1	4	3	11	12	19	38
Male	631	-	1	2	-	-	1	2	4	6	9	17
Female	1,147	-	-	1	1	1	3	1	7	6	10	21
Black, both sexes	689	-	1	-	-	1	3	8	5	12	16	15
Male	231	-	-	-	-	-	1	2	1	3	8	7
Female	458	-	1	-	-	1	2	6	4	9	8	8
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	3
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	1	1	1	1	1
Male	20	-	-	-	-	-	-	-	1	-	-	-
Female	29	-	-	-	-	-	-	1	-	1	1	1
Alabama												
All races, both sexes	54	-	-	-	-	-	1	-	1	-	2	2
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	38	-	-	-	-	-	1	-	-	-	2	2
White, both sexes	24	-	-	-	-	-	1	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Arkansas												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	136	-	-	-	1	-	-	2	1	-	2	8
Male	50	-	-	-	-	-	-	-	-	-	1	2
Female	86	-	-	-	1	-	-	2	1	-	1	6
White, both sexes	99	-	-	-	1	-	-	-	-	-	-	7
Male	38	-	-	-	-	-	-	-	-	-	-	2
Female	61	-	-	-	1	-	-	-	-	-	-	5
Black, both sexes	23	-	-	-	-	-	-	2	1	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Colorado												
All races, both sexes	35	-	-	-	-	-	1	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	33	-	-	-	-	-	1	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	29	-	1	-	-	-	-	-	-	-	-	-
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	1	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Florida												
All races, both sexes	121	-	1	-	-	-	-	1	1	1	3	-
Male	44	-	-	-	-	-	-	-	-	-	2	-
Female	77	-	1	-	-	-	-	1	1	1	1	-
White, both sexes	89	-	-	-	-	-	-	1	1	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	54	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	32	-	1	-	-	-	-	-	-	1	2	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	1	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	109	-	-	-	-	-	-	-	1	1	-	3
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	77	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	63	-	-	-	-	-	-	-	-	1	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	40	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	45	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Hawaii												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Illinois												
All races, both sexes	116	-	-	-	-	-	1	-	-	4	1	1
Male	31	-	-	-	-	-	1	-	-	-	-	1
Female	85	-	-	-	-	-	-	-	-	4	1	-
White, both sexes	76	-	-	-	-	-	-	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	38	-	-	-	-	-	1	-	-	3	-	1
Male	13	-	-	-	-	-	1	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	60	-	-	1	-	-	-	-	1	-	3	3
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	34	-	-	1	-	-	-	-	1	-	2	2
White, both sexes	49	-	-	1	-	-	-	-	1	-	3	3
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	1	-	-	-	-	1	-	2	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Iowa												
All races, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Kansas												
All races, both sexes	18	-	-	1	-	-	-	1	-	-	-	-
Male	6	-	-	1	-	-	-	1	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	1	-	-	-	-	-	-	-	-
Male	5	-	-	1	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Kentucky												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	78	-	-	-	-	-	2	1	1	1	1	1
Male	28	-	-	-	-	-	-	-	-	1	1	1
Female	50	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	39	-	-	-	-	-	1	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	39	-	-	-	-	-	1	1	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	1	1	1
Female	23	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Maine												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	2	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Massachusetts												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	114	-	-	-	-	1	-	-	1	1	1	3
Male	44	-	-	-	-	-	-	-	-	-	1	2
Female	70	-	-	-	-	1	-	-	1	1	-	1
White, both sexes	78	-	-	-	-	-	-	-	-	-	1	3
Male	31	-	-	-	-	-	-	-	-	-	1	2
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	34	-	-	-	-	1	-	-	1	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	1	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Minnesota												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	57	-	-	-	-	-	-	2	-	-	2	1
Male	23	-	-	-	-	-	-	1	-	-	2	1
Female	34	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	1	-	-	2	-
Male	14	-	-	-	-	-	-	1	-	-	2	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Missouri												
All races, both sexes	49	-	-	-	-	-	-	1	1	1	2	5
Male	22	-	-	-	-	-	-	-	-	1	1	3
Female	27	-	-	-	-	-	-	1	1	-	1	2
White, both sexes	37	-	-	-	-	-	-	-	-	1	1	4
Male	17	-	-	-	-	-	-	-	-	1	1	2
Female	20	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	12	-	-	-	-	-	-	1	1	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	24	-	-	-	-	-	-	-	1	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
New Hampshire												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	127	-	-	-	-	-	1	-	-	2	-	4
Male	47	-	-	-	-	-	-	-	-	1	-	2
Female	80	-	-	-	-	-	1	-	-	1	-	2
White, both sexes	93	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	33	-	-	-	-	-	1	-	-	2	-	3
Male	12	-	-	-	-	-	-	-	-	1	-	2
Female	21	-	-	-	-	-	1	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
New Mexico												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	160	-	-	-	-	-	-	1	1	1	-	1
Male	60	-	-	-	-	-	-	-	1	-	-	1
Female	100	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	129	-	-	-	-	-	-	-	-	1	-	1
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	79	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	30	-	-	-	-	-	-	1	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
North Carolina												
All races, both sexes	104	-	-	1	-	-	-	1	-	1	1	2
Male	37	-	-	1	-	-	-	-	-	1	-	-
Female	67	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	67	-	-	1	-	-	-	-	-	1	-	1
Male	28	-	-	1	-	-	-	-	-	1	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	36	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Ohio												
All races, both sexes	97	-	-	-	-	-	-	-	-	1	1	2
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	71	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	76	-	-	-	-	-	-	-	-	1	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	40	-	-	-	-	-	-	-	-	1	3	2
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	32	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	31	-	-	-	-	-	-	-	-	1	2	1
Male	6	-	-	-	-	-	-	-	-	1	1	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Oregon												
All races, both sexes	25	-	-	-	-	-	1	-	1	-	1	1
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	24	-	-	-	-	-	1	-	1	-	-	1
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	126	-	-	-	-	-	-	1	-	1	1	3
Male	43	-	-	-	-	-	-	1	-	1	-	1
Female	83	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	103	-	-	-	-	-	-	1	-	-	-	2
Male	35	-	-	-	-	-	-	1	-	-	-	1
Female	68	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	53	-	-	-	-	-	-	-	-	2	-	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	39	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	64	-	-	-	-	-	-	-	1	1	1	2
Male	27	-	-	-	-	-	-	-	1	-	-	1
Female	37	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	47	-	-	-	-	-	-	-	1	1	1	1
Male	20	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Texas												
All races, both sexes	168	-	-	-	-	1	-	1	2	3	3	4
Male	64	-	-	-	-	-	-	1	2	1	2	2
Female	104	-	-	-	-	1	-	-	-	2	1	2
White, both sexes	113	-	-	-	-	1	-	1	2	2	3	3
Male	37	-	-	-	-	-	-	1	2	1	2	1
Female	76	-	-	-	-	1	-	-	-	1	1	2
Black, both sexes	55	-	-	-	-	-	-	-	-	1	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Vermont												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	103	-	-	-	-	-	-	-	1	-	1	1
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	66	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	56	-	-	-	-	-	-	-	1	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	47	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Washington												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	32	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Wisconsin												
All races, both sexes	37	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	31	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
United States												
All races, both sexes	66	95	122	144	200	312	457	428	345	160	40	-
Male	31	50	48	69	82	122	161	121	86	38	10	-
Female	35	45	74	75	118	190	296	307	259	122	30	-
White, both sexes	37	55	79	97	123	209	340	332	270	118	25	-
Male	19	30	28	50	48	85	123	99	72	30	5	-
Female	18	25	51	47	75	124	217	233	198	88	20	-
Black, both sexes	28	38	40	44	68	93	105	88	69	41	14	-
Male	11	19	19	18	29	36	33	21	10	8	5	-
Female	17	19	21	26	39	57	72	67	59	33	9	-
American Indian 1/, both sexes	-	-	1	-	3	-	3	1	2	1	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	2	-	3	1	2	1	1	-
Asian or Pacific Islander 2/, both sexes	1	2	2	3	6	10	9	7	4	-	-	-
Male	1	1	1	1	4	1	5	1	4	-	-	-
Female	-	1	1	2	2	9	4	6	-	-	-	-
Alabama												
All races, both sexes	1	1	2	3	5	8	10	5	7	4	2	-
Male	-	1	1	1	2	2	6	-	1	-	1	-
Female	1	-	1	2	3	6	4	5	6	4	1	-
White, both sexes	-	1	1	-	2	3	5	2	4	1	1	-
Male	-	1	-	-	-	1	2	-	1	-	1	-
Female	-	-	1	-	2	2	3	2	3	1	-	-
Black, both sexes	1	-	1	3	2	5	5	3	3	3	1	-
Male	-	-	1	1	1	1	4	-	-	-	-	-
Female	1	-	-	2	1	4	1	3	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Alaska												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	2	3	2	4	7	4	6	8	2	1	-
Male	2	-	1	2	2	4	3	3	2	-	-	-
Female	1	2	2	-	2	3	1	3	6	2	1	-
White, both sexes	3	2	3	2	2	7	3	6	7	1	-	-
Male	2	-	1	2	1	4	3	3	2	-	-	-
Female	1	2	2	-	1	3	-	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	1	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Arkansas												
All races, both sexes	1	3	1	-	4	4	5	6	3	1	1	-
Male	-	1	1	-	-	1	2	1	-	-	-	-
Female	1	2	-	-	4	3	3	5	3	1	1	-
White, both sexes	1	1	-	-	2	2	1	5	3	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	2	2	-	5	3	1	-	-
Black, both sexes	-	2	1	-	2	2	4	1	-	-	1	-
Male	-	1	1	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	2	1	3	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	10	6	6	10	12	28	25	16	7	1	-
Male	1	6	3	4	6	5	6	7	8	1	-	-
Female	-	4	3	2	4	7	22	18	8	6	1	-
White, both sexes	-	6	6	5	5	7	21	21	13	7	-	-
Male	-	4	3	4	3	4	6	5	6	1	-	-
Female	-	2	3	1	2	3	15	16	7	6	-	-
Black, both sexes	1	3	-	-	3	3	5	2	1	-	1	-
Male	1	1	-	-	2	-	-	1	-	-	-	-
Female	-	2	-	-	1	3	5	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	2	2	2	2	-	-	-
Male	-	1	-	-	1	1	-	1	2	-	-	-
Female	-	-	-	1	1	1	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Colorado												
All races, both sexes	1	2	2	-	3	7	4	3	4	4	1	-
Male	-	2	-	-	1	1	1	3	-	1	1	-
Female	1	-	2	-	2	6	3	-	4	3	-	-
White, both sexes	1	2	2	-	2	7	4	3	4	4	1	-
Male	-	2	-	-	1	1	1	3	-	1	1	-
Female	1	-	2	-	1	6	3	-	4	3	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	1	4	3	8	2	7	2	-	-
Male	-	-	-	1	1	2	3	-	1	2	-	-
Female	-	-	1	-	3	1	5	2	6	-	-	-
White, both sexes	-	-	1	-	4	2	7	2	7	2	-	-
Male	-	-	-	-	1	1	2	-	1	2	-	-
Female	-	-	1	-	3	1	5	2	6	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Delaware												
All races, both sexes	-	-	-	3	-	1	1	2	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	2	1	-	-	-
White, both sexes	-	-	-	3	-	1	1	2	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	2	2	1	2	2	-	4	2	-	-
Male	-	-	-	-	-	2	-	-	1	2	-	-
Female	1	-	2	2	1	-	2	-	3	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	2	2	1	2	2	-	4	2	-	-
Male	-	-	-	-	-	2	-	-	1	2	-	-
Female	1	-	2	2	1	-	2	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Florida												
All races, both sexes	3	3	4	7	11	9	34	16	17	8	2	-
Male	1	1	2	3	5	5	10	6	5	4	-	-
Female	2	2	2	4	6	4	24	10	12	4	2	-
White, both sexes	-	2	1	5	8	8	27	14	13	8	-	-
Male	-	1	-	3	2	5	9	6	4	4	-	-
Female	-	1	1	2	6	3	18	8	9	4	-	-
Black, both sexes	3	1	3	2	3	1	7	2	4	-	2	-
Male	1	-	2	-	3	-	1	-	1	-	-	-
Female	2	1	1	2	-	1	6	2	3	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	9	6	4	6	12	11	32	16	5	2	-
Male	-	7	4	1	-	4	2	10	-	1	1	-
Female	1	2	2	3	6	8	9	22	16	4	1	-
White, both sexes	-	4	5	1	4	7	8	18	10	2	1	-
Male	-	4	3	-	-	4	1	9	-	-	-	-
Female	-	-	2	1	4	3	7	9	10	2	1	-
Black, both sexes	1	5	1	3	2	5	3	13	6	3	1	-
Male	-	3	1	1	-	-	1	1	-	1	1	-
Female	1	2	-	2	2	5	2	12	6	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Hawaii												
All races, both sexes	1	1	1	-	1	5	5	1	1	-	-	-
Male	1	1	1	-	-	-	3	-	1	-	-	-
Female	-	-	-	-	1	5	2	1	-	-	-	-
White, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	4	4	1	1	-	-	-
Male	1	-	1	-	-	-	3	-	1	-	-	-
Female	-	-	-	-	1	4	1	1	-	-	-	-
Idaho												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Illinois												
All races, both sexes	4	10	6	2	13	15	17	20	14	6	2	-
Male	2	3	1	1	3	6	4	6	3	-	-	-
Female	2	7	5	1	10	9	13	14	11	6	2	-
White, both sexes	1	6	2	2	7	9	10	17	14	4	2	-
Male	1	1	-	1	1	3	3	5	3	-	-	-
Female	-	5	2	1	6	6	7	12	11	4	2	-
Black, both sexes	3	3	4	-	6	5	7	3	-	2	-	-
Male	1	2	1	-	2	3	1	1	-	-	-	-
Female	2	1	3	-	4	2	6	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	1	3	5	7	6	6	8	8	5	3	-	-
Male	1	1	3	3	3	1	5	4	3	-	-	-
Female	-	2	2	4	3	5	3	4	2	3	-	-
White, both sexes	1	2	4	6	4	5	6	6	5	2	-	-
Male	1	-	2	3	1	1	5	4	3	-	-	-
Female	-	2	2	3	3	4	1	2	2	2	-	-
Black, both sexes	-	1	1	1	2	1	2	2	-	1	-	-
Male	-	1	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Iowa												
All races, both sexes	-	-	-	-	1	-	1	4	5	1	1	-
Male	-	-	-	-	1	-	-	2	4	1	-	-
Female	-	-	-	-	-	-	1	2	1	-	1	-
White, both sexes	-	-	-	-	1	-	1	4	5	1	1	-
Male	-	-	-	-	1	-	-	2	4	1	-	-
Female	-	-	-	-	-	-	1	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	2	2	2	5	3	1	-	-
Male	1	-	-	-	-	-	1	-	1	1	-	-
Female	-	-	-	-	2	2	1	5	2	-	-	-
White, both sexes	1	-	-	-	2	2	2	4	3	1	-	-
Male	1	-	-	-	-	-	1	-	1	1	-	-
Female	-	-	-	-	2	2	1	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Kentucky												
All races, both sexes	-	1	1	2	2	6	8	12	2	-	1	-
Male	-	-	-	2	1	3	3	2	-	-	-	-
Female	-	1	1	-	1	3	5	10	2	-	1	-
White, both sexes	-	1	1	1	2	3	6	10	2	-	1	-
Male	-	-	-	1	1	2	1	2	-	-	-	-
Female	-	1	1	-	1	1	5	8	2	-	1	-
Black, both sexes	-	-	-	1	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Louisiana												
All races, both sexes	4	3	6	7	5	8	12	9	10	6	1	-
Male	1	2	2	3	3	4	4	2	2	1	1	-
Female	3	1	4	4	2	4	8	7	8	5	-	-
White, both sexes	3	-	2	3	1	5	7	5	7	4	-	-
Male	1	-	-	2	-	3	3	2	1	-	-	-
Female	2	-	2	1	1	2	4	3	6	4	-	-
Black, both sexes	1	3	4	4	4	3	5	4	3	2	1	-
Male	-	2	2	1	3	1	1	-	1	1	1	-
Female	1	1	2	3	1	2	4	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Maine												
All races, both sexes	-	1	2	-	1	-	3	1	-	1	-	-
Male	-	1	1	-	1	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	3	1	-	-	-	-
White, both sexes	-	1	2	-	1	-	3	1	-	1	-	-
Male	-	1	1	-	1	-	-	-	-	1	-	-
Female	-	-	1	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	-	4	7	7	6	8	10	4	4	-	-
Male	-	-	2	1	1	3	2	3	1	1	-	-
Female	3	-	2	6	6	3	6	7	3	3	-	-
White, both sexes	-	-	1	3	3	2	3	5	2	2	-	-
Male	-	-	1	-	-	1	2	1	1	1	-	-
Female	-	-	-	3	3	1	1	4	1	1	-	-
Black, both sexes	3	-	3	4	4	4	5	5	2	2	-	-
Male	-	-	1	1	1	2	-	2	-	-	-	-
Female	3	-	2	3	3	2	5	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Massachusetts												
All races, both sexes	-	2	-	-	-	5	13	15	6	6	1	-
Male	-	1	-	-	-	-	2	5	1	4	-	-
Female	-	1	-	-	-	5	11	10	5	2	1	-
White, both sexes	-	2	-	-	-	4	8	15	5	6	1	-
Male	-	1	-	-	-	-	2	5	1	4	-	-
Female	-	1	-	-	-	4	6	10	4	2	1	-
Black, both sexes	-	-	-	-	-	1	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	2	4	8	8	11	20	20	20	9	3	2	-
Male	1	2	4	4	3	8	9	7	3	-	-	-
Female	1	2	4	4	8	12	11	13	6	3	2	-
White, both sexes	1	3	4	6	5	13	16	14	7	3	2	-
Male	-	1	2	3	1	4	8	6	3	-	-	-
Female	1	2	2	3	4	9	8	8	4	3	2	-
Black, both sexes	1	1	4	2	5	7	4	5	2	-	-	-
Male	1	1	2	1	1	4	1	1	-	-	-	-
Female	-	-	2	1	4	3	3	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Minnesota												
All races, both sexes	1	-	1	2	5	2	1	3	2	4	1	-
Male	1	-	-	-	1	-	1	1	-	3	1	-
Female	-	-	1	2	4	2	-	2	2	1	-	-
White, both sexes	1	-	1	2	5	2	1	3	2	4	1	-
Male	1	-	-	-	1	-	1	1	-	3	1	-
Female	-	-	1	2	4	2	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	4	-	2	4	10	11	7	7	3	1	-
Male	2	3	-	1	2	3	3	2	2	-	1	-
Female	1	1	-	1	2	7	8	5	5	3	-	-
White, both sexes	1	1	-	1	2	1	6	3	2	1	-	-
Male	1	1	-	-	1	1	-	2	1	-	-	-
Female	-	-	-	1	1	-	6	1	1	1	-	-
Black, both sexes	2	3	-	1	2	9	4	4	5	2	1	-
Male	1	2	-	1	1	2	2	-	1	-	1	-
Female	1	1	-	-	1	7	2	4	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Missouri												
All races, both sexes	3	-	2	5	1	4	12	4	5	2	1	-
Male	1	-	-	5	1	-	5	3	1	1	-	-
Female	2	-	2	-	-	4	7	1	4	1	1	-
White, both sexes	1	-	2	3	-	3	12	4	4	2	-	-
Male	-	-	-	3	-	-	5	3	1	1	-	-
Female	1	-	2	-	-	3	7	1	3	1	-	-
Black, both sexes	2	-	-	2	1	1	-	-	1	-	1	-
Male	1	-	-	2	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	-	2	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	2	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Nebraska												
All races, both sexes	-	-	-	1	1	2	2	2	3	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	1	1	2	2	2	-	-	-
White, both sexes	-	-	-	-	1	2	2	2	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	4	3	1	2	3	5	4	-	-	-
Male	-	-	1	1	1	-	-	3	2	-	-	-
Female	-	-	3	2	-	2	3	2	2	-	-	-
White, both sexes	-	-	2	2	1	2	2	5	3	-	-	-
Male	-	-	1	-	1	-	-	3	1	-	-	-
Female	-	-	1	2	-	2	2	2	2	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
New Hampshire												
All races, both sexes	-	-	2	-	-	1	1	1	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	1	-	-
White, both sexes	-	-	2	-	-	1	1	1	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	2	1	7	7	22	23	25	15	10	1	-
Male	2	-	1	3	6	10	10	6	6	-	-	-
Female	5	2	-	4	1	12	13	19	9	10	1	-
White, both sexes	4	-	1	6	5	16	16	22	13	8	1	-
Male	1	-	1	3	4	9	7	5	5	-	-	-
Female	3	-	-	3	1	7	9	17	8	8	1	-
Black, both sexes	3	2	-	1	2	6	7	2	2	2	-	-
Male	1	-	-	-	2	1	3	1	1	-	-	-
Female	2	2	-	1	-	5	4	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
New Mexico												
All races, both sexes	1	-	1	2	2	1	2	-	-	-	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	1	-	1	1	-	-	2	-	-	-	-	-
White, both sexes	1	-	1	2	1	1	1	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	1	-	1	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	5	4	9	7	22	32	30	27	14	3	-
Male	2	1	1	2	5	11	16	7	9	4	-	-
Female	1	4	3	7	2	11	16	23	18	10	3	-
White, both sexes	1	3	3	7	5	18	30	24	25	9	2	-
Male	1	1	-	2	4	9	16	6	8	2	-	-
Female	-	2	3	5	1	9	14	18	17	7	2	-
Black, both sexes	2	2	1	2	2	3	2	6	2	5	1	-
Male	1	-	1	-	1	2	-	1	1	2	-	-
Female	1	2	-	2	1	1	2	5	1	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
North Carolina												
All races, both sexes	3	3	5	7	10	15	19	14	15	7	-	-
Male	2	3	2	3	4	8	6	4	1	2	-	-
Female	1	-	3	4	6	7	13	10	14	5	-	-
White, both sexes	3	1	4	5	6	11	13	9	7	5	-	-
Male	2	1	2	2	3	6	4	3	1	2	-	-
Female	1	-	2	3	3	5	9	6	6	3	-	-
Black, both sexes	-	2	1	2	3	4	6	5	8	2	-	-
Male	-	2	-	1	1	2	2	1	-	-	-	-
Female	-	-	1	1	2	2	4	4	8	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	-	1	-	1	-	1	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	1	-	1	-	1	-	1	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Ohio												
All races, both sexes	3	5	4	4	6	11	10	23	23	4	-	-
Male	2	4	2	1	1	4	4	2	4	-	-	-
Female	1	1	2	3	5	7	6	21	19	4	-	-
White, both sexes	1	4	4	3	5	10	9	19	17	3	-	-
Male	-	3	2	1	1	3	4	2	4	-	-	-
Female	1	1	2	2	4	7	5	17	13	3	-	-
Black, both sexes	2	1	-	1	1	1	1	4	6	1	-	-
Male	2	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	4	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	3	2	1	5	11	5	4	1	-	-
Male	-	1	-	-	-	1	1	1	2	-	-	-
Female	1	-	3	2	1	4	10	4	2	1	-	-
White, both sexes	1	1	2	1	-	4	11	4	3	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	1	-	2	1	-	3	10	4	2	-	-	-
Black, both sexes	-	-	1	1	1	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	1	1	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Oregon												
All races, both sexes	-	-	-	2	1	3	3	3	5	2	2	-
Male	-	-	-	1	-	1	1	1	-	1	1	-
Female	-	-	-	1	1	2	2	2	5	1	1	-
White, both sexes	-	-	-	2	1	3	3	3	5	2	2	-
Male	-	-	-	1	-	1	1	1	-	1	1	-
Female	-	-	-	1	1	2	2	2	5	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	1	6	8	4	19	25	25	17	10	2	-
Male	3	-	2	3	3	9	13	4	2	1	-	-
Female	-	1	4	5	1	10	12	21	15	9	2	-
White, both sexes	2	1	4	7	3	14	23	22	15	8	1	-
Male	2	-	1	3	2	6	12	4	2	1	-	-
Female	-	1	3	4	1	8	11	18	13	7	1	-
Black, both sexes	1	-	2	1	1	4	2	3	2	2	1	-
Male	1	-	1	-	1	3	1	-	-	-	-	-
Female	-	-	1	1	-	1	1	3	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Rhode Island												
All races, both sexes	-	-	-	-	1	-	-	1	1	-	1	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	-	-	-	-	1	-	-	1	1	-	1	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	2	6	5	2	9	10	6	6	4	1	-
Male	-	1	3	2	1	1	2	1	1	1	-	-
Female	-	1	3	3	1	8	8	5	5	3	1	-
White, both sexes	-	1	3	2	-	1	5	3	6	2	1	-
Male	-	1	1	1	-	-	1	1	1	1	-	-
Female	-	-	2	1	-	1	4	2	5	1	1	-
Black, both sexes	-	1	3	3	2	8	5	3	-	2	-	-
Male	-	-	2	1	1	1	1	-	-	-	-	-
Female	-	1	1	2	1	7	4	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	2	4	5	9	9	12	9	3	-	-
Male	1	3	2	4	2	4	3	4	1	1	-	-
Female	1	1	-	-	3	5	6	8	8	2	-	-
White, both sexes	1	2	1	3	5	6	6	10	6	3	-	-
Male	1	2	1	3	2	3	2	3	-	1	-	-
Female	-	-	-	-	3	3	4	7	6	2	-	-
Black, both sexes	1	2	1	1	-	3	3	2	3	-	-	-
Male	-	1	1	1	-	1	1	1	1	-	-	-
Female	1	1	-	-	-	2	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Texas												
All races, both sexes	3	5	7	6	22	15	33	27	23	10	3	-
Male	1	1	2	5	8	5	13	9	8	2	2	-
Female	2	4	5	1	14	10	20	18	15	8	1	-
White, both sexes	3	3	5	3	14	11	23	17	17	5	-	-
Male	1	-	1	2	5	3	8	3	6	1	-	-
Female	2	3	4	1	9	8	15	14	11	4	-	-
Black, both sexes	-	2	2	3	8	4	10	10	6	5	3	-
Male	-	1	1	3	3	2	5	6	2	1	2	-
Female	-	1	1	-	5	2	5	4	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	1	-	3	2	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	3	1	1	-	-	-
White, both sexes	1	-	-	-	1	-	3	2	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Vermont												
All races, both sexes	-	-	1	-	-	-	1	2	1	1	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	1	-	-	-	1	2	1	1	-	-
Male	-	-	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	4	7	5	13	11	22	10	14	8	4	-
Male	1	1	1	4	5	6	8	4	3	1	1	-
Female	1	3	6	1	8	5	14	6	11	7	3	-
White, both sexes	2	-	2	2	4	6	13	5	11	5	4	-
Male	1	-	-	2	2	3	3	1	3	1	1	-
Female	1	-	2	-	2	3	10	4	8	4	3	-
Black, both sexes	-	4	5	3	9	5	9	5	3	3	-	-
Male	-	1	1	2	3	3	5	3	-	-	-	-
Female	-	3	4	1	6	2	4	2	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Washington												
All races, both sexes	1	3	1	4	3	-	2	7	8	3	1	-
Male	1	2	1	2	2	-	1	2	2	-	-	-
Female	-	1	-	2	1	-	1	5	6	3	1	-
White, both sexes	1	3	1	4	3	-	2	5	7	3	1	-
Male	1	2	1	2	2	-	1	2	2	-	-	-
Female	-	1	-	2	1	-	1	3	5	3	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	-	3	3	2	2	4	4	4	2	-	-
Male	-	-	-	2	1	1	2	1	1	-	-	-
Female	-	-	3	1	1	1	2	3	3	2	-	-
White, both sexes	-	-	3	3	1	1	4	4	3	2	-	-
Male	-	-	-	2	1	-	2	1	1	-	-	-
Female	-	-	3	1	-	1	2	3	2	2	-	-
Black, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Wisconsin												
All races, both sexes	-	-	-	1	3	6	8	4	7	4	1	-
Male	-	-	-	1	2	3	3	-	2	1	-	-
Female	-	-	-	-	1	3	5	4	5	3	1	-
White, both sexes	-	-	-	1	3	5	7	4	5	3	1	-
Male	-	-	-	1	2	2	2	-	2	-	-	-
Female	-	-	-	-	1	3	5	4	3	3	1	-
Black, both sexes	-	-	-	-	-	1	1	-	2	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
United States												
All races, both sexes	14,418	10	14	7	30	53	110	123	170	221	375	502
Male	4,361	9	6	2	9	21	31	27	47	66	124	177
Female	10,057	1	8	5	21	32	79	96	123	155	251	325
White, both sexes	12,235	8	8	6	20	30	54	64	81	120	228	319
Male	3,710	7	4	2	6	12	21	16	27	42	84	128
Female	8,525	1	4	4	14	18	33	48	54	78	144	191
Black, both sexes	1,846	2	6	-	9	21	47	55	74	95	134	165
Male	546	2	2	-	3	8	9	11	19	23	39	45
Female	1,300	-	4	-	6	13	38	44	55	72	95	120
American Indian 1/, both sexes	102	-	-	-	-	1	2	1	3	-	4	5
Male	35	-	-	-	-	-	-	-	1	-	1	1
Female	67	-	-	-	-	1	2	1	2	-	3	4
Asian or Pacific Islander 2/, both sexes	235	-	-	1	1	1	7	3	12	6	9	13
Male	70	-	-	-	-	1	1	-	-	1	-	3
Female	165	-	-	1	1	-	6	3	12	5	9	10
Alabama												
All races, both sexes	212	-	-	1	-	2	2	4	1	3	14	4
Male	62	-	-	-	-	1	1	1	-	1	4	1
Female	150	-	-	1	-	1	1	3	1	2	10	3
White, both sexes	159	-	-	1	-	2	1	1	-	1	6	3
Male	47	-	-	-	-	1	1	-	-	1	3	-
Female	112	-	-	1	-	1	-	1	-	-	3	3
Black, both sexes	53	-	-	-	-	-	1	3	1	2	8	1
Male	15	-	-	-	-	-	-	1	-	-	1	1
Female	38	-	-	-	-	-	1	2	1	2	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Alaska												
All races, both sexes	25	-	-	-	-	-	-	-	-	1	2	1
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	250	-	1	-	1	1	3	1	4	4	3	9
Male	80	-	1	-	1	-	1	1	1	-	1	7
Female	170	-	-	-	-	1	2	-	3	4	2	2
White, both sexes	234	-	1	-	1	1	1	1	2	4	2	9
Male	75	-	1	-	1	-	1	1	-	-	1	7
Female	159	-	-	-	-	1	-	-	2	4	1	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	1	-	2	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Arkansas												
All races, both sexes	124	-	-	-	-	-	2	-	1	2	5	1
Male	40	-	-	-	-	-	-	-	-	2	2	1
Female	84	-	-	-	-	-	2	-	1	-	3	-
White, both sexes	106	-	-	-	-	-	-	-	1	1	4	1
Male	36	-	-	-	-	-	-	-	-	1	2	1
Female	70	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	18	-	-	-	-	-	2	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,391	2	1	1	4	8	12	15	21	24	46	59
Male	420	1	1	-	1	1	2	2	6	5	18	21
Female	971	1	-	1	3	7	10	13	15	19	28	38
White, both sexes	1,158	2	1	1	2	5	7	9	13	17	24	44
Male	340	1	1	-	-	-	1	1	5	5	9	19
Female	818	1	-	1	2	5	6	8	8	12	15	25
Black, both sexes	134	-	-	-	2	2	2	5	3	4	17	12
Male	45	-	-	-	1	-	-	1	1	-	8	1
Female	89	-	-	-	1	2	2	4	2	4	9	11
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	89	-	-	-	-	1	3	1	5	3	4	2
Male	30	-	-	-	-	1	1	-	-	-	-	-
Female	59	-	-	-	-	-	2	1	5	3	4	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Colorado												
All races, both sexes	254	-	1	-	-	-	3	1	-	2	6	7
Male	70	-	-	-	-	-	1	-	-	1	1	1
Female	184	-	1	-	-	-	2	1	-	1	5	6
White, both sexes	239	-	1	-	-	-	2	-	-	2	6	4
Male	65	-	-	-	-	-	1	-	-	1	1	-
Female	174	-	1	-	-	-	1	-	-	1	5	4
Black, both sexes	12	-	-	-	-	-	1	1	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	1	1	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	170	-	-	-	-	2	-	2	-	1	7	5
Male	47	-	-	-	-	1	-	1	-	-	3	1
Female	123	-	-	-	-	1	-	1	-	1	4	4
White, both sexes	152	-	-	-	-	1	-	1	-	-	4	2
Male	38	-	-	-	-	1	-	-	-	-	1	-
Female	114	-	-	-	-	-	-	1	-	-	3	2
Black, both sexes	18	-	-	-	-	1	-	1	-	1	3	3
Male	9	-	-	-	-	-	-	1	-	-	2	1
Female	9	-	-	-	-	1	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Delaware												
All races, both sexes	42	-	1	-	-	-	-	-	-	-	4	2
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	3	2
White, both sexes	33	-	-	-	-	-	-	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	9	-	1	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	-	-	-	-	-	-	1	1	1	2	3
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	1	1	1	1	3
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	1	1	1	2	3
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	1	1	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Florida												
All races, both sexes	801	-	1	-	3	4	15	8	12	14	26	29
Male	227	-	1	-	-	1	3	2	3	6	7	6
Female	574	-	-	-	3	3	12	6	9	8	19	23
White, both sexes	700	-	1	-	1	3	9	5	6	10	16	20
Male	194	-	1	-	-	-	2	1	1	4	5	5
Female	506	-	-	-	1	3	7	4	5	6	11	15
Black, both sexes	97	-	-	-	2	1	5	3	6	4	10	8
Male	32	-	-	-	-	1	1	1	2	2	2	1
Female	65	-	-	-	2	-	4	2	4	2	8	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	1
Georgia												
All races, both sexes	374	-	1	-	-	2	9	6	12	9	13	23
Male	144	-	-	-	-	1	4	2	5	6	5	11
Female	230	-	1	-	-	1	5	4	7	3	8	12
White, both sexes	240	-	1	-	-	-	-	1	5	-	6	10
Male	93	-	-	-	-	-	-	1	3	-	3	4
Female	147	-	1	-	-	-	-	-	2	-	3	6
Black, both sexes	129	-	-	-	-	2	8	5	7	9	7	13
Male	50	-	-	-	-	1	4	1	2	6	2	7
Female	79	-	-	-	-	1	4	4	5	3	5	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Hawaii												
All races, both sexes	61	-	-	-	1	-	-	-	2	1	-	4
Male	25	-	-	-	-	-	-	-	1	-	-	4
Female	36	-	-	-	1	-	-	-	1	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	1	-	-	3
Male	9	-	-	-	-	-	-	-	1	-	-	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	1	-	-	-	1	1	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	1	-	-	-	1	1	-	-
Idaho												
All races, both sexes	94	-	-	-	-	-	-	2	1	-	2	2
Male	28	-	-	-	-	-	-	-	1	-	1	1
Female	66	-	-	-	-	-	-	2	-	-	1	1
White, both sexes	90	-	-	-	-	-	-	1	1	-	2	1
Male	27	-	-	-	-	-	-	-	1	-	1	1
Female	63	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Illinois												
All races, both sexes	665	3	1	-	1	4	3	7	9	13	12	22
Male	196	3	-	-	-	2	1	1	1	4	2	5
Female	469	-	1	-	1	2	2	6	8	9	10	17
White, both sexes	557	2	1	-	1	3	2	3	4	6	7	13
Male	166	2	-	-	-	1	1	1	1	2	-	3
Female	391	-	1	-	1	2	1	2	3	4	7	10
Black, both sexes	98	1	-	-	-	1	1	4	4	7	5	7
Male	27	1	-	-	-	1	-	-	-	2	2	2
Female	71	-	-	-	-	-	1	4	4	5	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	2
Indiana												
All races, both sexes	308	-	-	-	-	-	4	2	3	4	8	10
Male	98	-	-	-	-	-	-	1	-	2	4	5
Female	210	-	-	-	-	-	4	1	3	2	4	5
White, both sexes	271	-	-	-	-	-	2	1	-	2	5	6
Male	85	-	-	-	-	-	-	1	-	1	3	3
Female	186	-	-	-	-	-	2	-	-	1	2	3
Black, both sexes	36	-	-	-	-	-	2	1	3	2	3	4
Male	13	-	-	-	-	-	-	-	-	1	1	2
Female	23	-	-	-	-	-	2	1	3	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Iowa												
All races, both sexes	200	-	-	-	1	-	1	1	-	-	2	6
Male	74	-	-	-	-	-	-	1	-	-	1	2
Female	126	-	-	-	1	-	1	-	-	-	1	4
White, both sexes	196	-	-	-	1	-	-	1	-	-	2	4
Male	72	-	-	-	-	-	-	1	-	-	1	1
Female	124	-	-	-	1	-	-	-	-	-	1	3
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	189	-	1	1	1	-	1	1	-	1	2	5
Male	53	-	1	-	-	-	-	-	-	-	1	2
Female	136	-	-	1	1	-	1	1	-	1	1	3
White, both sexes	172	-	-	1	-	-	1	1	-	-	2	5
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	123	-	-	1	-	-	1	1	-	-	1	3
Black, both sexes	13	-	1	-	1	-	-	-	-	1	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Kentucky												
All races, both sexes	223	-	-	-	1	1	2	-	2	2	2	6
Male	71	-	-	-	-	-	-	-	-	1	2	1
Female	152	-	-	-	1	1	2	-	2	1	-	5
White, both sexes	208	-	-	-	1	1	2	-	1	1	1	4
Male	66	-	-	-	-	-	-	-	-	-	1	1
Female	142	-	-	-	1	1	2	-	1	1	-	3
Black, both sexes	15	-	-	-	-	-	-	-	1	1	1	2
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	220	-	-	-	-	1	3	3	3	6	6	13
Male	64	-	-	-	-	-	1	-	-	-	2	5
Female	156	-	-	-	-	1	2	3	3	6	4	8
White, both sexes	127	-	-	-	-	-	-	-	1	-	3	4
Male	50	-	-	-	-	-	-	-	-	-	1	4
Female	77	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	93	-	-	-	-	1	3	3	2	6	3	9
Male	14	-	-	-	-	-	1	-	-	-	1	1
Female	79	-	-	-	-	1	2	3	2	6	2	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Maine												
All races, both sexes	113	-	-	-	-	1	1	1	1	-	2	3
Male	35	-	-	-	-	1	-	-	-	-	-	1
Female	78	-	-	-	-	-	1	1	1	-	2	2
White, both sexes	112	-	-	-	-	1	1	1	1	-	2	3
Male	35	-	-	-	-	1	-	-	-	-	-	1
Female	77	-	-	-	-	-	1	1	1	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	294	-	-	-	2	1	1	2	1	3	10	12
Male	70	-	-	-	-	-	1	1	-	-	3	3
Female	224	-	-	-	2	1	-	1	1	3	7	9
White, both sexes	214	-	-	-	2	1	1	1	1	1	2	7
Male	45	-	-	-	-	-	1	-	-	-	1	1
Female	169	-	-	-	2	1	-	1	1	1	1	6
Black, both sexes	77	-	-	-	-	-	-	1	-	2	8	5
Male	25	-	-	-	-	-	-	1	-	-	2	2
Female	52	-	-	-	-	-	-	-	-	2	6	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Massachusetts												
All races, both sexes	257	-	-	-	-	-	-	-	2	1	4	8
Male	74	-	-	-	-	-	-	-	-	-	2	2
Female	183	-	-	-	-	-	-	-	2	1	2	6
White, both sexes	243	-	-	-	-	-	-	-	2	1	2	5
Male	71	-	-	-	-	-	-	-	-	-	1	1
Female	172	-	-	-	-	-	-	-	2	1	1	4
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	3
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	522	-	-	-	1	4	1	-	7	10	10	16
Male	175	-	-	-	-	2	-	-	2	4	2	7
Female	347	-	-	-	1	2	1	-	5	6	8	9
White, both sexes	439	-	-	-	-	1	-	-	1	4	5	6
Male	148	-	-	-	-	1	-	-	-	3	1	2
Female	291	-	-	-	-	-	-	-	1	1	4	4
Black, both sexes	78	-	-	-	1	3	1	-	4	6	5	10
Male	26	-	-	-	-	1	-	-	2	1	1	5
Female	52	-	-	-	1	2	1	-	2	5	4	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Minnesota												
All races, both sexes	336	-	1	-	2	1	-	2	2	1	4	5
Male	94	-	-	-	1	-	-	1	1	1	2	2
Female	242	-	1	-	1	1	-	1	1	-	2	3
White, both sexes	322	-	-	-	2	1	-	2	1	1	2	3
Male	91	-	-	-	1	-	-	1	1	1	2	1
Female	231	-	-	-	1	1	-	1	-	-	-	2
Black, both sexes	8	-	1	-	-	-	-	-	1	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	1	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	104	-	1	-	-	-	2	3	2	1	5	5
Male	40	-	1	-	-	-	-	2	1	1	3	3
Female	64	-	-	-	-	-	2	1	1	-	2	2
White, both sexes	63	-	-	-	-	-	-	3	-	-	2	1
Male	24	-	-	-	-	-	-	2	-	-	2	1
Female	39	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	41	-	1	-	-	-	2	-	2	1	3	4
Male	16	-	1	-	-	-	-	-	1	1	1	2
Female	25	-	-	-	-	-	2	-	1	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Missouri												
All races, both sexes	320	-	1	-	-	-	2	2	1	7	8	7
Male	97	-	1	-	-	-	-	-	-	2	3	1
Female	223	-	-	-	-	-	2	2	1	5	5	6
White, both sexes	284	-	1	-	-	-	-	2	-	7	5	6
Male	85	-	1	-	-	-	-	-	-	2	3	1
Female	199	-	-	-	-	-	-	2	-	5	2	5
Black, both sexes	32	-	-	-	-	-	2	-	-	-	3	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	2	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Nebraska												
All races, both sexes	131	-	-	-	1	-	-	-	1	-	-	2
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	95	-	-	-	1	-	-	-	1	-	-	2
White, both sexes	128	-	-	-	1	-	-	-	1	-	-	2
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	93	-	-	-	1	-	-	-	1	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	78	-	-	-	-	1	2	-	2	2	3	3
Male	27	-	-	-	-	-	-	-	-	1	2	-
Female	51	-	-	-	-	1	2	-	2	1	1	3
White, both sexes	68	-	-	-	-	-	1	-	1	1	1	3
Male	24	-	-	-	-	-	-	-	-	1	1	-
Female	44	-	-	-	-	-	1	-	1	-	-	3
Black, both sexes	5	-	-	-	-	-	-	-	1	1	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
New Hampshire												
All races, both sexes	66	-	-	-	-	-	-	-	1	-	-	-
Male	23	-	-	-	-	-	-	-	1	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	1	-	-	-
Male	23	-	-	-	-	-	-	-	1	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	409	-	-	1	-	2	2	5	5	12	11	21
Male	109	-	-	-	-	2	-	3	-	3	3	4
Female	300	-	-	1	-	-	2	2	5	9	8	17
White, both sexes	334	-	-	1	-	1	1	2	2	6	6	8
Male	94	-	-	-	-	1	-	2	-	2	2	1
Female	240	-	-	1	-	-	1	-	2	4	4	7
Black, both sexes	67	-	-	-	-	1	1	3	3	5	5	13
Male	14	-	-	-	-	1	-	1	-	1	1	3
Female	53	-	-	-	-	-	1	2	3	4	4	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
New Mexico												
All races, both sexes	121	-	-	-	1	-	3	1	-	1	7	6
Male	41	-	-	-	-	-	1	-	-	-	2	1
Female	80	-	-	-	1	-	2	1	-	1	5	5
White, both sexes	110	-	-	-	1	-	3	1	-	1	6	5
Male	35	-	-	-	-	-	1	-	-	-	2	1
Female	75	-	-	-	1	-	2	1	-	1	4	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	617	-	-	1	1	-	7	6	11	13	23	36
Male	200	-	-	-	-	-	3	-	5	2	9	17
Female	417	-	-	1	1	-	4	6	6	11	14	19
White, both sexes	504	-	-	-	1	-	5	3	5	9	17	23
Male	161	-	-	-	-	-	3	-	2	2	6	14
Female	343	-	-	-	1	-	2	3	3	7	11	9
Black, both sexes	103	-	-	-	-	-	2	2	6	4	6	9
Male	35	-	-	-	-	-	-	-	3	-	3	2
Female	68	-	-	-	-	-	2	2	3	4	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	1	-	-	-	1	-	-	-	4
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	1	-	-	-	1	-	-	-	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
North Carolina												
All races, both sexes	494	-	2	-	-	1	-	3	9	9	16	16
Male	131	-	-	-	-	-	-	-	3	1	4	6
Female	363	-	2	-	-	1	-	3	6	8	12	10
White, both sexes	359	-	1	-	-	-	-	1	5	1	6	9
Male	96	-	-	-	-	-	-	-	2	-	1	5
Female	263	-	1	-	-	-	-	1	3	1	5	4
Black, both sexes	126	-	1	-	-	1	-	2	4	8	9	7
Male	31	-	-	-	-	-	-	-	1	1	3	1
Female	95	-	1	-	-	1	-	2	3	7	6	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Ohio												
All races, both sexes	669	-	-	-	2	2	4	9	7	16	17	17
Male	201	-	-	-	1	2	-	2	1	2	3	6
Female	468	-	-	-	1	-	4	7	6	14	14	11
White, both sexes	584	-	-	-	1	2	1	5	6	6	14	14
Male	183	-	-	-	1	2	-	1	1	1	3	6
Female	401	-	-	-	-	-	1	4	5	5	11	8
Black, both sexes	85	-	-	-	1	-	3	4	1	10	3	3
Male	18	-	-	-	-	-	-	1	-	1	-	-
Female	67	-	-	-	1	-	3	3	1	9	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	201	-	-	-	-	1	2	1	-	2	1	12
Male	67	-	-	-	-	1	1	-	-	-	1	5
Female	134	-	-	-	-	-	1	1	-	2	-	7
White, both sexes	165	-	-	-	-	-	-	1	-	1	1	8
Male	54	-	-	-	-	-	-	-	-	-	1	5
Female	111	-	-	-	-	-	-	1	-	1	-	3
Black, both sexes	15	-	-	-	-	1	2	-	-	1	-	1
Male	6	-	-	-	-	1	1	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	1	-	1
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	3
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Oregon												
All races, both sexes	281	1	-	-	-	1	1	1	-	1	2	5
Male	86	1	-	-	-	1	-	-	-	1	2	3
Female	195	-	-	-	-	-	1	1	-	-	-	2
White, both sexes	268	1	-	-	-	1	1	-	-	1	2	5
Male	81	1	-	-	-	1	-	-	-	1	2	3
Female	187	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	670	-	-	1	2	2	2	5	4	2	11	21
Male	198	-	-	1	1	-	-	-	1	2	1	10
Female	472	-	-	-	1	2	2	5	3	-	10	11
White, both sexes	592	-	-	1	1	1	1	1	-	2	8	11
Male	175	-	-	1	-	-	-	-	-	2	1	6
Female	417	-	-	-	1	1	1	1	-	-	7	5
Black, both sexes	77	-	-	-	1	1	1	4	3	-	3	10
Male	23	-	-	-	1	-	-	-	1	-	-	4
Female	54	-	-	-	-	1	1	4	2	-	3	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Rhode Island												
All races, both sexes	36	-	-	-	-	-	-	-	-	2	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	2	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	225	-	-	-	-	3	2	-	6	7	6	9
Male	67	-	-	-	-	1	-	-	3	2	1	4
Female	158	-	-	-	-	2	2	-	3	5	5	5
White, both sexes	162	-	-	-	-	-	1	-	-	2	2	6
Male	48	-	-	-	-	-	-	-	-	1	1	3
Female	114	-	-	-	-	-	1	-	-	1	1	3
Black, both sexes	62	-	-	-	-	3	1	-	6	5	4	3
Male	18	-	-	-	-	1	-	-	3	1	-	1
Female	44	-	-	-	-	2	1	-	3	4	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
South Dakota												
All races, both sexes	71	-	-	-	-	-	-	-	-	-	2	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	69	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	361	1	-	-	1	-	2	3	3	4	11	15
Male	110	1	-	-	1	-	1	1	1	2	3	4
Female	251	-	-	-	-	-	1	2	2	2	8	11
White, both sexes	300	1	-	-	1	-	1	-	-	3	8	8
Male	89	1	-	-	1	-	1	-	-	1	2	3
Female	211	-	-	-	-	-	-	-	-	2	6	5
Black, both sexes	57	-	-	-	-	-	1	3	3	1	3	7
Male	19	-	-	-	-	-	-	1	1	1	1	1
Female	38	-	-	-	-	-	1	2	2	-	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Texas												
All races, both sexes	979	1	-	-	2	7	9	16	19	30	34	47
Male	294	1	-	-	1	3	6	3	4	10	11	16
Female	685	-	-	-	1	4	3	13	15	20	23	31
White, both sexes	829	1	-	-	2	5	7	12	10	20	25	31
Male	262	1	-	-	1	3	5	2	3	7	9	12
Female	567	-	-	-	1	2	2	10	7	13	16	19
Black, both sexes	133	-	-	-	-	2	2	4	7	10	6	16
Male	30	-	-	-	-	-	1	1	1	3	2	4
Female	103	-	-	-	-	2	1	3	6	7	4	12
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	2	-	3	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	2	-	3	-
Utah												
All races, both sexes	145	-	-	-	-	-	1	-	2	-	1	4
Male	47	-	-	-	-	-	1	-	-	-	-	2
Female	98	-	-	-	-	-	-	-	2	-	1	2
White, both sexes	143	-	-	-	-	-	1	-	2	-	1	4
Male	47	-	-	-	-	-	1	-	-	-	-	2
Female	96	-	-	-	-	-	-	-	2	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Vermont												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	351	1	-	1	1	-	4	3	5	4	9	6
Male	100	1	-	1	1	-	2	1	2	-	3	1
Female	251	-	-	-	-	-	2	2	3	4	6	5
White, both sexes	276	1	-	1	1	-	1	-	2	2	4	4
Male	75	1	-	1	1	-	1	-	1	-	2	1
Female	201	-	-	-	-	-	-	-	1	2	2	3
Black, both sexes	70	-	-	-	-	-	3	3	3	2	5	2
Male	23	-	-	-	-	-	1	1	1	-	1	-
Female	47	-	-	-	-	-	2	2	2	2	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Washington												
All races, both sexes	340	1	-	-	-	1	2	3	4	2	2	7
Male	108	1	-	-	-	1	1	1	2	2	1	2
Female	232	-	-	-	-	-	1	2	2	-	1	5
White, both sexes	315	-	-	-	-	-	1	2	3	2	1	6
Male	102	-	-	-	-	-	1	1	2	2	1	1
Female	213	-	-	-	-	-	-	1	1	-	-	5
Black, both sexes	11	1	-	-	-	1	-	1	1	-	-	1
Male	6	1	-	-	-	1	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	115	-	-	-	-	-	-	1	1	2	4	1
Male	30	-	-	-	-	-	-	-	1	1	1	-
Female	85	-	-	-	-	-	-	1	-	1	3	1
White, both sexes	112	-	-	-	-	-	-	1	1	2	3	1
Male	30	-	-	-	-	-	-	-	1	1	1	-
Female	82	-	-	-	-	-	-	1	-	1	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Wisconsin												
All races, both sexes	283	-	1	-	1	-	-	2	2	1	7	6
Male	86	-	-	-	1	-	-	-	1	-	3	2
Female	197	-	1	-	-	-	-	2	1	1	4	4
White, both sexes	266	-	-	-	-	-	-	1	1	1	5	4
Male	82	-	-	-	-	-	-	-	1	-	2	2
Female	184	-	-	-	-	-	-	1	-	1	3	2
Black, both sexes	17	-	1	-	1	-	-	1	1	-	2	2
Male	4	-	-	-	1	-	-	-	-	-	1	-
Female	13	-	1	-	-	-	-	1	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
United States												
All races, both sexes	639	792	828	967	1,257	1,744	2,275	2,065	1,429	641	165	1
Male	249	283	311	338	440	600	681	548	282	92	17	1
Female	390	509	517	629	817	1,144	1,594	1,517	1,147	549	148	-
White, both sexes	459	584	650	816	1,102	1,565	2,096	1,929	1,354	593	149	-
Male	191	214	242	288	385	534	627	512	273	81	14	-
Female	268	370	408	528	717	1,031	1,469	1,417	1,081	512	135	-
Black, both sexes	150	176	150	123	125	148	146	113	60	36	10	1
Male	46	58	58	42	43	54	41	27	6	8	1	1
Female	104	118	92	81	82	94	105	86	54	28	9	-
American Indian 1/, both sexes	12	15	14	9	12	10	9	3	2	-	-	-
Male	4	7	5	3	4	4	5	-	-	-	-	-
Female	8	8	9	6	8	6	4	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	18	17	14	19	18	21	24	20	13	12	6	-
Male	8	4	6	5	8	8	8	9	3	3	2	-
Female	10	13	8	14	10	13	16	11	10	9	4	-
Alabama												
All races, both sexes	8	13	17	19	19	22	39	24	17	3	-	-
Male	2	6	5	5	8	9	9	5	3	1	-	-
Female	6	7	12	14	11	13	30	19	14	2	-	-
White, both sexes	6	7	10	16	16	19	33	21	13	3	-	-
Male	2	4	3	4	7	8	7	3	2	1	-	-
Female	4	3	7	12	9	11	26	18	11	2	-	-
Black, both sexes	2	6	7	3	3	3	6	3	4	-	-	-
Male	-	2	2	1	1	1	2	2	1	-	-	-
Female	2	4	5	2	2	2	4	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Alaska												
All races, both sexes	3	1	3	3	3	1	2	4	-	1	-	-
Male	1	-	-	2	2	1	-	1	-	-	-	-
Female	2	1	3	1	1	-	2	3	-	1	-	-
White, both sexes	1	1	1	2	-	-	1	3	-	1	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	1	1	-	-	1	2	-	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	1	2	1	1	1	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	2	-	2	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	16	18	16	27	30	36	32	23	19	6	-	-
Male	6	6	4	7	7	15	10	8	4	-	-	-
Female	10	12	12	20	23	21	22	15	15	6	-	-
White, both sexes	13	16	16	26	27	36	30	23	19	6	-	-
Male	5	6	4	6	6	15	9	8	4	-	-	-
Female	8	10	12	20	21	21	21	15	15	6	-	-
Black, both sexes	3	1	-	1	2	-	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	2	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Arkansas												
All races, both sexes	9	10	11	7	13	14	19	10	12	7	1	-
Male	3	2	3	3	5	4	8	3	3	1	-	-
Female	6	8	8	4	8	10	11	7	9	6	1	-
White, both sexes	4	10	10	6	10	12	19	8	12	7	1	-
Male	3	2	3	2	4	4	8	2	3	1	-	-
Female	1	8	7	4	6	8	11	6	9	6	1	-
Black, both sexes	5	-	1	1	3	2	-	2	-	-	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	5	-	1	-	2	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	84	96	82	98	120	171	205	171	107	53	11	-
Male	30	31	29	34	37	55	64	51	21	10	-	-
Female	54	65	53	64	83	116	141	120	86	43	11	-
White, both sexes	63	71	63	81	106	154	179	159	101	46	10	-
Male	23	20	22	28	32	51	51	46	18	7	-	-
Female	40	51	41	53	74	103	128	113	83	39	10	-
Black, both sexes	12	16	12	9	8	9	13	5	1	2	-	-
Male	2	9	5	3	3	2	7	1	-	1	-	-
Female	10	7	7	6	5	7	6	4	1	1	-	-
American Indian 1/, both sexes	2	3	1	1	1	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	3	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	6	6	7	5	8	13	7	5	5	1	-
Male	3	2	1	3	2	2	6	4	3	2	-	-
Female	4	4	5	4	3	6	7	3	2	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Colorado												
All races, both sexes	9	15	13	13	22	24	44	45	33	14	2	-
Male	3	7	6	7	7	5	11	11	8	1	-	-
Female	6	8	7	6	15	19	33	34	25	13	2	-
White, both sexes	8	14	12	12	21	21	42	45	33	14	2	-
Male	3	7	5	6	7	3	11	11	8	1	-	-
Female	5	7	7	6	14	18	31	34	25	13	2	-
Black, both sexes	1	1	-	1	1	2	2	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	1	-	-	1	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	2	6	6	10	25	24	36	27	10	4	-
Male	2	-	2	2	2	9	8	9	6	1	-	-
Female	1	2	4	4	8	16	16	27	21	9	4	-
White, both sexes	2	2	6	5	10	23	21	35	26	10	4	-
Male	1	-	2	1	2	7	7	9	6	1	-	-
Female	1	2	4	4	8	16	14	26	20	9	4	-
Black, both sexes	1	-	-	1	-	2	3	1	1	-	-	-
Male	1	-	-	1	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Delaware												
All races, both sexes	3	5	2	2	2	4	4	4	2	6	1	-
Male	-	2	1	2	-	1	1	2	-	1	-	-
Female	3	3	1	-	2	3	3	2	2	5	1	-
White, both sexes	3	2	2	2	2	3	4	4	2	5	-	-
Male	-	-	1	2	-	1	1	2	-	-	-	-
Female	3	2	1	-	2	2	3	2	2	5	-	-
Black, both sexes	-	3	-	-	-	1	-	-	-	1	1	-
Male	-	2	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	1	2	-	1	4	3	1	-	-	-
Male	-	1	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	3	3	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	2	1	2	-	1	3	2	1	-	-	-
Male	-	1	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Florida												
All races, both sexes	36	50	40	58	71	109	117	100	63	39	6	-
Male	19	16	13	12	27	30	30	32	16	2	1	-
Female	17	34	27	46	44	79	87	68	47	37	5	-
White, both sexes	29	44	34	49	63	98	114	97	59	36	6	-
Male	16	15	11	8	24	25	29	29	15	2	1	-
Female	13	29	23	41	39	73	85	68	44	34	5	-
Black, both sexes	6	6	6	9	8	10	3	3	4	3	-	-
Male	3	1	2	4	3	4	1	3	1	-	-	-
Female	3	5	4	5	5	6	2	-	3	3	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	23	22	21	34	39	39	44	41	20	12	4	-
Male	12	11	11	18	15	14	13	10	5	1	-	-
Female	11	11	10	16	24	25	31	31	15	11	4	-
White, both sexes	12	15	14	25	23	32	33	37	13	11	2	-
Male	6	9	8	15	9	12	9	9	4	1	-	-
Female	6	6	6	10	14	20	24	28	9	10	2	-
Black, both sexes	10	7	6	8	15	7	11	4	7	1	2	-
Male	6	2	2	3	6	2	4	1	1	-	-	-
Female	4	5	4	5	9	5	7	3	6	1	2	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Hawaii												
All races, both sexes	6	6	3	2	3	5	7	7	6	5	3	-
Male	4	3	3	1	1	2	3	2	-	-	1	-
Female	2	3	-	1	2	3	4	5	6	5	2	-
White, both sexes	3	2	-	1	1	1	3	-	1	-	-	-
Male	1	1	-	1	-	-	2	-	-	-	-	-
Female	2	1	-	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	3	-	2	4	4	7	5	5	3	-
Male	3	2	3	-	1	2	1	2	-	-	1	-
Female	-	2	-	-	1	2	3	5	5	5	2	-
Idaho												
All races, both sexes	-	4	2	4	7	11	18	21	12	8	-	-
Male	-	2	-	4	1	5	3	4	4	2	-	-
Female	-	2	2	-	6	6	15	17	8	6	-	-
White, both sexes	-	4	1	4	7	10	18	21	12	8	-	-
Male	-	2	-	4	1	4	3	4	4	2	-	-
Female	-	2	1	-	6	6	15	17	8	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Illinois												
All races, both sexes	22	28	42	43	48	83	111	102	67	33	11	-
Male	8	11	17	13	14	32	36	28	13	4	1	-
Female	14	17	25	30	34	51	75	74	54	29	10	-
White, both sexes	14	20	32	37	38	68	101	97	65	33	10	-
Male	6	8	15	11	11	26	33	27	13	4	1	-
Female	8	12	17	26	27	42	68	70	52	29	9	-
Black, both sexes	8	7	10	6	7	14	9	5	1	-	1	-
Male	2	3	2	2	1	6	2	1	-	-	-	-
Female	6	4	8	4	6	8	7	4	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	1	-	-	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	1	-	-	-
Indiana												
All races, both sexes	10	17	18	20	26	39	46	51	36	10	4	-
Male	9	7	4	11	10	12	11	13	7	1	1	-
Female	1	10	14	9	16	27	35	38	29	9	3	-
White, both sexes	6	13	15	19	25	36	44	50	34	9	4	-
Male	5	5	4	11	9	11	10	13	7	1	1	-
Female	1	8	11	8	16	25	34	37	27	8	3	-
Black, both sexes	4	3	3	1	1	3	2	1	2	1	-	-
Male	4	2	-	-	1	1	1	-	-	-	-	-
Female	-	1	3	1	-	2	1	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Iowa												
All races, both sexes	2	8	12	14	13	18	41	36	36	7	2	-
Male	-	3	4	5	6	6	21	12	12	1	-	-
Female	2	5	8	9	7	12	20	24	24	6	2	-
White, both sexes	2	8	12	14	13	17	41	36	36	7	2	-
Male	-	3	4	5	6	5	21	12	12	1	-	-
Female	2	5	8	9	7	12	20	24	24	6	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	12	11	7	15	21	40	30	19	8	4	-
Male	2	1	5	1	6	6	12	7	6	3	-	-
Female	7	11	6	6	9	15	28	23	13	5	4	-
White, both sexes	7	11	9	4	13	19	39	30	18	8	4	-
Male	2	1	5	1	5	4	12	7	6	3	-	-
Female	5	10	4	3	8	15	27	23	12	5	4	-
Black, both sexes	1	1	1	2	2	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	1	1	2	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Kentucky												
All races, both sexes	11	12	9	20	25	23	40	34	19	9	5	-
Male	7	4	4	10	6	10	7	10	6	3	-	-
Female	4	8	5	10	19	13	33	24	13	6	5	-
White, both sexes	8	12	9	17	25	22	39	34	19	8	4	-
Male	5	4	4	9	6	10	7	10	6	3	-	-
Female	3	8	5	8	19	12	32	24	13	5	4	-
Black, both sexes	3	-	-	3	-	1	1	-	-	1	1	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	2	-	1	1	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	10	13	13	8	29	25	31	27	20	7	2	-
Male	2	5	4	2	14	8	8	5	6	2	-	-
Female	8	8	9	6	15	17	23	22	14	5	2	-
White, both sexes	2	5	8	5	17	18	24	18	15	5	2	-
Male	1	3	3	1	11	7	7	5	6	1	-	-
Female	1	2	5	4	6	11	17	13	9	4	2	-
Black, both sexes	8	8	5	3	12	7	7	9	5	2	-	-
Male	1	2	1	1	3	1	1	-	-	1	-	-
Female	7	6	4	2	9	6	6	9	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Maine												
All races, both sexes	3	4	4	9	8	11	21	21	10	9	4	-
Male	3	1	3	-	3	4	11	4	2	2	-	-
Female	-	3	1	9	5	7	10	17	8	7	4	-
White, both sexes	3	4	4	9	8	10	21	21	10	9	4	-
Male	3	1	3	-	3	4	11	4	2	2	-	-
Female	-	3	1	9	5	6	10	17	8	7	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	15	19	21	12	22	35	40	56	28	12	2	-
Male	7	1	7	5	4	13	7	11	5	1	1	-
Female	8	18	14	7	18	22	33	45	23	11	1	-
White, both sexes	4	7	10	8	17	30	36	45	28	12	1	-
Male	1	-	4	4	3	11	7	6	5	1	-	-
Female	3	7	6	4	14	19	29	39	23	11	1	-
Black, both sexes	11	12	10	4	5	5	3	10	-	-	1	-
Male	6	1	3	1	1	2	-	5	-	-	1	-
Female	5	11	7	3	4	3	3	5	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Massachusetts												
All races, both sexes	5	13	11	12	17	38	53	40	35	14	4	-
Male	-	5	3	1	6	17	18	10	8	1	1	-
Female	5	8	8	11	11	21	35	30	27	13	3	-
White, both sexes	5	11	9	12	15	38	52	40	35	13	3	-
Male	-	5	3	1	6	17	18	10	8	1	-	-
Female	5	6	6	11	9	21	34	30	27	12	3	-
Black, both sexes	-	2	1	-	2	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	2	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	14	39	33	32	47	56	93	86	40	27	6	-
Male	6	15	13	12	19	23	33	19	12	3	3	-
Female	8	24	20	20	28	33	60	67	28	24	3	-
White, both sexes	8	26	27	29	45	50	87	82	36	26	6	-
Male	3	10	11	11	19	22	31	18	11	2	3	-
Female	5	16	16	18	26	28	56	64	25	24	3	-
Black, both sexes	5	12	6	3	2	6	6	3	4	1	-	-
Male	3	5	2	1	-	1	2	-	1	1	-	-
Female	2	7	4	2	2	5	4	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Minnesota												
All races, both sexes	4	12	11	17	25	44	55	58	55	28	9	-
Male	2	5	3	7	9	16	18	13	5	7	1	-
Female	2	7	8	10	16	28	37	45	50	21	8	-
White, both sexes	3	10	10	14	25	44	54	58	55	28	9	-
Male	2	5	2	6	9	16	18	13	5	7	1	-
Female	1	5	8	8	16	28	36	45	50	21	8	-
Black, both sexes	1	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	7	9	6	12	11	16	10	4	4	-	-
Male	-	2	2	4	9	1	6	4	-	1	-	-
Female	6	5	7	2	3	10	10	6	4	3	-	-
White, both sexes	2	3	5	5	10	6	13	8	4	1	-	-
Male	-	-	-	3	7	1	5	3	-	-	-	-
Female	2	3	5	2	3	5	8	5	4	1	-	-
Black, both sexes	4	4	4	1	2	5	3	2	-	3	-	-
Male	-	2	2	1	2	-	1	1	-	1	-	-
Female	4	2	2	-	-	5	2	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Missouri												
All races, both sexes	6	14	16	24	29	43	64	39	36	18	3	-
Male	3	3	8	11	15	16	16	7	6	3	2	-
Female	3	11	8	13	14	27	48	32	30	15	1	-
White, both sexes	6	10	12	17	26	39	63	36	35	16	3	-
Male	3	2	5	7	13	15	16	7	6	2	2	-
Female	3	8	7	10	13	24	47	29	29	14	1	-
Black, both sexes	-	3	4	6	3	4	1	2	1	2	-	-
Male	-	1	3	4	2	1	-	-	-	1	-	-
Female	-	2	1	2	1	3	1	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	2	2	2	2	5	11	13	15	22	3	2	-
Male	-	2	-	1	-	4	4	4	4	1	-	-
Female	2	-	2	1	5	7	9	11	18	2	2	-
White, both sexes	2	2	1	2	4	11	13	15	22	3	2	-
Male	-	2	-	1	-	4	4	4	4	1	-	-
Female	2	-	1	1	4	7	9	11	18	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Nebraska												
All races, both sexes	4	2	4	6	11	20	19	24	22	10	5	-
Male	-	1	-	3	5	7	6	7	5	2	-	-
Female	4	1	4	3	6	13	13	17	17	8	5	-
White, both sexes	3	2	4	6	10	20	19	23	22	10	5	-
Male	-	1	-	3	5	7	6	6	5	2	-	-
Female	3	1	4	3	5	13	13	17	17	8	5	-
Black, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	4	1	10	10	11	10	10	4	-	-	-
Male	3	1	1	4	5	3	3	3	1	-	-	-
Female	2	3	-	6	5	8	7	7	3	-	-	-
White, both sexes	5	2	1	10	9	10	10	10	4	-	-	-
Male	3	1	1	4	4	2	3	3	1	-	-	-
Female	2	1	-	6	5	8	7	7	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
New Hampshire												
All races, both sexes	3	6	2	4	6	12	12	6	11	3	-	-
Male	1	2	-	1	3	7	4	3	1	-	-	-
Female	2	4	2	3	3	5	8	3	10	3	-	-
White, both sexes	2	6	2	4	6	12	12	6	11	3	-	-
Male	1	2	-	1	3	7	4	3	1	-	-	-
Female	1	4	2	3	3	5	8	3	10	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	16	25	27	24	35	47	61	54	34	24	3	-
Male	6	6	9	4	8	14	18	19	9	1	-	-
Female	10	19	18	20	27	33	43	35	25	23	3	-
White, both sexes	11	12	25	18	34	44	53	52	32	23	3	-
Male	5	4	8	3	8	14	17	17	9	1	-	-
Female	6	8	17	15	26	30	36	35	23	22	3	-
Black, both sexes	5	12	2	4	-	3	7	2	1	-	-	-
Male	1	2	1	1	-	-	-	2	-	-	-	-
Female	4	10	1	3	-	3	7	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	2	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
New Mexico												
All races, both sexes	9	5	6	10	11	13	16	14	13	5	-	-
Male	4	3	3	4	2	6	8	4	3	-	-	-
Female	5	2	3	6	9	7	8	10	10	5	-	-
White, both sexes	7	3	5	9	10	13	15	14	12	5	-	-
Male	3	1	2	3	1	6	8	4	3	-	-	-
Female	4	2	3	6	9	7	7	10	9	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	2	1	1	1	-	1	-	1	-	-	-
Male	1	2	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	37	35	28	52	50	75	85	63	59	30	5	-
Male	13	13	13	19	19	30	31	13	9	3	1	-
Female	24	22	15	33	31	45	54	50	50	27	4	-
White, both sexes	24	28	19	41	38	68	77	60	55	26	5	-
Male	11	9	8	15	13	26	28	13	8	2	1	-
Female	13	19	11	26	25	42	49	47	47	24	4	-
Black, both sexes	13	7	8	9	11	7	8	3	4	4	-	-
Male	2	4	4	3	5	4	3	-	1	1	-	-
Female	11	3	4	6	6	3	5	3	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
North Carolina												
All races, both sexes	22	32	26	24	42	51	98	78	43	15	7	-
Male	9	7	14	5	16	12	28	14	7	4	1	-
Female	13	25	12	19	26	39	70	64	36	11	6	-
White, both sexes	14	17	16	18	33	41	77	65	36	13	6	-
Male	6	2	10	3	13	9	23	11	7	3	1	-
Female	8	15	6	15	20	32	54	54	29	10	5	-
Black, both sexes	7	13	10	6	8	10	18	13	7	1	1	-
Male	3	3	4	2	3	3	4	3	-	-	-	-
Female	4	10	6	4	5	7	14	10	7	1	1	-
American Indian 1/, both sexes	1	2	-	-	1	-	1	-	-	-	-	-
Male	-	2	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	2	3	-	5	9	8	8	4	-	-
Male	-	1	2	2	-	3	2	4	1	1	-	-
Female	2	1	-	1	-	2	7	4	7	3	-	-
White, both sexes	2	2	2	3	-	5	8	8	8	4	-	-
Male	-	1	2	2	-	3	2	4	1	1	-	-
Female	2	1	-	1	-	2	6	4	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Ohio												
All races, both sexes	23	32	31	39	54	89	93	113	77	31	13	-
Male	7	20	11	11	27	37	24	27	14	5	1	-
Female	16	12	20	28	27	52	69	86	63	26	12	-
White, both sexes	20	26	22	35	47	77	88	105	74	29	12	-
Male	7	16	8	11	25	32	23	27	13	5	1	-
Female	13	10	14	24	22	45	65	78	61	24	11	-
Black, both sexes	3	6	9	4	7	12	5	8	3	2	1	-
Male	-	4	3	-	2	5	1	-	1	-	-	-
Female	3	2	6	4	5	7	4	8	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	8	17	22	15	22	34	29	17	13	4	1	-
Male	4	7	6	7	10	12	9	3	1	-	-	-
Female	4	10	16	8	12	22	20	14	12	4	1	-
White, both sexes	6	13	14	12	20	29	27	17	12	4	-	-
Male	3	6	4	4	9	10	8	3	1	-	-	-
Female	3	7	10	8	11	19	19	14	11	4	-	-
Black, both sexes	-	1	4	2	-	1	-	-	1	-	1	-
Male	-	-	1	2	-	1	-	-	-	-	-	-
Female	-	1	3	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	2	3	4	1	2	4	2	-	-	-	-	-
Male	1	1	1	1	1	1	1	-	-	-	-	-
Female	1	2	3	-	1	3	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Oregon												
All races, both sexes	5	15	13	17	23	34	50	49	43	15	4	1
Male	3	10	5	7	8	12	15	11	6	-	-	1
Female	2	5	8	10	15	22	35	38	37	15	4	-
White, both sexes	5	13	11	16	21	33	49	47	43	15	4	-
Male	3	9	4	7	7	12	15	10	6	-	-	-
Female	2	4	7	9	14	21	34	37	37	15	4	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	1
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	-	1	1	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	37	28	46	41	67	81	112	109	64	29	6	-
Male	14	11	14	17	21	26	28	37	13	1	-	-
Female	23	17	32	24	46	55	84	72	51	28	6	-
White, both sexes	32	18	40	35	63	75	106	103	63	25	6	-
Male	13	8	13	16	19	22	26	34	13	1	-	-
Female	19	10	27	19	44	53	80	69	50	24	6	-
Black, both sexes	5	10	6	6	4	6	6	6	1	4	-	-
Male	1	3	1	1	2	4	2	3	-	-	-	-
Female	4	7	5	5	2	2	4	3	1	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Rhode Island												
All races, both sexes	1	-	-	4	3	7	10	2	5	1	1	-
Male	-	-	-	2	2	2	2	1	2	-	-	-
Female	1	-	-	2	1	5	8	1	3	1	1	-
White, both sexes	1	-	-	4	3	7	10	2	5	1	1	-
Male	-	-	-	2	2	2	2	1	2	-	-	-
Female	1	-	-	2	1	5	8	1	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	13	18	15	14	24	26	34	23	20	3	2	-
Male	4	7	3	2	10	11	8	5	6	-	-	-
Female	9	11	12	12	14	15	26	18	14	3	2	-
White, both sexes	8	8	10	11	21	22	29	17	20	3	2	-
Male	2	2	3	2	8	9	7	4	6	-	-	-
Female	6	6	7	9	13	13	22	13	14	3	2	-
Black, both sexes	5	9	5	3	3	4	5	6	-	-	-	-
Male	2	4	-	-	2	2	1	1	-	-	-	-
Female	3	5	5	3	1	2	4	5	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
South Dakota												
All races, both sexes	2	1	-	5	4	9	11	11	14	10	2	-
Male	-	1	-	3	1	4	2	6	4	-	1	-
Female	2	-	-	2	3	5	9	5	10	10	1	-
White, both sexes	2	1	-	5	4	8	11	11	14	10	2	-
Male	-	1	-	3	1	4	2	6	4	-	1	-
Female	2	-	-	2	3	4	9	5	10	10	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	22	21	18	37	37	53	49	53	18	9	4	-
Male	9	8	8	15	11	12	15	15	1	2	-	-
Female	13	13	10	22	26	41	34	38	17	7	4	-
White, both sexes	14	17	15	29	35	45	45	50	17	8	3	-
Male	6	6	7	11	11	8	14	14	1	2	-	-
Female	8	11	8	18	24	37	31	36	16	6	3	-
Black, both sexes	7	4	3	8	1	8	3	2	1	1	1	-
Male	2	2	1	4	-	4	1	-	-	-	-	-
Female	5	2	2	4	1	4	2	2	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Texas												
All races, both sexes	55	59	81	76	86	103	129	118	71	32	4	-
Male	16	18	34	24	27	32	39	31	12	6	-	-
Female	39	41	47	52	59	71	90	87	59	26	4	-
White, both sexes	41	47	65	67	73	95	118	109	68	29	4	-
Male	14	17	25	24	25	31	34	31	12	6	-	-
Female	27	30	40	43	48	64	84	78	56	23	4	-
Black, both sexes	13	12	15	6	11	6	10	8	2	3	-	-
Male	2	1	9	-	1	1	4	-	-	-	-	-
Female	11	11	6	6	10	5	6	8	2	3	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	2	1	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	2	1	1	-	-	1	-	-	-
Utah												
All races, both sexes	6	4	10	8	13	18	26	32	14	6	-	-
Male	2	2	4	5	4	4	9	10	2	2	-	-
Female	4	2	6	3	9	14	17	22	12	4	-	-
White, both sexes	5	4	10	8	13	17	26	32	14	6	-	-
Male	2	2	4	5	4	4	9	10	2	2	-	-
Female	3	2	6	3	9	13	17	22	12	4	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Vermont												
All races, both sexes	-	3	1	2	4	7	5	8	12	1	1	-
Male	-	-	1	-	1	4	-	1	-	-	-	-
Female	-	3	-	2	3	3	5	7	12	1	1	-
White, both sexes	-	3	1	2	4	7	5	8	12	1	1	-
Male	-	-	1	-	1	4	-	1	-	-	-	-
Female	-	3	-	2	3	3	5	7	12	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	14	12	26	32	29	39	59	42	44	16	4	-
Male	7	3	15	10	6	14	17	12	2	2	-	-
Female	7	9	11	22	23	25	42	30	42	14	4	-
White, both sexes	10	7	18	22	28	33	52	35	38	13	4	-
Male	4	2	9	8	5	10	15	11	2	1	-	-
Female	6	5	9	14	23	23	37	24	36	12	4	-
Black, both sexes	3	5	8	8	1	4	7	7	6	3	-	-
Male	2	1	6	2	1	3	2	1	-	1	-	-
Female	1	4	2	6	-	1	5	6	6	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Washington												
All races, both sexes	16	18	19	18	34	36	52	60	41	17	7	-
Male	6	6	7	3	11	10	22	18	10	3	1	-
Female	10	12	12	15	23	26	30	42	31	14	6	-
White, both sexes	12	16	18	16	31	34	52	58	40	17	6	-
Male	6	5	6	3	11	10	22	17	10	3	1	-
Female	6	11	12	13	20	24	30	41	30	14	5	-
Black, both sexes	1	1	1	-	1	1	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	2	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	2	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	4	4	5	4	13	21	23	19	9	3	1	-
Male	1	3	2	3	2	9	4	2	1	-	-	-
Female	3	1	3	1	11	12	19	17	8	3	1	-
White, both sexes	4	4	5	4	13	21	23	18	8	3	1	-
Male	1	3	2	3	2	9	4	2	1	-	-	-
Female	3	1	3	1	11	12	19	16	7	3	1	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Wisconsin												
All races, both sexes	14	4	13	18	19	26	53	53	40	18	5	-
Male	7	1	4	3	8	10	17	15	8	6	-	-
Female	7	3	9	15	11	16	36	38	32	12	5	-
White, both sexes	13	4	12	17	19	25	52	50	39	18	5	-
Male	7	1	3	3	8	9	17	15	8	6	-	-
Female	6	3	9	14	11	16	35	35	31	12	5	-
Black, both sexes	1	-	1	1	-	1	1	3	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	1	3	3	-	7	7	5	4	2	-	-
Male	2	-	-	3	-	1	1	2	2	1	-	-
Female	-	1	3	-	-	6	6	3	2	1	-	-
White, both sexes	2	1	3	3	-	7	7	5	4	2	-	-
Male	2	-	-	3	-	1	1	2	2	1	-	-
Female	-	1	3	-	-	6	6	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
United States												
All races, both sexes	2,538	-	1	-	1	-	3	5	5	8	18	32
Male	596	-	-	-	-	-	-	2	2	2	3	10
Female	1,942	-	1	-	1	-	3	3	3	6	15	22
White, both sexes	2,330	-	-	-	1	-	1	4	5	5	16	29
Male	554	-	-	-	-	-	-	2	2	2	3	10
Female	1,776	-	-	-	1	-	1	2	3	3	13	19
Black, both sexes	155	-	1	-	-	-	1	1	-	3	2	3
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	127	-	1	-	-	-	1	1	-	3	2	3
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	1	-	-	-	-	-
Alabama												
All races, both sexes	54	-	-	-	-	-	-	-	-	2	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	2	1	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	-	-	-	-	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	53	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Arkansas												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	231	-	-	-	-	-	1	-	2	-	2	3
Male	48	-	-	-	-	-	-	-	1	-	-	2
Female	183	-	-	-	-	-	1	-	1	-	2	1
White, both sexes	207	-	-	-	-	-	-	-	2	-	2	3
Male	41	-	-	-	-	-	-	-	1	-	-	2
Female	166	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Colorado												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Florida												
All races, both sexes	123	-	-	-	-	-	-	-	-	-	1	3
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	94	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	3
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	89	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Hawaii												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	1	-
Male	6	-	-	-	-	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Illinois												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	87	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	100	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	81	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	67	-	-	-	-	-	-	2	-	-	-	1
Male	13	-	-	-	-	-	-	1	-	-	-	-
Female	54	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	63	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Iowa												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Kentucky												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	36	-	-	-	-	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	49	-	-	-	-	-	-	-	1	1	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	45	-	-	-	-	-	-	-	1	1	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Massachusetts												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	103	-	-	-	-	-	-	-	-	-	3	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	75	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	99	-	-	-	-	-	-	-	-	-	3	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Minnesota												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Missouri												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	2	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	44	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	2	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	44	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Nebraska												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	51	-	-	-	-	-	-	1	-	-	-	2
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	49	-	-	-	-	-	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
New Mexico												
All races, both sexes	24	-	-	-	1	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	1	-	-	-	-	-	1	1
White, both sexes	23	-	-	-	1	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	1	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	67	-	-	-	-	-	-	-	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
North Carolina												
All races, both sexes	105	-	1	-	-	-	-	-	-	-	1	3
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	80	-	1	-	-	-	-	-	-	-	1	2
White, both sexes	85	-	-	-	-	-	-	-	-	-	1	3
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	67	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	16	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Ohio												
All races, both sexes	112	-	-	-	-	-	-	-	-	-	2	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	86	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	104	-	-	-	-	-	-	-	-	-	2	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	78	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Oregon												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	113	-	-	-	-	-	-	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	95	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	109	-	-	-	-	-	-	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	91	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	43	-	-	-	-	-	1	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
South Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Texas												
All races, both sexes	173	-	-	-	-	-	1	2	-	-	1	2
Male	42	-	-	-	-	-	-	-	-	-	1	-
Female	131	-	-	-	-	-	1	2	-	-	-	2
White, both sexes	159	-	-	-	-	-	1	2	-	-	1	1
Male	41	-	-	-	-	-	-	-	-	-	1	-
Female	118	-	-	-	-	-	1	2	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	28	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Vermont												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	62	-	-	-	-	-	-	-	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	1	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Washington												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Wisconsin												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
United States												
All races, both sexes	64	109	136	208	315	485	562	370	173	40	3	-
Male	23	35	36	55	95	131	116	63	19	4	-	-
Female	41	74	100	153	220	354	446	307	154	36	3	-
White, both sexes	55	96	108	195	287	448	532	344	162	39	3	-
Male	21	31	30	54	89	122	106	60	18	4	-	-
Female	34	65	78	141	198	326	426	284	144	35	3	-
Black, both sexes	8	6	22	11	23	25	18	22	8	1	-	-
Male	2	2	5	-	5	5	7	2	-	-	-	-
Female	6	4	17	11	18	20	11	20	8	1	-	-
American Indian 1/, both sexes	-	4	5	1	3	6	3	1	1	-	-	-
Male	-	2	1	1	1	3	1	-	-	-	-	-
Female	-	2	4	-	2	3	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	1	2	6	9	3	2	-	-	-
Male	-	-	-	-	-	1	2	1	1	-	-	-
Female	1	3	1	1	2	5	7	2	1	-	-	-
Alabama												
All races, both sexes	2	2	4	8	4	10	11	9	1	-	-	-
Male	1	1	1	3	-	3	2	2	-	-	-	-
Female	1	1	3	5	4	7	9	7	1	-	-	-
White, both sexes	2	2	2	8	3	8	10	8	-	-	-	-
Male	1	1	1	3	-	2	2	2	-	-	-	-
Female	1	1	1	5	3	6	8	6	-	-	-	-
Black, both sexes	-	-	2	-	1	2	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Alaska												
All races, both sexes	-	1	2	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	1	2	4	16	12	8	1	5	-	-	-
Male	1	-	-	1	3	6	1	1	1	-	-	-
Female	2	1	2	3	13	6	7	-	4	-	-	-
White, both sexes	3	1	2	4	15	12	8	1	5	-	-	-
Male	1	-	-	1	3	6	1	1	1	-	-	-
Female	2	1	2	3	12	6	7	-	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Arkansas												
All races, both sexes	-	3	3	-	5	3	5	2	1	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	-	3	2	-	4	2	5	1	1	-	-	-
White, both sexes	-	3	2	-	4	3	5	1	1	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	3	1	-	3	2	5	1	1	-	-	-
Black, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10	14	12	13	26	44	52	31	17	4	-	-
Male	3	6	1	2	7	8	13	4	1	-	-	-
Female	7	8	11	11	19	36	39	27	16	4	-	-
White, both sexes	9	13	10	13	25	38	46	28	14	4	-	-
Male	3	6	-	2	7	7	10	3	-	-	-	-
Female	6	7	10	11	18	31	36	25	14	4	-	-
Black, both sexes	1	-	2	-	1	4	2	2	1	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	1	-	1	-	1	4	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	4	1	2	-	-	-
Male	-	-	-	-	-	1	2	-	1	-	-	-
Female	-	1	-	-	-	1	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Colorado												
All races, both sexes	-	2	5	2	7	6	11	8	1	1	1	-
Male	-	-	2	1	4	-	3	2	1	-	-	-
Female	-	2	3	1	3	6	8	6	-	1	1	-
White, both sexes	-	1	4	2	7	5	10	8	1	1	1	-
Male	-	-	1	1	4	-	3	2	1	-	-	-
Female	-	1	3	1	3	5	7	6	-	1	1	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	1	4	4	3	7	1	-	1	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	1	3	4	2	7	1	-	1	-
White, both sexes	-	-	1	1	4	4	2	6	1	-	1	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	1	1	3	4	1	6	1	-	1	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Delaware												
All races, both sexes	-	1	1	1	-	-	2	2	1	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	2	1	1	-	-	-
White, both sexes	-	1	1	1	-	-	2	2	1	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Florida												
All races, both sexes	3	3	5	11	18	30	22	17	8	2	-	-
Male	2	1	2	2	5	5	6	5	1	-	-	-
Female	1	2	3	9	13	25	16	12	7	2	-	-
White, both sexes	3	3	4	11	15	28	21	17	8	2	-	-
Male	2	1	1	2	4	5	5	5	1	-	-	-
Female	1	2	3	9	11	23	16	12	7	2	-	-
Black, both sexes	-	-	1	-	3	2	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	2	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	3	4	8	9	11	14	5	2	1	-	-
Male	1	1	1	2	3	4	4	-	-	-	-	-
Female	1	2	3	6	6	7	10	5	2	1	-	-
White, both sexes	1	3	1	7	5	11	13	4	2	1	-	-
Male	-	1	-	2	1	4	3	-	-	-	-	-
Female	1	2	1	5	4	7	10	4	2	1	-	-
Black, both sexes	1	-	3	1	4	-	1	1	-	-	-	-
Male	1	-	1	-	2	-	1	-	-	-	-	-
Female	-	-	2	1	2	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Hawaii												
All races, both sexes	1	1	-	-	1	2	4	2	1	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	-	1	2	3	1	1	-	-	-
White, both sexes	1	-	-	-	-	1	2	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	1	1	2	1	-	-	-	-
Idaho												
All races, both sexes	-	-	-	1	1	3	6	2	3	-	-	-
Male	-	-	-	1	-	1	1	-	1	-	-	-
Female	-	-	-	-	1	2	5	2	2	-	-	-
White, both sexes	-	-	-	1	1	2	6	2	3	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	2	5	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Illinois												
All races, both sexes	2	3	4	9	8	19	27	22	10	1	-	-
Male	-	-	2	2	1	3	3	8	-	-	-	-
Female	2	3	2	7	7	16	24	14	10	1	-	-
White, both sexes	1	3	3	9	7	18	26	22	9	1	-	-
Male	-	-	2	2	1	3	3	8	-	-	-	-
Female	1	3	1	7	6	15	23	14	9	1	-	-
Black, both sexes	1	-	1	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	1	4	5	4	10	10	9	13	8	-	-	-
Male	-	2	2	1	1	1	2	2	1	-	-	-
Female	1	2	3	3	9	9	7	11	7	-	-	-
White, both sexes	1	4	4	4	10	8	9	13	8	-	-	-
Male	-	2	2	1	1	-	2	2	1	-	-	-
Female	1	2	2	3	9	8	7	11	7	-	-	-
Black, both sexes	-	-	1	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Iowa												
All races, both sexes	-	2	5	1	4	8	12	3	4	-	-	-
Male	-	1	2	-	2	3	3	1	-	-	-	-
Female	-	1	3	1	2	5	9	2	4	-	-	-
White, both sexes	-	2	5	1	4	7	12	3	4	-	-	-
Male	-	1	2	-	2	2	3	1	-	-	-	-
Female	-	1	3	1	2	5	9	2	4	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	5	2	-	6	5	15	5	3	1	-	-
Male	1	-	-	-	3	1	2	1	2	-	-	-
Female	-	5	2	-	3	4	13	4	1	1	-	-
White, both sexes	1	5	1	-	6	4	15	5	3	1	-	-
Male	1	-	-	-	3	-	2	1	2	-	-	-
Female	-	5	1	-	3	4	13	4	1	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Kentucky												
All races, both sexes	1	1	2	6	8	6	8	6	3	1	-	-
Male	1	-	-	3	2	1	1	2	1	-	-	-
Female	-	1	2	3	6	5	7	4	2	1	-	-
White, both sexes	1	1	2	6	8	6	8	6	3	1	-	-
Male	1	-	-	3	2	1	1	2	1	-	-	-
Female	-	1	2	3	6	5	7	4	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	3	2	4	7	4	9	2	-	-	-
Male	-	-	2	-	1	1	1	-	-	-	-	-
Female	2	-	1	2	3	6	3	9	2	-	-	-
White, both sexes	-	-	1	1	3	4	4	5	-	-	-	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	1	2	3	3	5	-	-	-	-
Black, both sexes	2	-	2	1	1	3	-	4	2	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	1	1	1	3	-	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Maine												
All races, both sexes	-	1	1	3	4	3	5	4	-	1	-	-
Male	-	-	1	-	-	2	2	-	-	-	-	-
Female	-	1	-	3	4	1	3	4	-	1	-	-
White, both sexes	-	1	1	3	4	3	5	4	-	1	-	-
Male	-	-	1	-	-	2	2	-	-	-	-	-
Female	-	1	-	3	4	1	3	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	2	3	4	10	13	8	3	-	-	-
Male	-	-	-	2	1	3	2	-	1	-	-	-
Female	1	2	2	1	3	7	11	8	2	-	-	-
White, both sexes	1	1	1	3	3	10	13	7	3	-	-	-
Male	-	-	-	2	1	3	2	-	1	-	-	-
Female	1	1	1	1	2	7	11	7	2	-	-	-
Black, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Massachusetts												
All races, both sexes	1	-	1	3	3	10	13	7	4	1	-	-
Male	-	-	-	-	-	4	4	1	-	-	-	-
Female	1	-	1	3	3	6	9	6	4	1	-	-
White, both sexes	1	-	1	3	2	10	13	7	4	1	-	-
Male	-	-	-	-	-	4	4	1	-	-	-	-
Female	1	-	1	3	2	6	9	6	4	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	4	6	7	12	21	26	13	5	3	-	-
Male	1	-	1	2	5	8	5	4	1	-	-	-
Female	1	4	5	5	7	13	21	9	4	3	-	-
White, both sexes	1	4	5	7	12	20	25	13	5	3	-	-
Male	1	-	1	2	5	8	5	4	1	-	-	-
Female	-	4	4	5	7	12	20	9	4	3	-	-
Black, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Minnesota												
All races, both sexes	-	1	2	4	11	15	15	16	7	2	-	-
Male	-	-	1	1	3	5	1	4	-	-	-	-
Female	-	1	1	3	8	10	14	12	7	2	-	-
White, both sexes	-	-	2	3	11	15	15	16	7	2	-	-
Male	-	-	1	1	3	5	1	4	-	-	-	-
Female	-	-	1	2	8	10	14	12	7	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	3	-	3	1	2	1	-	-	-	-
Male	-	-	-	-	3	-	-	-	-	-	-	-
Female	-	1	3	-	-	1	2	1	-	-	-	-
White, both sexes	-	1	2	-	3	1	2	-	-	-	-	-
Male	-	-	-	-	3	-	-	-	-	-	-	-
Female	-	1	2	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Missouri												
All races, both sexes	1	1	1	8	6	15	11	7	5	2	-	-
Male	-	-	-	3	3	3	3	1	1	-	-	-
Female	1	1	1	5	3	12	8	6	4	2	-	-
White, both sexes	1	1	1	8	6	15	11	7	5	2	-	-
Male	-	-	-	3	3	3	3	1	1	-	-	-
Female	1	1	1	5	3	12	8	6	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	2	2	-	2	3	6	3	-	-	-	-
Male	-	2	-	-	-	2	2	2	-	-	-	-
Female	1	-	2	-	2	1	4	1	-	-	-	-
White, both sexes	1	2	1	-	1	3	6	3	-	-	-	-
Male	-	2	-	-	-	2	2	2	-	-	-	-
Female	1	-	1	-	1	1	4	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Nebraska												
All races, both sexes	1	-	1	1	4	6	2	3	3	-	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	1	-	1	1	2	4	2	3	3	-	-	-
White, both sexes	1	-	1	1	4	6	2	3	3	-	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	1	-	1	1	2	4	2	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	2	3	3	1	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	2	3	1	2	-	-	-	-
White, both sexes	-	1	-	2	3	3	1	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	2	3	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
New Hampshire												
All races, both sexes	-	1	1	1	-	3	5	-	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	1	1	-	2	4	-	2	-	-	-
White, both sexes	-	1	1	1	-	3	5	-	2	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	1	1	-	2	4	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	3	4	6	12	12	5	5	-	-	-
Male	-	-	2	-	2	1	-	-	1	-	-	-
Female	-	1	1	4	4	11	12	5	4	-	-	-
White, both sexes	-	1	3	4	6	12	11	5	5	-	-	-
Male	-	-	2	-	2	1	-	-	1	-	-	-
Female	-	1	1	4	4	11	11	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
New Mexico												
All races, both sexes	2	2	1	1	2	6	3	4	-	-	-	-
Male	1	1	1	1	-	3	1	-	-	-	-	-
Female	1	1	-	-	2	3	2	4	-	-	-	-
White, both sexes	2	2	1	1	2	6	2	4	-	-	-	-
Male	1	1	1	1	-	3	1	-	-	-	-	-
Female	1	1	-	-	2	3	1	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	5	-	9	7	14	11	11	5	3	-	-
Male	1	1	-	3	3	5	1	1	1	-	-	-
Female	-	4	-	6	4	9	10	10	4	3	-	-
White, both sexes	-	4	-	9	6	13	11	11	5	3	-	-
Male	-	1	-	3	2	4	1	1	1	-	-	-
Female	-	3	-	6	4	9	10	10	4	3	-	-
Black, both sexes	1	1	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
North Carolina												
All races, both sexes	2	4	2	4	11	15	37	18	6	1	-	-
Male	1	3	1	-	4	1	12	1	-	1	-	-
Female	1	1	1	4	7	14	25	17	6	-	-	-
White, both sexes	2	1	2	3	10	12	29	17	4	1	-	-
Male	1	-	1	-	4	-	9	1	-	1	-	-
Female	1	1	1	3	6	12	20	16	4	-	-	-
Black, both sexes	-	1	-	1	1	3	6	1	2	-	-	-
Male	-	1	-	-	-	1	3	-	-	-	-	-
Female	-	-	-	1	1	2	3	1	2	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	2	-	3	2	1	1	-	-	-
Male	-	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	1	1	-	-	-
White, both sexes	-	1	-	2	-	3	2	1	1	-	-	-
Male	-	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Ohio												
All races, both sexes	2	2	6	11	12	20	20	24	9	2	1	-
Male	1	2	1	2	4	8	3	3	-	1	-	-
Female	1	-	5	9	8	12	17	21	9	1	1	-
White, both sexes	2	2	6	9	11	18	19	22	9	2	1	-
Male	1	2	1	2	4	8	3	3	-	1	-	-
Female	1	-	5	7	7	10	16	19	9	1	1	-
Black, both sexes	-	-	-	2	1	2	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	2	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	4	5	4	3	8	9	-	2	1	-	-
Male	-	1	-	1	2	3	2	-	-	-	-	-
Female	-	3	5	3	1	5	7	-	2	1	-	-
White, both sexes	-	2	2	4	2	7	7	-	2	1	-	-
Male	-	1	-	1	1	3	1	-	-	-	-	-
Female	-	1	2	3	1	4	6	-	2	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	2	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	2	2	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Oregon												
All races, both sexes	2	3	3	4	8	9	13	7	5	1	-	-
Male	-	2	1	1	2	5	4	-	1	-	-	-
Female	2	1	2	3	6	4	9	7	4	1	-	-
White, both sexes	2	3	3	4	8	9	13	7	5	1	-	-
Male	-	2	1	1	2	5	4	-	1	-	-	-
Female	2	1	2	3	6	4	9	7	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	4	8	9	13	19	29	16	8	3	-	-
Male	-	2	3	1	1	4	4	2	-	-	-	-
Female	2	2	5	8	12	15	25	14	8	3	-	-
White, both sexes	2	4	7	8	13	18	29	15	8	3	-	-
Male	-	2	3	1	1	4	4	2	-	-	-	-
Female	2	2	4	7	12	14	25	13	8	3	-	-
Black, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Rhode Island												
All races, both sexes	-	-	-	1	-	-	4	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	4	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	3	7	7	9	5	6	3	-	-	-
Male	-	-	1	1	5	1	1	2	1	-	-	-
Female	2	-	2	6	2	8	4	4	2	-	-	-
White, both sexes	2	-	2	6	6	9	5	5	3	-	-	-
Male	-	-	1	1	4	1	1	2	1	-	-	-
Female	2	-	1	5	2	8	4	3	2	-	-	-
Black, both sexes	-	-	1	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
South Dakota												
All races, both sexes	-	-	-	1	1	1	4	1	2	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	4	1	2	1	-	-
White, both sexes	-	-	-	1	1	1	4	1	2	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	4	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	2	4	1	12	14	17	7	4	-	-	-
Male	-	1	1	-	2	2	5	2	-	-	-	-
Female	2	1	3	1	10	12	12	5	4	-	-	-
White, both sexes	1	2	3	1	11	13	16	7	4	-	-	-
Male	-	1	1	-	2	2	5	2	-	-	-	-
Female	1	1	2	1	9	11	11	5	4	-	-	-
Black, both sexes	1	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Texas												
All races, both sexes	9	17	14	20	21	30	26	23	7	-	-	-
Male	4	5	4	5	6	9	5	3	-	-	-	-
Female	5	12	10	15	15	21	21	20	7	-	-	-
White, both sexes	8	16	12	18	17	28	26	22	7	-	-	-
Male	4	5	3	5	6	9	5	3	-	-	-	-
Female	4	11	9	13	11	19	21	19	7	-	-	-
Black, both sexes	-	1	2	2	4	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	2	4	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	2	5	6	6	3	3	1	1	-	-
Male	-	-	-	2	4	1	-	1	-	1	-	-
Female	-	-	2	3	2	5	3	2	1	-	-	-
White, both sexes	-	-	2	5	6	6	3	3	1	1	-	-
Male	-	-	-	2	4	1	-	1	-	1	-	-
Female	-	-	2	3	2	5	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Vermont												
All races, both sexes	-	-	-	1	1	4	4	2	1	1	-	-
Male	-	-	-	-	-	4	-	-	-	-	-	-
Female	-	-	-	1	1	-	4	2	1	1	-	-
White, both sexes	-	-	-	1	1	4	4	2	1	1	-	-
Male	-	-	-	-	-	4	-	-	-	-	-	-
Female	-	-	-	1	1	-	4	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	2	2	9	4	10	19	7	4	3	-	-
Male	-	1	1	3	1	3	7	1	-	-	-	-
Female	1	1	1	6	3	7	12	6	4	3	-	-
White, both sexes	1	1	2	8	4	8	18	5	4	2	-	-
Male	-	-	1	3	1	3	6	1	-	-	-	-
Female	1	1	1	5	3	5	12	4	4	2	-	-
Black, both sexes	-	1	-	1	-	1	1	2	-	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Washington												
All races, both sexes	2	-	-	2	7	8	8	10	5	1	-	-
Male	1	-	-	1	3	1	3	3	2	1	-	-
Female	1	-	-	1	4	7	5	7	3	-	-	-
White, both sexes	2	-	-	2	6	7	8	10	4	1	-	-
Male	1	-	-	1	3	1	3	3	2	1	-	-
Female	1	-	-	1	3	6	5	7	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	-	2	8	13	4	4	1	-	-	-
Male	-	-	-	1	2	5	1	-	-	-	-	-
Female	-	1	-	1	6	8	3	4	1	-	-	-
White, both sexes	-	1	-	2	8	13	4	4	-	-	-	-
Male	-	-	-	1	2	5	1	-	-	-	-	-
Female	-	1	-	1	6	8	3	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Wisconsin												
All races, both sexes	2	1	3	5	3	8	16	9	2	2	-	-
Male	2	1	1	-	2	2	2	1	-	-	-	-
Female	-	-	2	5	1	6	14	8	2	2	-	-
White, both sexes	2	1	3	4	3	8	16	9	2	2	-	-
Male	2	1	1	-	2	2	2	1	-	-	-	-
Female	-	-	2	4	1	6	14	8	2	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	2	-	5	2	-	2	-	-	-
Male	-	-	-	2	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	4	2	-	1	-	-	-
White, both sexes	-	-	-	2	-	5	2	-	2	-	-	-
Male	-	-	-	2	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	4	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
United States												
All races, both sexes	1,392	-	1	2	9	26	67	63	71	81	126	131
Male	186	-	-	-	1	7	14	7	6	7	13	18
Female	1,206	-	1	2	8	19	53	56	65	74	113	113
White, both sexes	854	-	1	1	4	8	26	27	27	36	60	57
Male	122	-	-	-	-	1	7	3	3	2	9	8
Female	732	-	1	1	4	7	19	24	24	34	51	49
Black, both sexes	472	-	-	-	4	16	35	33	36	41	60	66
Male	59	-	-	-	1	5	6	4	3	5	4	9
Female	413	-	-	-	3	11	29	29	33	36	56	57
American Indian 1/, both sexes	14	-	-	-	-	1	1	1	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	1	1	1	-	-	1	2
Asian or Pacific Islander 2/, both sexes	52	-	-	1	1	1	5	2	8	4	5	6
Male	4	-	-	-	-	1	1	-	-	-	-	1
Female	48	-	-	1	1	-	4	2	8	4	5	5
Alabama												
All races, both sexes	20	-	-	-	-	-	1	2	1	-	5	1
Male	4	-	-	-	-	-	-	1	-	-	1	-
Female	16	-	-	-	-	-	1	1	1	-	4	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	1	2	1	-	4	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	1	1	1	-	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	33	-	-	-	-	-	1	-	1	2	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	1	-	1	2	-	2
White, both sexes	30	-	-	-	-	-	-	-	1	2	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	1	2	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Arkansas												
All races, both sexes	21	-	-	-	-	-	2	-	-	1	3	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	2	-	-	-	3	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	8	-	-	-	-	-	2	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	182	-	-	-	1	7	10	10	8	10	17	20
Male	19	-	-	-	-	1	2	1	-	-	3	2
Female	163	-	-	-	1	6	8	9	8	10	14	18
White, both sexes	114	-	-	-	-	4	6	4	3	6	5	12
Male	11	-	-	-	-	-	1	-	-	-	1	2
Female	103	-	-	-	-	4	5	4	3	6	4	10
Black, both sexes	47	-	-	-	1	2	2	5	2	1	10	7
Male	5	-	-	-	-	-	-	1	-	-	2	-
Female	42	-	-	-	1	2	2	4	2	1	8	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	1	2	1	3	3	2	1
Male	3	-	-	-	-	1	1	-	-	-	-	-
Female	17	-	-	-	-	-	1	1	3	3	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Colorado												
All races, both sexes	11	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	1	-	-	-	2	-	-	-	-	-
White, both sexes	9	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	9	-	-	-	-	2	-	1	-	-	-	-
Male	3	-	-	-	-	1	-	1	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	1	1	-	1	3
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	1	-	1	3
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	1	1	-	1	3
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	1	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Florida												
All races, both sexes	106	-	-	-	1	2	7	7	8	7	14	7
Male	15	-	-	-	-	-	-	1	1	2	2	-
Female	91	-	-	-	1	2	7	6	7	5	12	7
White, both sexes	73	-	-	-	-	2	3	5	5	4	7	3
Male	12	-	-	-	-	-	-	1	-	1	2	-
Female	61	-	-	-	-	2	3	4	5	3	5	3
Black, both sexes	32	-	-	-	1	-	3	2	3	3	7	4
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	29	-	-	-	1	-	3	2	2	2	7	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	50	-	-	-	-	2	8	2	3	2	7	4
Male	13	-	-	-	-	1	3	1	-	1	1	2
Female	37	-	-	-	-	1	5	1	3	1	6	2
White, both sexes	17	-	-	-	-	-	-	1	-	-	2	2
Male	6	-	-	-	-	-	-	1	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	31	-	-	-	-	2	7	1	3	2	5	2
Male	7	-	-	-	-	1	3	-	-	1	1	1
Female	24	-	-	-	-	1	4	1	3	1	4	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Hawaii												
All races, both sexes	8	-	-	-	1	-	-	-	1	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	1	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	1	-	-	-	1	-	-	-
Idaho												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Illinois												
All races, both sexes	55	-	-	-	1	1	1	4	4	2	1	11
Male	4	-	-	-	-	1	-	-	-	-	-	1
Female	51	-	-	-	1	-	1	4	4	2	1	10
White, both sexes	30	-	-	-	1	-	1	1	2	-	1	6
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	1	-	1	1	2	-	1	6
Black, both sexes	22	-	-	-	-	1	-	3	1	2	-	4
Male	3	-	-	-	-	1	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	3	1	2	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
Indiana												
All races, both sexes	27	-	-	-	-	-	4	-	2	2	3	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	4	-	2	2	3	2
White, both sexes	19	-	-	-	-	-	2	-	-	1	2	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	2	-	-	1	2	2
Black, both sexes	8	-	-	-	-	-	2	-	2	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	2	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Iowa												
All races, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	17	-	-	1	-	-	-	1	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	1	-	-	-	1	-	1	1	-
White, both sexes	13	-	-	1	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	1	-	-	-	1	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Kentucky												
All races, both sexes	14	-	-	-	-	-	-	-	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	30	-	-	-	-	1	2	2	1	2	1	4
Male	7	-	-	-	-	-	1	-	-	-	-	1
Female	23	-	-	-	-	1	1	2	1	2	1	3
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	1	2	2	1	2	1	3
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	20	-	-	-	-	1	1	2	1	2	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	26	-	-	-	1	-	-	-	-	1	3	4
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	1	-	-	-	-	1	3	3
White, both sexes	9	-	-	-	1	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	1	3	3
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Massachusetts												
All races, both sexes	14	-	-	-	-	-	-	-	2	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	12	-	-	-	-	-	-	-	2	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	2	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	38	-	-	-	-	3	1	-	3	3	3	4
Male	4	-	-	-	-	1	-	-	-	-	-	1
Female	34	-	-	-	-	2	1	-	3	3	3	3
White, both sexes	18	-	-	-	-	-	-	-	1	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	1	1	1	-
Black, both sexes	19	-	-	-	-	3	1	-	1	2	2	4
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	17	-	-	-	-	2	1	-	1	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Minnesota												
All races, both sexes	12	-	-	-	-	1	-	-	1	-	2	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	1	-	-	1	-	1	1
White, both sexes	7	-	-	-	-	1	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	2	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	1	-	2	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	1	-	2	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Missouri												
All races, both sexes	26	-	-	-	-	-	1	-	1	3	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	1	-	1	3	1	1
White, both sexes	21	-	-	-	-	-	-	-	-	3	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	3	1	-
Black, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	1	1	-	1	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	1	1	-	1	1	-	-
White, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	50	-	-	-	-	-	2	2	3	6	3	6
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	45	-	-	-	-	-	2	2	3	6	3	5
White, both sexes	24	-	-	-	-	-	1	-	2	1	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	1	-	2	1	1	2
Black, both sexes	25	-	-	-	-	-	1	2	1	4	2	4
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	1	2	1	4	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
New Mexico												
All races, both sexes	20	-	-	-	-	-	3	1	-	1	3	1
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	18	-	-	-	-	-	2	1	-	1	3	1
White, both sexes	18	-	-	-	-	-	3	1	-	1	2	1
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	16	-	-	-	-	-	2	1	-	1	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	81	-	-	1	-	-	4	4	5	4	8	10
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	75	-	-	1	-	-	3	4	5	4	8	10
White, both sexes	52	-	-	-	-	-	2	3	2	2	7	4
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	49	-	-	-	-	-	1	3	2	2	7	4
Black, both sexes	26	-	-	-	-	-	2	1	3	2	1	4
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	2	1	3	2	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	1	-	-	-	-	-	-	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
North Carolina												
All races, both sexes	52	-	-	-	-	1	-	2	2	3	8	3
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	1	-	2	2	3	8	3
White, both sexes	22	-	-	-	-	-	-	1	-	-	2	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	29	-	-	-	-	1	-	1	2	3	5	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	1	-	1	2	3	5	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Ohio												
All races, both sexes	51	-	-	-	1	-	1	5	2	7	3	3
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	46	-	-	-	1	-	1	5	2	6	3	3
White, both sexes	37	-	-	-	-	-	-	3	2	2	2	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	3	2	2	2	2
Black, both sexes	14	-	-	-	1	-	1	2	-	5	1	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	1	-	1	2	-	4	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	31	-	-	-	-	1	2	-	-	1	-	4
Male	5	-	-	-	-	1	1	-	-	-	-	-
Female	26	-	-	-	-	-	1	-	-	1	-	4
White, both sexes	20	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	7	-	-	-	-	1	2	-	-	-	-	1
Male	3	-	-	-	-	1	1	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Oregon												
All races, both sexes	9	-	-	-	-	-	1	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	1	-	-	-	1
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	59	-	-	-	2	-	1	3	1	-	4	6
Male	8	-	-	-	1	-	-	-	-	-	-	2
Female	51	-	-	-	1	-	1	3	1	-	4	4
White, both sexes	37	-	-	-	1	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	1	-	-	-	-	-	2	1
Black, both sexes	22	-	-	-	1	-	1	3	1	-	2	5
Male	3	-	-	-	1	-	-	-	-	-	-	2
Female	19	-	-	-	-	-	1	3	1	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	27	-	-	-	-	1	-	-	3	4	2	1
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	1	-	-	2	4	2	1
White, both sexes	13	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	14	-	-	-	-	1	-	-	3	3	1	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	1	-	-	2	3	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	35	-	-	-	-	-	1	1	2	1	3	8
Male	7	-	-	-	-	-	-	-	1	-	-	3
Female	28	-	-	-	-	-	1	1	1	1	3	5
White, both sexes	22	-	-	-	-	-	-	-	-	1	2	3
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	13	-	-	-	-	-	1	1	2	-	1	5
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	1	1	1	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Texas												
All races, both sexes	125	-	-	-	1	2	5	6	8	12	12	9
Male	18	-	-	-	-	-	3	1	1	1	1	-
Female	107	-	-	-	1	2	2	5	7	11	11	9
White, both sexes	85	-	-	-	1	-	3	5	4	8	8	3
Male	16	-	-	-	-	-	2	1	1	1	1	-
Female	69	-	-	-	1	-	1	4	3	7	7	3
Black, both sexes	37	-	-	-	-	2	2	1	3	4	3	6
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	35	-	-	-	-	2	1	1	3	4	3	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
Utah												
All races, both sexes	13	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	1	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	1	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	32	-	-	-	-	-	2	3	3	2	4	3
Male	4	-	-	-	-	-	-	1	1	-	-	-
Female	28	-	-	-	-	-	2	2	2	2	4	3
White, both sexes	17	-	-	-	-	-	-	-	1	-	1	2
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	14	-	-	-	-	-	2	3	2	2	3	1
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	2	2	2	2	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Washington												
All races, both sexes	23	-	-	-	-	1	2	2	1	-	1	-
Male	6	-	-	-	-	1	1	-	1	-	-	-
Female	17	-	-	-	-	-	1	2	-	-	1	-
White, both sexes	16	-	-	-	-	-	1	1	1	-	-	-
Male	5	-	-	-	-	-	1	-	1	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	3	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	2	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Wisconsin												
All races, both sexes	13	-	-	-	-	-	-	2	1	1	2	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	2	1	1	2	1
White, both sexes	9	-	-	-	-	-	-	1	-	1	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	4	-	-	-	-	-	-	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
United States												
All races, both sexes	143	142	111	93	90	91	79	38	22	4	2	-
Male	21	20	16	13	13	15	11	3	1	-	-	-
Female	122	122	95	80	77	76	68	35	21	4	2	-
White, both sexes	90	92	80	72	72	71	73	34	18	4	1	-
Male	17	16	13	11	11	8	10	2	1	-	-	-
Female	73	76	67	61	61	63	63	32	17	4	1	-
Black, both sexes	45	44	29	18	17	17	6	1	4	-	-	-
Male	4	4	2	1	2	7	1	1	-	-	-	-
Female	41	40	27	17	15	10	5	-	4	-	-	-
American Indian 1/, both sexes	4	3	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	4	3	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	1	3	1	3	-	3	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	4	3	1	2	1	3	-	3	-	-	1	-
Alabama												
All races, both sexes	-	-	3	1	2	2	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	3	1	1	2	-	1	-	-	-	-
White, both sexes	-	-	2	1	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	1	1	2	-	1	-	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6	5	5	4	2	1	2	1	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	5	4	5	3	2	1	2	1	-	-	-	-
White, both sexes	5	4	5	4	2	1	2	1	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	4	3	5	3	2	1	2	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Arkansas												
All races, both sexes	4	3	2	1	1	2	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	4	3	2	1	-	2	-	1	-	-	-	-
White, both sexes	1	3	2	1	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	3	2	1	-	2	-	1	-	-	-	-
Black, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23	18	13	11	10	8	9	4	2	-	1	-
Male	3	2	-	2	1	1	1	-	-	-	-	-
Female	20	16	13	9	9	7	8	4	2	-	1	-
White, both sexes	17	11	9	8	9	8	7	3	2	-	-	-
Male	3	1	-	1	1	1	-	-	-	-	-	-
Female	14	10	9	7	8	7	7	3	2	-	-	-
Black, both sexes	5	4	3	2	1	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	5	3	3	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	-	-	-	1	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	2	1	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Colorado												
All races, both sexes	2	2	1	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	1	-	2	-	-	1	-	-	-	-
White, both sexes	2	2	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	1	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	1	1	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	1	1	-	-	-
White, both sexes	-	1	-	1	1	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Delaware												
All races, both sexes	2	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Florida												
All races, both sexes	7	9	7	1	8	8	8	4	1	-	-	-
Male	3	1	1	-	3	-	1	-	-	-	-	-
Female	4	8	6	1	5	8	7	4	1	-	-	-
White, both sexes	5	6	5	1	7	7	8	4	1	-	-	-
Male	2	1	1	-	3	-	1	-	-	-	-	-
Female	3	5	4	1	4	7	7	4	1	-	-	-
Black, both sexes	2	3	2	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	3	2	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	6	5	-	4	4	-	1	-	2	-	-	-
Male	-	2	-	2	-	-	-	-	-	-	-	-
Female	6	3	-	2	4	-	1	-	2	-	-	-
White, both sexes	3	3	-	4	2	-	-	-	-	-	-	-
Male	-	2	-	2	-	-	-	-	-	-	-	-
Female	3	1	-	2	2	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	2	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	2	-	1	-	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Hawaii												
All races, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Idaho												
All races, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Illinois												
All races, both sexes	3	3	5	5	5	4	3	1	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	3	2	5	5	5	3	3	1	-	1	-	-
White, both sexes	1	3	2	4	2	1	3	1	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	2	2	4	2	1	3	1	-	1	-	-
Black, both sexes	2	-	3	1	2	3	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	3	1	2	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	3	2	-	2	3	1	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	2	2	-	2	3	1	1	-	-	-
White, both sexes	1	-	2	2	-	2	3	1	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	2	3	1	1	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Iowa												
All races, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	2	1	2	2	1	2	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	2	2	1	2	1	1	2	-	1	-	-	-
White, both sexes	1	2	1	1	2	1	2	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	2	1	1	1	1	2	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Kentucky												
All races, both sexes	-	3	2	2	1	1	1	1	-	-	-	-
Male	-	1	1	-	-	-	1	1	-	-	-	-
Female	-	2	1	2	1	1	-	-	-	-	-	-
White, both sexes	-	3	2	1	1	1	1	1	-	-	-	-
Male	-	1	1	-	-	-	1	1	-	-	-	-
Female	-	2	1	1	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	5	3	1	1	2	1	-	-	-	-	-
Male	2	2	-	-	-	1	-	-	-	-	-	-
Female	2	3	3	1	1	1	1	-	-	-	-	-
White, both sexes	1	2	1	1	-	1	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
Black, both sexes	3	3	2	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	2	3	2	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Maine												
All races, both sexes	-	-	1	-	-	1	-	1	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Maryland												
All races, both sexes	3	3	3	2	1	3	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	3	3	3	2	1	3	1	-	-	-	-	-
White, both sexes	-	2	-	2	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	2	-	2	1	-	-	-	-	-
Black, both sexes	3	1	3	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	3	1	3	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Massachusetts												
All races, both sexes	1	1	4	-	-	1	1	-	2	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	1	3	-	-	-	1	-	2	-	-	-
White, both sexes	1	1	3	-	-	1	1	-	2	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	1	2	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	8	2	1	4	-	-	2	1	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	2	7	1	1	4	-	-	2	1	1	-	-
White, both sexes	1	5	2	-	3	-	-	2	1	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	1	4	1	-	3	-	-	2	1	1	-	-
Black, both sexes	1	3	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	3	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Minnesota												
All races, both sexes	-	2	1	3	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	3	-	-	-	1	-	-	-	-
White, both sexes	-	1	1	2	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	2	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	2	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	2	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Missouri												
All races, both sexes	2	3	1	2	5	1	5	-	-	-	-	-
Male	1	1	1	1	2	1	-	-	-	-	-	-
Female	1	2	-	1	3	-	5	-	-	-	-	-
White, both sexes	2	3	1	-	5	1	5	-	-	-	-	-
Male	1	1	1	-	2	1	-	-	-	-	-	-
Female	1	2	-	-	3	-	5	-	-	-	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Nebraska												
All races, both sexes	2	-	1	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	2	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	2	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
New Hampshire												
All races, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	9	6	3	3	-	2	-	-	-	-	-
Male	1	3	-	-	-	-	-	-	-	-	-	-
Female	4	6	6	3	3	-	2	-	-	-	-	-
White, both sexes	3	3	5	2	3	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	5	2	3	-	1	-	-	-	-	-
Black, both sexes	2	6	1	1	-	-	1	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	2	4	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
New Mexico												
All races, both sexes	3	-	1	3	2	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	1	2	2	1	-	-	1	-	-	-
White, both sexes	2	-	1	3	2	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	1	2	2	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	9	4	5	4	6	3	1	1	1	-	-
Male	2	-	-	-	-	2	1	-	-	-	-	-
Female	9	9	4	5	4	4	2	1	1	1	-	-
White, both sexes	6	9	2	2	4	4	3	1	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	5	9	2	2	4	4	2	1	-	1	-	-
Black, both sexes	5	-	2	3	-	2	-	-	1	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	4	-	2	3	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
North Carolina												
All races, both sexes	3	9	3	4	4	5	2	1	2	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	9	3	4	4	5	2	1	2	-	-	-
White, both sexes	-	3	-	4	3	4	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	3	-	4	3	4	1	1	2	-	-	-
Black, both sexes	3	6	3	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	6	3	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Ohio												
All races, both sexes	5	1	2	4	3	7	4	2	1	-	-	-
Male	1	-	-	-	-	2	-	-	1	-	-	-
Female	4	1	2	4	3	5	4	2	-	-	-	-
White, both sexes	4	-	2	4	3	6	4	2	1	-	-	-
Male	1	-	-	-	-	2	-	-	1	-	-	-
Female	3	-	2	4	3	4	4	2	-	-	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	2	7	1	3	2	4	2	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	2	2	5	1	3	2	3	2	-	-	-	-
White, both sexes	1	2	3	1	3	2	4	2	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	2	3	1	3	2	3	2	-	-	-	-
Black, both sexes	-	-	3	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Oregon												
All races, both sexes	1	1	1	-	-	1	-	2	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	2	-	-	-	-
White, both sexes	1	-	1	-	-	1	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	11	7	6	4	2	3	5	1	2	1	-	-
Male	2	1	-	1	1	-	-	-	-	-	-	-
Female	9	6	6	3	1	3	5	1	2	1	-	-
White, both sexes	9	3	5	3	2	2	5	1	2	1	-	-
Male	2	1	-	1	1	-	-	-	-	-	-	-
Female	7	2	5	2	1	2	5	1	2	1	-	-
Black, both sexes	2	4	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	4	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	4	2	3	2	2	2	-	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	1	3	2	2	2	1	2	-	-	-	-	-
White, both sexes	1	3	2	1	2	-	2	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	2	2	-	2	-	2	-	-	-	-	-
Black, both sexes	-	1	-	2	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
South Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	-	1	1	4	5	2	1	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	5	-	-	1	3	4	2	1	-	-	-	-
White, both sexes	3	-	1	1	4	4	2	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	3	-	-	1	3	4	2	1	-	-	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Texas												
All races, both sexes	12	9	11	11	6	6	11	2	2	-	-	-
Male	1	2	2	1	1	2	2	-	-	-	-	-
Female	11	7	9	10	5	4	9	2	2	-	-	-
White, both sexes	7	5	9	8	4	5	11	2	2	-	-	-
Male	1	2	1	1	1	2	2	-	-	-	-	-
Female	6	3	8	7	3	3	9	2	2	-	-	-
Black, both sexes	5	4	2	2	2	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	5	4	1	2	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	1	1	-	1	3	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	1	3	-	2	-	-	-	-
White, both sexes	2	1	1	-	1	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	1	-	1	2	-	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Vermont												
All races, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	2	2	2	3	2	-	1	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	2	1	-	2	2	3	2	-	1	-	-	-
White, both sexes	2	-	2	1	2	3	2	-	1	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	2	3	2	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Washington												
All races, both sexes	4	3	1	2	3	2	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	3	3	-	2	3	2	-	-	-	-	-	-
White, both sexes	2	3	1	2	2	2	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	1	3	-	2	2	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Wisconsin												
All races, both sexes	2	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	2	-	1	1	-	-	-	-	-
White, both sexes	1	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	2	-	1	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
United States												
All races, both sexes	4,789	2	6	2	8	10	25	36	65	93	146	202
Male	1,458	2	3	1	1	6	9	8	22	28	57	70
Female	3,331	-	3	1	7	4	16	28	43	65	89	132
White, both sexes	4,026	2	4	2	5	5	15	18	31	48	86	126
Male	1,225	2	2	1	1	3	7	3	12	14	32	48
Female	2,801	-	2	1	4	2	8	15	19	34	54	78
Black, both sexes	646	-	2	-	3	5	9	18	30	43	54	67
Male	193	-	1	-	-	3	2	5	10	13	24	20
Female	453	-	1	-	3	2	7	13	20	30	30	47
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	-	-	3	2
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	2	2
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	1	-	4	2	3	7
Male	33	-	-	-	-	-	-	-	-	1	-	2
Female	52	-	-	-	-	-	1	-	4	1	3	5
Alabama												
All races, both sexes	60	-	-	-	-	-	-	1	-	-	4	2
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	-	1	-	-	3	1
White, both sexes	45	-	-	-	-	-	-	1	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	3	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Alaska												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	82	-	-	-	-	1	1	1	1	1	1	3
Male	26	-	-	-	-	-	1	1	-	-	-	3
Female	56	-	-	-	-	1	-	-	1	1	1	-
White, both sexes	79	-	-	-	-	1	1	1	1	1	1	3
Male	26	-	-	-	-	-	1	1	-	-	-	3
Female	53	-	-	-	-	1	-	-	1	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Arkansas												
All races, both sexes	37	-	-	-	-	-	-	-	1	1	2	-
Male	17	-	-	-	-	-	-	-	-	1	2	-
Female	20	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	1	1	2	-
Male	17	-	-	-	-	-	-	-	-	1	2	-
Female	16	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	472	1	1	1	1	1	1	2	6	10	15	21
Male	146	1	1	-	-	-	-	-	1	2	6	7
Female	326	-	-	1	1	1	1	2	5	8	9	14
White, both sexes	394	1	1	1	1	1	1	2	4	7	7	16
Male	117	1	1	-	-	-	-	-	1	2	1	6
Female	277	-	-	1	1	1	1	2	3	5	6	10
Black, both sexes	45	-	-	-	-	-	-	-	-	3	5	4
Male	19	-	-	-	-	-	-	-	-	-	4	1
Female	26	-	-	-	-	-	-	-	-	3	1	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	2	-	2	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	2	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Colorado												
All races, both sexes	69	-	-	-	-	-	1	1	-	1	5	3
Male	16	-	-	-	-	-	1	-	-	-	-	1
Female	53	-	-	-	-	-	-	1	-	1	5	2
White, both sexes	64	-	-	-	-	-	1	-	-	1	5	-
Male	14	-	-	-	-	-	1	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	1	5	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	6	4
Male	22	-	-	-	-	-	-	-	-	-	2	1
Female	42	-	-	-	-	-	-	-	-	1	4	3
White, both sexes	56	-	-	-	-	-	-	-	-	-	3	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	3	3
Male	4	-	-	-	-	-	-	-	-	-	2	1
Female	4	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Delaware												
All races, both sexes	18	-	1	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	2	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	1	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Florida												
All races, both sexes	297	-	1	-	1	1	5	1	3	6	6	16
Male	79	-	1	-	-	1	1	1	1	3	3	4
Female	218	-	-	-	1	-	4	-	2	3	3	12
White, both sexes	266	-	1	-	-	-	4	-	1	5	5	12
Male	69	-	1	-	-	-	1	-	1	2	2	3
Female	197	-	-	-	-	-	3	-	-	3	3	9
Black, both sexes	28	-	-	-	1	1	1	1	2	1	1	3
Male	9	-	-	-	-	1	-	1	-	1	1	1
Female	19	-	-	-	1	-	1	-	2	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	131	-	1	-	-	-	1	4	4	6	4	9
Male	52	-	-	-	-	-	1	1	2	4	3	3
Female	79	-	1	-	-	-	-	3	2	2	1	6
White, both sexes	85	-	1	-	-	-	-	-	4	-	2	3
Male	34	-	-	-	-	-	-	-	2	-	2	-
Female	51	-	1	-	-	-	-	-	2	-	-	3
Black, both sexes	44	-	-	-	-	-	1	4	-	6	2	6
Male	17	-	-	-	-	-	1	1	-	4	1	3
Female	27	-	-	-	-	-	-	3	-	2	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Hawaii												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Illinois												
All races, both sexes	228	1	-	-	-	-	2	2	5	6	8	6
Male	72	1	-	-	-	-	1	-	1	2	2	3
Female	156	-	-	-	-	-	1	2	4	4	6	3
White, both sexes	184	1	-	-	-	-	1	1	2	2	4	3
Male	59	1	-	-	-	-	1	-	1	-	-	2
Female	125	-	-	-	-	-	-	1	1	2	4	1
Black, both sexes	39	-	-	-	-	-	1	1	3	4	4	2
Male	11	-	-	-	-	-	-	-	-	2	2	1
Female	28	-	-	-	-	-	1	1	3	2	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	94	-	-	-	-	-	-	-	1	-	3	4
Male	36	-	-	-	-	-	-	-	-	-	3	2
Female	58	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	83	-	-	-	-	-	-	-	-	-	2	1
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	1	-	1	3
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Iowa												
All races, both sexes	75	-	-	-	-	-	-	1	-	-	1	2
Male	29	-	-	-	-	-	-	1	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	74	-	-	-	-	-	-	1	-	-	1	1
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	63	-	-	-	1	-	1	-	-	-	-	4
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	1	-	1	-	-	-	-	3
White, both sexes	56	-	-	-	-	-	1	-	-	-	-	4
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	1	-	-	-	-	3
Black, both sexes	5	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Kentucky												
All races, both sexes	77	-	-	-	-	-	2	-	2	1	1	2
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	56	-	-	-	-	-	2	-	2	1	-	2
White, both sexes	71	-	-	-	-	-	2	-	1	1	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	-	-	2	-	1	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	75	-	-	-	-	-	1	1	2	3	4	5
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	54	-	-	-	-	-	1	1	2	3	3	4
White, both sexes	48	-	-	-	-	-	-	-	1	-	2	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	27	-	-	-	-	-	1	1	1	3	2	4
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	1	1	1	3	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Maine												
All races, both sexes	25	-	-	-	-	-	-	1	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	25	-	-	-	-	-	-	1	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	93	-	-	-	1	-	-	1	-	1	3	4
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	72	-	-	-	1	-	-	1	-	1	2	4
White, both sexes	66	-	-	-	1	-	-	1	-	-	-	3
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	1	-	-	1	-	-	-	3
Black, both sexes	26	-	-	-	-	-	-	-	-	1	3	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Massachusetts												
All races, both sexes	81	-	-	-	-	-	-	-	-	1	2	2
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	74	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	189	-	-	-	1	-	-	-	3	2	3	5
Male	66	-	-	-	-	-	-	-	2	-	1	1
Female	123	-	-	-	1	-	-	-	1	2	2	4
White, both sexes	157	-	-	-	-	-	-	-	-	-	1	2
Male	54	-	-	-	-	-	-	-	-	-	1	-
Female	103	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	29	-	-	-	1	-	-	-	3	2	2	3
Male	11	-	-	-	-	-	-	-	2	-	-	1
Female	18	-	-	-	1	-	-	-	1	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Minnesota												
All races, both sexes	108	-	-	-	-	-	-	-	-	1	2	3
Male	33	-	-	-	-	-	-	-	-	1	1	2
Female	75	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	104	-	-	-	-	-	-	-	-	1	1	2
Male	31	-	-	-	-	-	-	-	-	1	1	1
Female	73	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	40	-	1	-	-	-	1	2	1	1	2	4
Male	18	-	1	-	-	-	-	1	1	1	2	2
Female	22	-	-	-	-	-	1	1	-	-	-	2
White, both sexes	22	-	-	-	-	-	-	2	-	-	1	1
Male	10	-	-	-	-	-	-	1	-	-	1	1
Female	12	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	18	-	1	-	-	-	1	-	1	1	1	3
Male	8	-	1	-	-	-	-	-	1	1	1	1
Female	10	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Missouri												
All races, both sexes	98	-	-	-	-	-	1	2	-	3	5	3
Male	27	-	-	-	-	-	-	-	-	1	2	-
Female	71	-	-	-	-	-	1	2	-	2	3	3
White, both sexes	83	-	-	-	-	-	-	2	-	3	2	3
Male	25	-	-	-	-	-	-	-	-	1	2	-
Female	58	-	-	-	-	-	-	2	-	2	-	3
Black, both sexes	13	-	-	-	-	-	1	-	-	-	3	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	1	-	-	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Nebraska												
All races, both sexes	39	-	-	-	1	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	1	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	25	-	-	-	-	-	1	-	1	-	2	3
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	1	-	1	-	1	3
White, both sexes	22	-	-	-	-	-	-	-	1	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	158	-	-	-	-	2	-	1	2	5	7	11
Male	43	-	-	-	-	2	-	1	-	2	3	2
Female	115	-	-	-	-	-	-	-	2	3	4	9
White, both sexes	126	-	-	-	-	1	-	-	-	4	4	4
Male	37	-	-	-	-	1	-	-	-	1	2	1
Female	89	-	-	-	-	-	-	-	-	3	2	3
Black, both sexes	27	-	-	-	-	1	-	1	2	1	3	7
Male	5	-	-	-	-	1	-	1	-	1	1	1
Female	22	-	-	-	-	-	-	-	2	-	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
New Mexico												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	2	2
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	37	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	212	-	-	-	1	-	1	1	4	6	8	15
Male	77	-	-	-	-	-	1	-	3	1	4	9
Female	135	-	-	-	1	-	-	1	1	5	4	6
White, both sexes	169	-	-	-	1	-	1	-	2	4	5	9
Male	63	-	-	-	-	-	1	-	1	1	3	7
Female	106	-	-	-	1	-	-	-	1	3	2	2
Black, both sexes	38	-	-	-	-	-	-	1	2	2	3	4
Male	11	-	-	-	-	-	-	-	2	-	1	1
Female	27	-	-	-	-	-	-	1	-	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
North Carolina												
All races, both sexes	160	-	1	-	-	-	-	1	3	6	4	3
Male	43	-	-	-	-	-	-	-	1	1	2	-
Female	117	-	1	-	-	-	-	1	2	5	2	3
White, both sexes	118	-	1	-	-	-	-	-	1	1	1	1
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	87	-	1	-	-	-	-	-	1	1	1	1
Black, both sexes	40	-	-	-	-	-	-	1	2	5	3	2
Male	11	-	-	-	-	-	-	-	1	1	2	-
Female	29	-	-	-	-	-	-	1	1	4	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Ohio												
All races, both sexes	226	-	-	-	-	1	3	2	3	7	7	10
Male	62	-	-	-	-	1	-	1	1	-	-	4
Female	164	-	-	-	-	-	3	1	2	7	7	6
White, both sexes	191	-	-	-	-	1	1	-	2	3	5	8
Male	57	-	-	-	-	1	-	-	1	-	-	4
Female	134	-	-	-	-	-	1	-	1	3	5	4
Black, both sexes	35	-	-	-	-	-	2	2	1	4	2	2
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	30	-	-	-	-	-	2	1	1	4	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	53	-	-	-	-	-	-	1	-	1	-	4
Male	22	-	-	-	-	-	-	-	-	-	-	3
Female	31	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	45	-	-	-	-	-	-	1	-	-	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	3
Female	25	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Oregon												
All races, both sexes	92	-	-	-	-	-	-	-	-	1	1	1
Male	29	-	-	-	-	-	-	-	-	1	1	-
Female	63	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	87	-	-	-	-	-	-	-	-	1	1	1
Male	27	-	-	-	-	-	-	-	-	1	1	-
Female	60	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	217	-	-	-	-	1	-	2	3	-	2	9
Male	68	-	-	-	-	-	-	-	1	-	-	6
Female	149	-	-	-	-	1	-	2	2	-	2	3
White, both sexes	185	-	-	-	-	-	-	1	-	-	1	5
Male	58	-	-	-	-	-	-	-	-	-	-	4
Female	127	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	31	-	-	-	-	1	-	1	2	-	1	4
Male	10	-	-	-	-	-	-	-	1	-	-	2
Female	21	-	-	-	-	1	-	1	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	52	-	-	-	-	2	-	-	3	2	1	2
Male	14	-	-	-	-	1	-	-	2	1	-	1
Female	38	-	-	-	-	1	-	-	1	1	1	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	17	-	-	-	-	2	-	-	3	2	1	-
Male	5	-	-	-	-	1	-	-	2	1	-	-
Female	12	-	-	-	-	1	-	-	1	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
South Dakota												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	116	-	-	-	-	-	-	2	1	3	4	3
Male	40	-	-	-	-	-	-	1	-	2	-	-
Female	76	-	-	-	-	-	-	1	1	1	4	3
White, both sexes	95	-	-	-	-	-	-	-	-	2	3	3
Male	31	-	-	-	-	-	-	-	-	1	-	-
Female	64	-	-	-	-	-	-	-	-	1	3	3
Black, both sexes	20	-	-	-	-	-	-	2	1	1	1	-
Male	8	-	-	-	-	-	-	1	-	1	-	-
Female	12	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Texas												
All races, both sexes	338	-	-	-	1	1	2	5	9	10	17	21
Male	90	-	-	-	1	1	2	-	3	3	7	7
Female	248	-	-	-	-	-	-	5	6	7	10	14
White, both sexes	285	-	-	-	1	1	2	3	4	6	14	14
Male	78	-	-	-	1	1	2	-	2	2	6	4
Female	207	-	-	-	-	-	-	3	2	4	8	10
Black, both sexes	47	-	-	-	-	-	-	2	4	4	2	7
Male	11	-	-	-	-	-	-	-	1	1	1	3
Female	36	-	-	-	-	-	-	2	3	3	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
Utah												
All races, both sexes	36	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	36	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	142	-	-	1	-	-	1	-	1	1	4	2
Male	39	-	-	1	-	-	1	-	-	-	2	-
Female	103	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	105	-	-	1	-	-	-	-	1	1	2	1
Male	25	-	-	1	-	-	-	-	-	-	1	-
Female	80	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	35	-	-	-	-	-	1	-	-	-	2	1
Male	13	-	-	-	-	-	1	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Washington												
All races, both sexes	110	-	-	-	-	-	-	-	3	1	1	3
Male	30	-	-	-	-	-	-	-	1	1	1	-
Female	80	-	-	-	-	-	-	-	2	-	-	3
White, both sexes	102	-	-	-	-	-	-	-	2	1	1	3
Male	29	-	-	-	-	-	-	-	1	1	1	-
Female	73	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	29	-	-	-	-	-	-	1	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	28	-	-	-	-	-	-	1	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Wisconsin												
All races, both sexes	104	-	-	-	-	-	-	-	1	-	3	4
Male	37	-	-	-	-	-	-	-	1	-	2	2
Female	67	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	100	-	-	-	-	-	-	-	1	-	2	3
Male	36	-	-	-	-	-	-	-	1	-	1	2
Female	64	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
United States												
All races, both sexes	243	330	326	393	427	549	656	590	400	213	67	-
Male	84	106	126	137	129	173	192	168	94	34	8	-
Female	159	224	200	256	298	376	464	422	306	179	59	-
White, both sexes	172	236	262	329	384	502	606	559	380	194	60	-
Male	64	78	99	120	116	164	176	156	92	29	6	-
Female	108	158	163	209	268	338	430	403	288	165	54	-
Black, both sexes	62	81	50	47	32	39	41	25	16	17	5	-
Male	20	23	21	15	8	4	11	7	2	4	-	-
Female	42	58	29	32	24	35	30	18	14	13	5	-
American Indian 1/, both sexes	4	4	5	6	3	3	1	-	1	-	-	-
Male	-	2	1	1	-	1	1	-	-	-	-	-
Female	4	2	4	5	3	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	9	9	11	8	5	8	6	3	2	2	-
Male	-	3	5	1	5	4	4	5	-	1	2	-
Female	5	6	4	10	3	1	4	1	3	1	-	-
Alabama												
All races, both sexes	3	5	4	4	7	5	13	5	5	2	-	-
Male	-	2	1	-	4	3	1	1	3	1	-	-
Female	3	3	3	4	3	2	12	4	2	1	-	-
White, both sexes	1	3	3	2	6	5	11	5	4	2	-	-
Male	-	1	1	-	4	3	1	1	2	1	-	-
Female	1	2	2	2	2	2	10	4	2	1	-	-
Black, both sexes	2	2	1	2	1	-	2	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	2	1	1	2	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Alaska												
All races, both sexes	2	-	1	1	2	-	-	2	-	-	-	-
Male	1	-	-	-	1	-	-	1	-	-	-	-
Female	1	-	1	1	1	-	-	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	6	5	14	4	12	14	7	4	2	-	-
Male	2	2	2	2	1	5	5	1	1	-	-	-
Female	3	4	3	12	3	7	9	6	3	2	-	-
White, both sexes	4	5	5	14	4	12	13	7	4	2	-	-
Male	2	2	2	2	1	5	5	1	1	-	-	-
Female	2	3	3	12	3	7	8	6	3	2	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Arkansas												
All races, both sexes	4	1	5	2	3	6	6	2	3	1	-	-
Male	2	-	2	1	1	2	4	1	1	-	-	-
Female	2	1	3	1	2	4	2	1	2	1	-	-
White, both sexes	2	1	5	2	3	4	6	2	3	1	-	-
Male	2	-	2	1	1	2	4	1	1	-	-	-
Female	-	1	3	1	2	2	2	1	2	1	-	-
Black, both sexes	2	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	22	40	28	45	42	67	77	48	22	17	4	-
Male	2	11	13	18	10	24	27	16	4	3	-	-
Female	20	29	15	27	32	43	50	32	18	14	4	-
White, both sexes	14	30	22	37	37	62	65	44	21	16	4	-
Male	1	7	11	16	8	22	20	14	4	2	-	-
Female	13	23	11	21	29	40	45	30	17	14	4	-
Black, both sexes	5	7	3	3	2	3	7	2	-	1	-	-
Male	1	3	1	1	2	1	4	-	-	1	-	-
Female	4	4	2	2	-	2	3	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	3	4	2	2	5	2	1	-	-	-
Male	-	1	1	1	-	1	3	2	-	-	-	-
Female	3	1	2	3	2	1	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Colorado												
All races, both sexes	4	7	1	3	6	6	11	12	6	2	-	-
Male	1	4	-	1	-	2	2	2	2	-	-	-
Female	3	3	1	2	6	4	9	10	4	2	-	-
White, both sexes	4	7	1	3	6	5	11	12	6	2	-	-
Male	1	4	-	1	-	1	2	2	2	-	-	-
Female	3	3	1	2	6	4	9	10	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	1	2	3	2	13	7	8	12	2	1	-
Male	1	-	-	1	1	6	2	4	4	-	-	-
Female	1	1	2	2	1	7	5	4	8	2	1	-
White, both sexes	1	1	2	3	2	13	7	8	12	2	1	-
Male	-	-	-	1	1	6	2	4	4	-	-	-
Female	1	1	2	2	1	7	5	4	8	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Delaware												
All races, both sexes	1	2	-	1	2	1	1	2	-	4	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	1	-	-	2	1	1	1	-	4	-	-
White, both sexes	1	1	-	1	2	1	1	2	-	4	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	1	-	-	2	1	1	1	-	4	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Florida												
All races, both sexes	19	28	20	27	26	32	40	27	20	17	1	-
Male	8	8	6	2	6	8	9	7	8	1	1	-
Female	11	20	14	25	20	24	31	20	12	16	-	-
White, both sexes	15	26	18	23	25	30	39	26	19	16	1	-
Male	7	8	6	1	6	7	9	6	7	1	1	-
Female	8	18	12	22	19	23	30	20	12	15	-	-
Black, both sexes	3	2	2	4	1	1	1	1	1	1	-	-
Male	1	-	-	1	-	-	-	1	1	-	-	-
Female	2	2	2	3	1	1	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	11	9	10	15	15	11	14	7	4	1	-
Male	3	5	6	5	5	4	2	6	2	-	-	-
Female	2	6	3	5	10	11	9	8	5	4	1	-
White, both sexes	2	7	7	6	12	12	6	13	5	4	1	-
Male	1	4	5	3	5	4	1	5	2	-	-	-
Female	1	3	2	3	7	8	5	8	3	4	1	-
Black, both sexes	3	4	1	3	3	3	5	1	2	-	-	-
Male	2	1	-	2	-	-	1	1	-	-	-	-
Female	1	3	1	1	3	3	4	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Hawaii												
All races, both sexes	1	2	2	-	1	-	-	1	1	-	1	-
Male	-	2	2	-	1	-	-	1	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	-	1	-	-	1	1	-	1	-
Male	-	2	2	-	1	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Idaho												
All races, both sexes	-	3	2	2	3	2	7	3	2	2	-	-
Male	-	1	-	2	-	1	2	1	1	1	-	-
Female	-	2	2	-	3	1	5	2	1	1	-	-
White, both sexes	-	3	1	2	3	2	7	3	2	2	-	-
Male	-	1	-	2	-	1	2	1	1	1	-	-
Female	-	2	1	-	3	1	5	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Illinois												
All races, both sexes	9	14	20	17	18	25	28	29	19	14	5	-
Male	4	5	11	7	3	9	8	8	5	2	-	-
Female	5	9	9	10	15	16	20	21	14	12	5	-
White, both sexes	7	7	17	14	15	20	25	29	18	14	4	-
Male	4	3	9	6	2	8	7	8	5	2	-	-
Female	3	4	8	8	13	12	18	21	13	12	4	-
Black, both sexes	2	6	3	3	2	5	2	-	-	-	1	-
Male	-	2	2	1	-	1	-	-	-	-	-	-
Female	2	4	1	2	2	4	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	3	6	4	10	7	12	12	11	14	4	3	-
Male	3	3	1	7	4	2	2	3	5	-	1	-
Female	-	3	3	3	3	10	10	8	9	4	2	-
White, both sexes	2	3	4	10	7	11	12	11	14	3	3	-
Male	2	1	1	7	4	2	2	3	5	-	1	-
Female	-	2	3	3	3	9	10	8	9	3	2	-
Black, both sexes	1	2	-	-	-	1	-	-	-	1	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Iowa												
All races, both sexes	1	4	5	10	5	5	11	11	13	4	2	-
Male	-	2	1	3	1	2	7	4	6	1	-	-
Female	1	2	4	7	4	3	4	7	7	3	2	-
White, both sexes	1	4	5	10	5	5	11	11	13	4	2	-
Male	-	2	1	3	1	2	7	4	6	1	-	-
Female	1	2	4	7	4	3	4	7	7	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	5	5	4	5	10	7	10	3	3	1	-
Male	-	1	3	1	2	2	3	3	1	2	-	-
Female	4	4	2	3	3	8	4	7	2	1	1	-
White, both sexes	3	4	4	2	4	10	7	10	3	3	1	-
Male	-	1	3	1	1	2	3	3	1	2	-	-
Female	3	3	1	1	3	8	4	7	2	1	1	-
Black, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Kentucky												
All races, both sexes	2	6	2	5	11	8	13	11	5	4	2	-
Male	1	2	1	3	3	3	2	3	1	1	-	-
Female	1	4	1	2	8	5	11	8	4	3	2	-
White, both sexes	1	6	2	5	11	7	12	11	5	4	1	-
Male	-	2	1	3	3	3	2	3	1	1	-	-
Female	1	4	1	2	8	4	10	8	4	3	1	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	6	3	3	9	8	9	8	7	2	-	-
Male	-	2	2	-	4	4	2	2	3	-	-	-
Female	4	4	1	3	5	4	7	6	4	2	-	-
White, both sexes	1	3	3	2	6	7	8	6	6	2	-	-
Male	-	1	2	-	4	4	2	2	3	-	-	-
Female	1	2	1	2	2	3	6	4	3	2	-	-
Black, both sexes	3	3	-	1	3	1	1	2	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	3	2	-	1	3	1	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Maine												
All races, both sexes	-	1	1	2	1	-	2	8	4	2	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	1	2	1	-	2	8	3	2	1	-
White, both sexes	-	1	1	2	1	-	2	8	4	2	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	1	2	1	-	2	8	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	9	7	4	6	10	6	19	12	4	-	-
Male	3	1	2	2	1	4	2	3	2	-	-	-
Female	3	8	5	2	5	6	4	16	10	4	-	-
White, both sexes	2	3	4	2	4	8	6	16	12	4	-	-
Male	1	-	1	1	-	4	2	2	2	-	-	-
Female	1	3	3	1	4	4	4	14	10	4	-	-
Black, both sexes	4	6	3	2	2	2	-	2	-	-	-	-
Male	2	1	1	1	1	-	-	1	-	-	-	-
Female	2	5	2	1	1	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Massachusetts												
All races, both sexes	3	8	4	5	5	12	13	15	6	3	2	-
Male	-	4	-	-	1	5	3	2	-	1	1	-
Female	3	4	4	5	4	7	10	13	6	2	1	-
White, both sexes	3	7	3	5	5	12	13	15	6	2	1	-
Male	-	4	-	-	1	5	3	2	-	1	-	-
Female	3	3	3	5	4	7	10	13	6	1	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8	13	18	15	19	20	26	35	9	10	2	-
Male	5	4	9	6	8	8	9	10	1	1	1	-
Female	3	9	9	9	11	12	17	25	8	9	1	-
White, both sexes	4	8	16	13	18	19	25	32	8	9	2	-
Male	2	3	8	5	8	8	8	9	1	-	1	-
Female	2	5	8	8	10	11	17	23	7	9	1	-
Black, both sexes	3	4	2	2	1	1	1	2	1	1	-	-
Male	3	1	1	1	-	-	1	-	-	1	-	-
Female	-	3	1	1	1	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Minnesota												
All races, both sexes	2	5	6	7	8	11	17	15	13	15	3	-
Male	-	2	2	4	3	2	8	2	1	4	1	-
Female	2	3	4	3	5	9	9	13	12	11	2	-
White, both sexes	1	5	5	7	8	11	17	15	13	15	3	-
Male	-	2	1	4	3	2	8	2	1	4	1	-
Female	1	3	4	3	5	9	9	13	12	11	2	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	3	2	2	3	5	7	1	2	1	-	-
Male	-	1	1	1	2	1	4	-	-	-	-	-
Female	2	2	1	1	1	4	3	1	2	1	-	-
White, both sexes	-	1	1	2	2	2	7	1	2	-	-	-
Male	-	-	-	1	1	1	4	-	-	-	-	-
Female	-	1	1	1	1	1	3	1	2	-	-	-
Black, both sexes	2	2	1	-	1	3	-	-	-	1	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	2	1	-	-	-	3	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Missouri												
All races, both sexes	1	8	9	8	9	9	18	9	9	3	1	-
Male	1	-	2	4	3	3	4	3	4	-	-	-
Female	-	8	7	4	6	6	14	6	5	3	1	-
White, both sexes	1	5	7	6	9	7	18	8	9	2	1	-
Male	1	-	1	3	3	3	4	3	4	-	-	-
Female	-	5	6	3	6	4	14	5	5	2	1	-
Black, both sexes	-	2	2	1	-	2	-	1	-	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	2	1	-	-	2	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	1	3	2	2	1	4	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	3	2	1	1	4	-	1	-
White, both sexes	1	-	-	1	3	2	2	1	4	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	3	2	1	1	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Nebraska												
All races, both sexes	1	2	1	2	2	7	7	5	5	4	2	-
Male	-	1	-	1	1	2	2	1	1	2	-	-
Female	1	1	1	1	1	5	5	4	4	2	2	-
White, both sexes	1	2	1	2	2	7	7	4	5	4	2	-
Male	-	1	-	1	1	2	2	-	1	2	-	-
Female	1	1	1	1	1	5	5	4	4	2	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	1	3	1	4	3	1	1	-	-	-
Male	2	-	1	2	-	1	1	1	1	-	-	-
Female	1	1	-	1	1	3	2	-	-	-	-	-
White, both sexes	3	-	1	3	1	4	3	1	1	-	-	-
Male	2	-	1	2	-	1	1	1	1	-	-	-
Female	1	-	-	1	1	3	2	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
New Hampshire												
All races, both sexes	2	3	-	1	4	5	4	4	2	2	-	-
Male	-	1	-	-	3	2	1	1	-	-	-	-
Female	2	2	-	1	1	3	3	3	2	2	-	-
White, both sexes	1	3	-	1	4	5	4	4	2	2	-	-
Male	-	1	-	-	3	2	1	1	-	-	-	-
Female	1	2	-	1	1	3	3	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	11	11	9	13	18	19	18	12	12	3	-
Male	-	1	3	2	3	6	7	8	3	-	-	-
Female	4	10	8	7	10	12	12	10	9	12	3	-
White, both sexes	2	4	11	6	13	16	17	18	12	11	3	-
Male	-	1	3	2	3	6	6	8	3	-	-	-
Female	2	3	8	4	10	10	11	10	9	11	3	-
Black, both sexes	2	6	-	1	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	6	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	2	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
New Mexico												
All races, both sexes	2	2	2	5	3	4	8	2	7	2	-	-
Male	1	1	1	2	1	1	5	-	2	-	-	-
Female	1	1	1	3	2	3	3	2	5	2	-	-
White, both sexes	2	1	2	4	3	4	8	2	6	2	-	-
Male	1	-	1	1	1	1	5	-	2	-	-	-
Female	1	1	1	3	2	3	3	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	14	6	15	21	18	27	27	19	17	9	3	-
Male	6	3	9	8	5	9	9	5	3	2	-	-
Female	8	3	6	13	13	18	18	14	14	7	3	-
White, both sexes	10	4	8	17	12	25	26	19	16	7	3	-
Male	6	3	4	6	4	9	9	5	3	1	-	-
Female	4	1	4	11	8	16	17	14	13	6	3	-
Black, both sexes	4	2	6	3	5	2	1	-	1	2	-	-
Male	-	-	4	2	-	-	-	-	-	1	-	-
Female	4	2	2	1	5	2	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
North Carolina												
All races, both sexes	12	15	12	10	14	18	22	24	6	5	4	-
Male	5	1	5	4	6	6	3	4	3	2	-	-
Female	7	14	7	6	8	12	19	20	3	3	4	-
White, both sexes	8	10	9	9	12	16	17	20	6	3	3	-
Male	3	-	4	3	5	6	3	3	3	1	-	-
Female	5	10	5	6	7	10	14	17	3	2	3	-
Black, both sexes	3	5	3	1	2	2	5	4	-	1	1	-
Male	2	1	1	1	1	-	-	1	-	-	-	-
Female	1	4	2	-	1	2	5	3	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	1	1	-	-	-	4	3	1	-	-
Male	-	-	1	1	-	-	-	3	-	-	-	-
Female	2	-	-	-	-	-	-	1	3	1	-	-
White, both sexes	2	-	1	1	-	-	-	4	3	1	-	-
Male	-	-	1	1	-	-	-	3	-	-	-	-
Female	2	-	-	-	-	-	-	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Ohio												
All races, both sexes	11	14	15	15	14	24	22	30	32	8	8	-
Male	4	8	4	4	6	9	5	5	8	1	1	-
Female	7	6	11	11	8	15	17	25	24	7	7	-
White, both sexes	9	11	9	13	12	22	21	29	31	7	7	-
Male	4	5	3	4	6	9	5	5	8	1	1	-
Female	5	6	6	9	6	13	16	24	23	6	6	-
Black, both sexes	2	3	6	2	2	2	1	1	1	1	1	-
Male	-	3	1	-	-	-	-	-	-	-	-	-
Female	2	-	5	2	2	2	1	1	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	5	7	1	9	8	5	6	3	-	-	-
Male	1	3	3	1	4	1	4	1	1	-	-	-
Female	2	2	4	-	5	7	1	5	2	-	-	-
White, both sexes	3	4	6	-	8	6	5	6	3	-	-	-
Male	1	3	3	-	4	-	4	1	1	-	-	-
Female	2	1	3	-	4	6	1	5	2	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Oregon												
All races, both sexes	1	6	4	8	8	11	21	11	11	6	2	-
Male	1	4	2	5	1	2	6	4	2	-	-	-
Female	-	2	2	3	7	9	15	7	9	6	2	-
White, both sexes	1	6	3	7	7	10	20	11	11	6	2	-
Male	1	4	1	5	-	2	6	4	2	-	-	-
Female	-	2	2	2	7	8	14	7	9	6	2	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	16	7	17	19	29	22	28	35	15	10	2	-
Male	7	3	4	9	8	6	5	15	3	1	-	-
Female	9	4	13	10	21	16	23	20	12	9	2	-
White, both sexes	13	5	14	16	26	21	25	33	15	8	2	-
Male	6	2	3	9	7	5	4	14	3	1	-	-
Female	7	3	11	7	19	16	21	19	12	7	2	-
Black, both sexes	3	2	3	3	3	1	3	2	-	2	-	-
Male	1	1	1	-	1	1	1	1	-	-	-	-
Female	2	1	2	3	2	-	2	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Rhode Island												
All races, both sexes	1	-	-	1	1	3	1	1	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	1	-	-
White, both sexes	1	-	-	1	1	3	1	1	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	8	5	3	6	7	2	3	3	1	-	-
Male	1	2	-	-	1	3	-	-	2	-	-	-
Female	3	6	5	3	5	4	2	3	1	1	-	-
White, both sexes	2	3	3	3	6	7	2	2	3	1	-	-
Male	-	1	-	-	1	3	-	-	2	-	-	-
Female	2	2	3	3	5	4	2	2	1	1	-	-
Black, both sexes	2	4	2	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	4	2	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
South Dakota												
All races, both sexes	1	-	-	1	1	4	2	3	2	3	1	-
Male	-	-	-	1	-	2	-	1	-	-	1	-
Female	1	-	-	-	1	2	2	2	2	3	-	-
White, both sexes	1	-	-	1	1	3	2	3	2	3	1	-
Male	-	-	-	1	-	2	-	1	-	-	1	-
Female	1	-	-	-	1	1	2	2	2	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	9	7	21	9	14	10	19	4	1	3	-
Male	3	5	3	6	4	4	4	7	-	1	-	-
Female	3	4	4	15	5	10	6	12	4	-	3	-
White, both sexes	4	7	5	15	9	13	10	18	3	1	2	-
Male	2	3	2	4	4	4	4	6	-	1	-	-
Female	2	4	3	11	5	9	6	12	3	-	2	-
Black, both sexes	2	2	2	6	-	1	-	-	1	-	1	-
Male	1	2	1	2	-	-	-	-	-	-	-	-
Female	1	-	1	4	-	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Texas												
All races, both sexes	23	24	29	28	28	36	40	33	20	11	-	-
Male	4	3	10	10	8	9	9	9	2	2	-	-
Female	19	21	19	18	20	27	31	24	18	9	-	-
White, both sexes	17	17	25	25	25	35	36	33	18	9	-	-
Male	3	2	9	10	7	9	7	9	2	2	-	-
Female	14	15	16	15	18	26	29	24	16	7	-	-
Black, both sexes	6	7	3	1	2	1	4	-	2	2	-	-
Male	1	1	1	-	-	-	2	-	-	-	-	-
Female	5	6	2	1	2	1	2	-	2	2	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	2	4	3	1	5	7	5	5	1	-	-
Male	-	2	1	3	-	-	3	1	1	-	-	-
Female	-	-	3	-	1	5	4	4	4	1	-	-
White, both sexes	-	2	4	3	1	5	7	5	5	1	-	-
Male	-	2	1	3	-	-	3	1	1	-	-	-
Female	-	-	3	-	1	5	4	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Vermont												
All races, both sexes	-	1	-	1	3	1	1	4	4	-	1	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	1	2	1	1	3	4	-	1	-
White, both sexes	-	1	-	1	3	1	1	4	4	-	1	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	1	-	1	2	1	1	3	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	8	16	17	14	12	22	8	20	6	2	-
Male	4	1	9	4	4	2	4	4	1	2	-	-
Female	3	7	7	13	10	10	18	4	19	4	2	-
White, both sexes	4	5	10	10	14	10	18	6	16	4	2	-
Male	2	1	5	3	4	-	3	3	1	1	-	-
Female	2	4	5	7	10	10	15	3	15	3	2	-
Black, both sexes	3	3	6	6	-	1	4	2	4	2	-	-
Male	2	-	4	1	-	1	1	1	-	1	-	-
Female	1	3	2	5	-	-	3	1	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Washington												
All races, both sexes	4	8	7	10	13	11	21	14	9	4	1	-
Male	2	3	2	2	3	3	9	2	-	1	-	-
Female	2	5	5	8	10	8	12	12	9	3	1	-
White, both sexes	3	6	7	8	12	10	21	14	9	4	1	-
Male	2	2	2	2	3	3	9	2	-	1	-	-
Female	1	4	5	6	9	7	12	12	9	3	1	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	1	-	-	4	4	5	5	4	1	-	-
Male	1	1	-	-	-	-	1	1	1	-	-	-
Female	2	-	-	-	4	4	4	4	3	1	-	-
White, both sexes	3	1	-	-	4	4	5	4	4	1	-	-
Male	1	1	-	-	-	-	1	1	1	-	-	-
Female	2	-	-	-	4	4	4	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Wisconsin												
All races, both sexes	8	3	6	8	10	6	16	21	11	3	4	-
Male	4	-	1	1	4	4	4	8	4	2	-	-
Female	4	3	5	7	6	2	12	13	7	1	4	-
White, both sexes	8	3	6	8	10	6	16	19	11	3	4	-
Male	4	-	1	1	4	4	4	8	4	2	-	-
Female	4	3	5	7	6	2	12	11	7	1	4	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	1	-	-	2	3	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	3	1	1	-	-	-
White, both sexes	1	-	1	-	-	2	3	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
United States												
All races, both sexes	1,115	1	1	2	7	10	7	6	7	9	12	24
Male	479	1	-	-	5	3	2	1	5	6	5	13
Female	636	-	1	2	2	7	5	5	2	3	7	11
White, both sexes	982	1	-	2	6	10	5	5	5	7	10	20
Male	411	1	-	-	4	3	2	1	4	4	3	12
Female	571	-	-	2	2	7	3	4	1	3	7	8
Black, both sexes	113	-	1	-	1	-	1	-	1	2	2	4
Male	61	-	-	-	1	-	-	-	1	2	2	1
Female	52	-	1	-	-	-	1	-	-	-	-	3
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Alabama												
All races, both sexes	9	-	-	1	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	1	-	1	-	-	-	-	-	1
White, both sexes	9	-	-	1	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	1	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	21	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Arkansas												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	90	-	-	-	-	-	-	1	3	2	-	4
Male	40	-	-	-	-	-	-	-	2	1	-	3
Female	50	-	-	-	-	-	-	1	1	1	-	1
White, both sexes	79	-	-	-	-	-	-	1	2	2	-	3
Male	35	-	-	-	-	-	-	-	1	1	-	3
Female	44	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Colorado												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Florida												
All races, both sexes	61	-	-	-	1	1	1	-	-	1	2	1
Male	32	-	-	-	-	-	1	-	-	1	-	1
Female	29	-	-	-	1	1	-	-	-	-	2	-
White, both sexes	59	-	-	-	1	1	1	-	-	1	2	1
Male	30	-	-	-	-	-	1	-	-	1	-	1
Female	29	-	-	-	1	1	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Illinois												
All races, both sexes	49	-	-	-	-	2	-	-	-	1	-	1
Male	22	-	-	-	-	1	-	-	-	-	-	1
Female	27	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	41	-	-	-	-	2	-	-	-	1	-	1
Male	19	-	-	-	-	1	-	-	-	-	-	1
Female	22	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Iowa												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Kentucky												
All races, both sexes	16	-	-	-	1	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	16	-	-	-	1	1	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Maine												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	21	-	-	-	-	1	1	-	-	-	-	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	11	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	1	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Massachusetts												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	48	-	-	-	-	1	-	-	1	2	-	1
Male	23	-	-	-	-	1	-	-	-	2	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	39	-	-	-	-	1	-	-	-	1	-	1
Male	17	-	-	-	-	1	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Minnesota												
All races, both sexes	31	-	-	-	1	-	-	1	1	-	-	-
Male	15	-	-	-	1	-	-	-	1	-	-	-
Female	16	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	30	-	-	-	1	-	-	1	1	-	-	-
Male	15	-	-	-	1	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	8	-	-	-	-	-	1	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Missouri												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Nebraska												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	30	-	-	1	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	1	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
New Mexico												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	59	-	-	-	-	-	1	1	-	-	2	1
Male	17	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	52	-	-	-	-	-	1	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	38	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
North Carolina												
All races, both sexes	34	-	-	-	-	-	-	-	1	-	1	1
Male	14	-	-	-	-	-	-	-	1	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Ohio												
All races, both sexes	58	-	-	-	1	-	-	1	-	-	2	2
Male	30	-	-	-	1	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	54	-	-	-	1	-	-	1	-	-	2	2
Male	27	-	-	-	1	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Oregon												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	64	-	-	-	-	-	1	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	58	-	-	-	-	-	1	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	22	-	-	-	1	-	-	-	-	-	-	1
Male	9	-	-	-	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	1	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Texas												
All races, both sexes	66	-	-	-	-	3	-	-	-	3	1	2
Male	31	-	-	-	-	1	-	-	-	2	1	1
Female	35	-	-	-	-	2	-	-	-	1	-	1
White, both sexes	59	-	-	-	-	3	-	-	-	2	-	2
Male	27	-	-	-	-	1	-	-	-	1	-	1
Female	32	-	-	-	-	2	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	22	1	-	-	1	-	-	-	-	-	-	-
Male	13	1	-	-	1	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	1	-	-	1	-	-	-	-	-	-	-
Male	11	1	-	-	1	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Washington												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
Wisconsin												
All races, both sexes	27	-	1	-	1	-	-	-	-	-	2	-
Male	9	-	-	-	1	-	-	-	-	-	1	-
Female	18	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
United States												
All races, both sexes	39	67	58	65	111	147	203	183	109	42	5	-
Male	26	45	25	32	56	78	76	62	27	11	-	-
Female	13	22	33	33	55	69	127	121	82	31	5	-
White, both sexes	28	52	44	53	91	134	188	175	103	38	5	-
Male	18	32	17	25	46	68	74	60	27	10	-	-
Female	10	20	27	28	45	66	114	115	76	28	5	-
Black, both sexes	9	15	13	10	14	12	14	7	4	3	-	-
Male	6	13	8	6	8	10	1	1	-	1	-	-
Female	3	2	5	4	6	2	13	6	4	2	-	-
American Indian 1/, both sexes	1	-	-	1	3	-	1	-	-	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	3	1	-	1	2	1	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	3	1	-	-	2	1	-	-
Alabama												
All races, both sexes	-	-	-	1	2	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
White, both sexes	-	-	-	1	2	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Alaska												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	1	1	-	5	1	3	5	2	-	-	-
Male	2	1	-	-	1	-	1	2	-	-	-	-
Female	-	-	1	-	4	1	2	3	2	-	-	-
White, both sexes	1	1	1	-	4	1	2	5	2	-	-	-
Male	1	1	-	-	1	-	-	2	-	-	-	-
Female	-	-	1	-	3	1	2	3	2	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Arkansas												
All races, both sexes	-	2	-	-	2	1	1	2	1	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	2	1	1	-	-
White, both sexes	-	2	-	-	2	1	1	2	1	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	8	4	6	11	8	10	13	11	3	1	-
Male	4	4	-	3	4	4	3	6	5	1	-	-
Female	1	4	4	3	7	4	7	7	6	2	1	-
White, both sexes	3	7	3	4	9	7	10	13	11	3	1	-
Male	2	3	-	2	4	4	3	6	5	1	-	-
Female	1	4	3	2	5	3	7	7	6	2	1	-
Black, both sexes	-	1	-	2	1	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Colorado												
All races, both sexes	1	1	4	1	1	2	6	2	6	2	-	-
Male	-	1	2	1	-	-	2	1	1	-	-	-
Female	1	-	2	-	1	2	4	1	5	2	-	-
White, both sexes	-	1	4	1	1	2	5	2	6	2	-	-
Male	-	1	2	1	-	-	2	1	1	-	-	-
Female	-	-	2	-	1	2	3	1	5	2	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	1	1	5	2	2	1	-	-
Male	-	-	-	1	-	1	3	-	1	-	-	-
Female	-	-	-	-	1	-	2	2	1	1	-	-
White, both sexes	-	-	-	-	1	-	5	2	2	1	-	-
Male	-	-	-	-	-	-	3	-	1	-	-	-
Female	-	-	-	-	1	-	2	2	1	1	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Florida												
All races, both sexes	2	1	1	4	7	11	10	15	3	-	-	-
Male	2	1	1	2	5	6	4	8	-	-	-	-
Female	-	-	-	2	2	5	6	7	3	-	-	-
White, both sexes	1	1	1	4	6	11	10	15	3	-	-	-
Male	1	1	1	2	4	6	4	8	-	-	-	-
Female	-	-	-	2	2	5	6	7	3	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	2	4	3	5	3	4	5	-	2	1	-
Male	1	2	2	3	2	1	1	1	-	1	-	-
Female	-	-	2	-	3	2	3	4	-	1	1	-
White, both sexes	-	1	2	3	1	1	3	4	-	2	1	-
Male	-	1	1	3	1	-	1	1	-	1	-	-
Female	-	-	1	-	-	1	2	3	-	1	1	-
Black, both sexes	1	1	2	-	3	2	1	1	-	-	-	-
Male	1	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	2	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Hawaii												
All races, both sexes	-	-	-	1	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Idaho												
All races, both sexes	-	1	-	-	-	-	-	3	2	2	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	2	2	-	-
White, both sexes	-	1	-	-	-	-	-	3	2	2	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Illinois												
All races, both sexes	2	1	3	4	3	8	14	4	2	3	1	-
Male	1	1	-	2	2	6	6	-	1	1	-	-
Female	1	-	3	2	1	2	8	4	1	2	1	-
White, both sexes	-	1	2	4	3	7	12	2	2	3	1	-
Male	-	1	-	2	2	5	5	-	1	1	-	-
Female	-	-	2	2	1	2	7	2	1	2	1	-
Black, both sexes	2	-	1	-	-	1	2	2	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	2	2	5	1	1	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	5	1	1	-	-	-
White, both sexes	1	-	-	-	2	2	4	1	1	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Iowa												
All races, both sexes	1	-	2	2	-	1	6	4	2	-	-	-
Male	-	-	1	1	-	1	3	2	1	-	-	-
Female	1	-	1	1	-	-	3	2	1	-	-	-
White, both sexes	1	-	2	2	-	1	6	4	2	-	-	-
Male	-	-	1	1	-	1	3	2	1	-	-	-
Female	1	-	1	1	-	-	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	1	2	2	3	1	1	-	-
Male	-	-	-	-	-	2	2	-	-	1	-	-
Female	-	-	-	1	1	-	-	3	1	-	-	-
White, both sexes	-	-	-	1	1	1	2	3	1	1	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	-	1	1	-	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Kentucky												
All races, both sexes	2	-	1	-	2	2	1	4	1	-	1	-
Male	1	-	-	-	-	2	-	2	-	-	-	-
Female	1	-	1	-	2	-	1	2	1	-	1	-
White, both sexes	2	-	1	-	2	2	1	4	1	-	1	-
Male	1	-	-	-	-	2	-	2	-	-	-	-
Female	1	-	1	-	2	-	1	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	1	3	1	2	-	4	-	-	-
Male	-	-	-	1	2	1	-	-	2	-	-	-
Female	-	-	-	-	1	-	2	-	2	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	3	-	-	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	1	3	-	1	-	1	-	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Maine												
All races, both sexes	-	2	-	-	-	-	3	1	1	2	-	-
Male	-	1	-	-	-	-	1	1	1	1	-	-
Female	-	1	-	-	-	-	2	-	-	1	-	-
White, both sexes	-	2	-	-	-	-	3	1	1	2	-	-
Male	-	1	-	-	-	-	1	1	1	1	-	-
Female	-	1	-	-	-	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	2	-	1	4	3	2	4	1	-	-
Male	1	-	1	-	-	4	2	-	1	-	-	-
Female	-	1	1	-	1	-	1	2	3	1	-	-
White, both sexes	-	-	1	-	1	3	3	2	4	1	-	-
Male	-	-	1	-	-	3	2	-	1	-	-	-
Female	-	-	-	-	1	-	1	2	3	1	-	-
Black, both sexes	1	1	1	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Massachusetts												
All races, both sexes	-	-	-	1	2	2	6	5	7	-	-	-
Male	-	-	-	-	2	1	2	3	4	-	-	-
Female	-	-	-	1	-	1	4	2	3	-	-	-
White, both sexes	-	-	-	1	2	2	6	5	7	-	-	-
Male	-	-	-	-	2	1	2	3	4	-	-	-
Female	-	-	-	1	-	1	4	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	10	2	1	4	5	10	8	2	1	-	-
Male	-	6	1	-	4	2	5	1	1	-	-	-
Female	-	4	1	1	-	3	5	7	1	1	-	-
White, both sexes	-	5	1	1	4	4	10	8	2	1	-	-
Male	-	2	-	-	4	2	5	1	1	-	-	-
Female	-	3	1	1	-	2	5	7	1	1	-	-
Black, both sexes	-	5	1	-	-	1	-	-	-	-	-	-
Male	-	4	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Minnesota												
All races, both sexes	-	1	1	1	1	6	8	3	5	2	-	-
Male	-	1	-	1	1	3	4	1	-	2	-	-
Female	-	-	1	-	-	3	4	2	5	-	-	-
White, both sexes	-	1	1	1	1	6	7	3	5	2	-	-
Male	-	1	-	1	1	3	4	1	-	2	-	-
Female	-	-	1	-	-	3	3	2	5	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	-	1	2	2	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Missouri												
All races, both sexes	1	1	1	2	3	3	4	4	3	1	-	-
Male	-	1	1	1	2	3	-	1	-	1	-	-
Female	1	-	-	1	1	-	4	3	3	-	-	-
White, both sexes	1	1	-	1	1	3	4	4	2	-	-	-
Male	-	1	-	-	1	3	-	1	-	-	-	-
Female	1	-	-	1	-	-	4	3	2	-	-	-
Black, both sexes	-	-	1	1	2	-	-	-	1	1	-	-
Male	-	-	1	1	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	2	1	1	2	1	-	-
Male	-	-	-	1	-	2	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	1	-	2	1	1	2	1	-	-
Male	-	-	-	1	-	2	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Nebraska												
All races, both sexes	-	-	1	1	1	2	5	1	1	-	1	-
Male	-	-	-	1	1	2	3	-	1	-	-	-
Female	-	-	1	-	-	-	2	1	-	-	1	-
White, both sexes	-	-	1	1	1	2	5	1	1	-	1	-
Male	-	-	-	1	1	2	3	-	1	-	-	-
Female	-	-	1	-	-	-	2	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	2	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	2	1	-	-	-	-
White, both sexes	-	-	-	2	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
New Hampshire												
All races, both sexes	-	1	-	-	1	1	-	-	2	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	1	-	-	1	1	-	-	2	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	1	-	3	1	7	6	3	2	-	-
Male	1	1	-	-	1	1	4	2	2	-	-	-
Female	2	1	1	-	2	-	3	4	1	2	-	-
White, both sexes	3	2	1	-	3	1	6	6	3	2	-	-
Male	1	1	-	-	1	1	4	2	2	-	-	-
Female	2	1	1	-	2	-	2	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
New Mexico												
All races, both sexes	-	-	-	-	3	2	2	2	1	1	-	-
Male	-	-	-	-	1	2	-	1	-	-	-	-
Female	-	-	-	-	2	-	2	1	1	1	-	-
White, both sexes	-	-	-	-	2	2	2	2	1	1	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	2	-	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	5	3	7	5	9	9	8	7	1	-	-
Male	-	4	1	2	2	2	2	2	-	-	-	-
Female	-	1	2	5	3	7	7	6	7	1	-	-
White, both sexes	-	3	3	4	5	9	8	8	7	1	-	-
Male	-	2	1	1	2	2	2	2	-	-	-	-
Female	-	1	2	3	3	7	6	6	7	1	-	-
Black, both sexes	-	2	-	2	-	-	1	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
North Carolina												
All races, both sexes	-	3	3	2	3	6	7	4	2	1	-	-
Male	-	2	3	-	2	3	2	-	-	-	-	-
Female	-	1	-	2	1	3	5	4	2	1	-	-
White, both sexes	-	2	1	1	2	4	6	3	1	1	-	-
Male	-	1	1	-	1	1	2	-	-	-	-	-
Female	-	1	-	1	1	3	4	3	1	1	-	-
Black, both sexes	-	1	2	1	1	2	1	1	1	-	-	-
Male	-	1	2	-	1	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Ohio												
All races, both sexes	1	5	4	1	8	13	6	8	2	4	-	-
Male	1	4	3	-	5	5	3	5	-	1	-	-
Female	-	1	1	1	3	8	3	3	2	3	-	-
White, both sexes	1	5	3	1	7	12	6	8	2	3	-	-
Male	1	4	2	-	4	4	3	5	-	1	-	-
Female	-	1	1	1	3	8	3	3	2	2	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	1	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	2	-	4	2	6	3	1	1	-	-	-
Male	1	1	-	3	1	4	-	-	-	-	-	-
Female	-	1	-	1	1	2	3	1	1	-	-	-
White, both sexes	1	2	-	4	2	5	3	1	1	-	-	-
Male	1	1	-	3	1	3	-	-	-	-	-	-
Female	-	1	-	1	1	2	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Oregon												
All races, both sexes	-	1	1	3	2	1	3	6	4	-	-	-
Male	-	1	-	1	1	1	2	1	-	-	-	-
Female	-	-	1	2	1	-	1	5	4	-	-	-
White, both sexes	-	1	1	3	1	1	3	5	4	-	-	-
Male	-	1	-	1	1	1	2	-	-	-	-	-
Female	-	-	1	2	-	-	1	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	5	4	3	9	8	13	12	2	2	-	-
Male	3	3	1	1	5	3	4	3	-	-	-	-
Female	-	2	3	2	4	5	9	9	2	2	-	-
White, both sexes	3	4	3	3	8	8	11	12	2	1	-	-
Male	3	2	1	1	4	3	4	3	-	-	-	-
Female	-	2	2	2	4	5	7	9	2	1	-	-
Black, both sexes	-	1	1	-	1	-	2	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	2	1	-	2	2	4	2	2	-	-	-
Male	1	1	-	-	1	2	1	1	-	-	-	-
Female	1	1	1	-	1	-	3	1	2	-	-	-
White, both sexes	1	1	-	-	2	2	3	-	2	-	-	-
Male	1	-	-	-	1	2	1	-	-	-	-	-
Female	-	1	-	-	1	-	2	-	2	-	-	-
Black, both sexes	1	1	1	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
South Dakota												
All races, both sexes	-	-	-	1	1	2	3	3	-	-	-	-
Male	-	-	-	1	1	-	2	2	-	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	2	3	3	-	-	-	-
Male	-	-	-	1	1	-	2	2	-	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	2	5	-	5	3	4	-	-	-	-
Male	-	-	1	3	-	2	-	2	-	-	-	-
Female	-	1	1	2	-	3	3	2	-	-	-	-
White, both sexes	-	1	2	4	-	4	3	4	-	-	-	-
Male	-	-	1	2	-	1	-	2	-	-	-	-
Female	-	1	1	2	-	3	3	2	-	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Texas												
All races, both sexes	4	4	4	2	7	6	10	12	6	2	-	-
Male	3	3	3	1	3	1	6	3	2	1	-	-
Female	1	1	1	1	4	5	4	9	4	1	-	-
White, both sexes	3	4	3	2	6	6	10	11	5	2	-	-
Male	2	3	2	1	3	1	6	3	2	1	-	-
Female	1	1	1	1	3	5	4	8	3	1	-	-
Black, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Utah												
All races, both sexes	1	1	1	-	1	1	2	4	1	1	-	-
Male	1	-	1	-	-	-	-	2	-	1	-	-
Female	-	1	-	-	1	1	2	2	1	-	-	-
White, both sexes	1	1	1	-	1	1	2	4	1	1	-	-
Male	1	-	1	-	-	-	-	2	-	1	-	-
Female	-	1	-	-	1	1	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	1	2	-	4	3	7	2	-	-	-
Male	1	-	1	2	-	2	2	3	-	-	-	-
Female	-	-	-	-	-	2	1	4	2	-	-	-
White, both sexes	1	-	-	1	-	4	3	7	2	-	-	-
Male	1	-	-	1	-	2	2	3	-	-	-	-
Female	-	-	-	-	-	2	1	4	2	-	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Washington												
All races, both sexes	3	1	3	1	2	1	3	7	4	1	-	-
Male	1	1	1	-	2	-	-	3	1	-	-	-
Female	2	-	2	1	-	1	3	4	3	1	-	-
White, both sexes	3	1	3	1	2	1	3	7	4	1	-	-
Male	1	1	1	-	2	-	-	3	1	-	-	-
Female	2	-	2	1	-	1	3	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	1	-	-	2	4	1	-	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	1	-	1	-	-	-	3	1	-	1	-	-
White, both sexes	1	-	1	-	-	2	4	1	-	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	1	-	1	-	-	-	3	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Wisconsin												
All races, both sexes	-	-	-	-	2	4	5	5	5	2	-	-
Male	-	-	-	-	-	2	3	1	1	-	-	-
Female	-	-	-	-	2	2	2	4	4	2	-	-
White, both sexes	-	-	-	-	2	4	5	5	4	2	-	-
Male	-	-	-	-	-	2	3	1	1	-	-	-
Female	-	-	-	-	2	2	2	4	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
United States												
All races, both sexes	1,277	7	3	1	5	4	4	12	15	15	50	69
Male	674	6	1	1	2	3	4	9	9	12	31	38
Female	603	1	2	-	3	1	-	3	6	3	19	31
White, both sexes	1,072	5	2	1	4	4	3	10	9	12	40	53
Male	553	4	1	1	1	3	3	7	4	11	26	27
Female	519	1	1	-	3	1	-	3	5	1	14	26
Black, both sexes	173	2	1	-	1	-	1	2	5	3	9	15
Male	96	2	-	-	1	-	1	2	4	1	5	10
Female	77	-	1	-	-	-	-	-	1	2	4	5
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	25	-	1	-	1	-	-	-	1	-	1	4
Male	17	-	1	-	1	-	-	-	1	-	-	4
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	1	-	1	-	-	-	-	-	-	4
Male	14	-	1	-	1	-	-	-	-	-	-	4
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Arkansas												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	131	1	-	-	2	-	-	2	1	2	9	9
Male	74	-	-	-	1	-	-	1	1	2	7	6
Female	57	1	-	-	1	-	-	1	-	-	2	3
White, both sexes	110	1	-	-	1	-	-	2	1	2	8	8
Male	59	-	-	-	-	-	-	1	1	2	6	5
Female	51	1	-	-	1	-	-	1	-	-	2	3
Black, both sexes	10	-	-	-	1	-	-	-	-	-	1	-
Male	6	-	-	-	1	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Colorado												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Florida												
All races, both sexes	59	-	-	-	-	-	1	-	1	-	3	1
Male	28	-	-	-	-	-	1	-	1	-	2	-
Female	31	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	46	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	13	-	-	-	-	-	1	-	1	-	1	1
Male	9	-	-	-	-	-	1	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	34	-	-	-	-	-	-	-	3	-	1	3
Male	24	-	-	-	-	-	-	-	2	-	1	1
Female	10	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	21	-	-	-	-	-	-	-	1	-	1	1
Male	15	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	13	-	-	-	-	-	-	-	2	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Hawaii												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Illinois												
All races, both sexes	64	2	1	-	-	1	-	1	-	1	1	4
Male	28	2	-	-	-	-	-	1	-	-	-	-
Female	36	-	1	-	-	1	-	-	-	1	1	4
White, both sexes	55	1	1	-	-	1	-	1	-	-	1	3
Male	25	1	-	-	-	-	-	1	-	-	-	-
Female	30	-	1	-	-	1	-	-	-	-	1	3
Black, both sexes	8	1	-	-	-	-	-	-	-	1	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	35	-	-	-	-	-	-	-	-	2	-	2
Male	22	-	-	-	-	-	-	-	-	2	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Iowa												
All races, both sexes	12	-	-	-	1	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	1	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Kentucky												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Maine												
All races, both sexes	16	-	-	-	-	1	-	-	1	-	-	-
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	16	-	-	-	-	1	-	-	1	-	-	-
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	33	-	-	-	-	-	-	1	-	-	3	2
Male	17	-	-	-	-	-	-	1	-	-	2	1
Female	16	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	1	-	-	2	1
Male	11	-	-	-	-	-	-	1	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Massachusetts												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	3
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	40	-	-	-	-	-	-	-	-	2	1	3
Male	23	-	-	-	-	-	-	-	-	2	1	3
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	2	-	1
Male	19	-	-	-	-	-	-	-	-	2	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Minnesota												
All races, both sexes	20	-	1	-	1	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	1	-	1	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	1	-	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Missouri												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	26	-	-	-	-	-	-	-	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Nebraska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
New Hampshire												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	31	-	-	-	-	-	-	1	-	1	1	-
Male	14	-	-	-	-	-	-	1	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	1	-	1	1	-
Male	12	-	-	-	-	-	-	1	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
New Mexico												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	70	-	-	-	-	-	1	-	2	1	1	6
Male	38	-	-	-	-	-	1	-	2	-	1	4
Female	32	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	65	-	-	-	-	-	1	-	1	1	1	6
Male	35	-	-	-	-	-	1	-	1	-	1	4
Female	30	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
North Carolina												
All races, both sexes	37	-	-	-	-	-	-	-	3	-	2	5
Male	20	-	-	-	-	-	-	-	1	-	1	4
Female	17	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	30	-	-	-	-	-	-	-	3	-	2	3
Male	18	-	-	-	-	-	-	-	1	-	1	3
Female	12	-	-	-	-	-	-	-	2	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Ohio												
All races, both sexes	60	-	-	-	-	1	-	1	1	2	2	1
Male	27	-	-	-	-	1	-	1	-	1	1	-
Female	33	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	53	-	-	-	-	1	-	1	1	1	2	1
Male	25	-	-	-	-	1	-	1	-	1	1	-
Female	28	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Oregon												
All races, both sexes	15	1	-	-	-	1	-	-	-	-	-	1
Male	9	1	-	-	-	1	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	1	-	-	-	1	-	-	-	-	-	1
Male	9	1	-	-	-	1	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	61	-	-	1	-	-	-	-	-	-	2	3
Male	31	-	-	1	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	58	-	-	1	-	-	-	-	-	-	2	3
Male	29	-	-	1	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Rhode Island												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	3	4
Male	20	-	-	-	-	-	-	-	-	-	1	2
Female	16	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	2	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	30	1	-	-	-	-	-	-	-	-	1	-
Male	16	1	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	1	-	-	-	-	-	-	-	-	1	-
Male	14	1	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Texas												
All races, both sexes	78	1	-	-	-	-	1	3	1	-	2	6
Male	38	1	-	-	-	-	1	2	-	-	-	3
Female	40	-	-	-	-	-	-	1	1	-	2	3
White, both sexes	67	1	-	-	-	-	1	2	1	-	1	6
Male	35	1	-	-	-	-	1	1	-	-	-	3
Female	32	-	-	-	-	-	-	1	1	-	1	3
Black, both sexes	10	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	34	-	-	-	-	-	1	-	1	-	1	-
Male	16	-	-	-	-	-	1	-	1	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	1	-	-	-	1	-
Male	12	-	-	-	-	-	1	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Washington												
All races, both sexes	37	1	-	-	-	-	-	1	-	1	-	2
Male	20	1	-	-	-	-	-	1	-	1	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	1	-	1	-	1
Male	17	-	-	-	-	-	-	1	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	1	-	-	-	-	-	-	-	-	-	1
Male	3	1	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Wisconsin												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
United States												
All races, both sexes	92	68	98	95	123	162	180	150	82	37	5	-
Male	59	36	56	49	64	86	82	82	33	10	1	-
Female	33	32	42	46	59	76	98	68	49	27	4	-
White, both sexes	67	50	82	74	99	138	164	139	78	33	5	-
Male	40	26	44	37	49	75	78	73	33	9	1	-
Female	27	24	38	37	50	63	86	66	45	24	4	-
Black, both sexes	16	13	13	19	20	22	15	9	4	3	-	-
Male	10	7	9	10	12	10	3	8	-	1	-	-
Female	6	6	4	9	8	12	12	1	4	2	-	-
American Indian 1/, both sexes	3	3	2	-	1	-	-	1	-	-	-	-
Male	3	2	2	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	2	1	2	3	2	1	1	-	1	-	-
Male	6	1	1	2	3	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-
Alabama												
All races, both sexes	-	2	4	3	2	2	3	1	-	-	-	-
Male	-	1	3	2	1	1	1	1	-	-	-	-
Female	-	1	1	1	1	1	2	-	-	-	-	-
White, both sexes	-	1	2	2	2	1	3	-	-	-	-	-
Male	-	1	1	1	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	2	-	-	-	-	-
Black, both sexes	-	1	2	1	-	1	-	1	-	-	-	-
Male	-	-	2	1	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	1	4	3	4	1	1	1	-	-	-
Male	-	-	1	3	2	2	-	1	1	-	-	-
Female	-	2	-	1	1	2	1	-	-	-	-	-
White, both sexes	-	2	1	3	2	4	1	1	1	-	-	-
Male	-	-	1	2	1	2	-	1	1	-	-	-
Female	-	2	-	1	1	2	1	-	-	-	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Arkansas												
All races, both sexes	1	-	1	1	-	1	1	-	2	1	-	-
Male	1	-	-	-	-	1	-	-	1	1	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
White, both sexes	1	-	1	1	-	1	1	-	2	1	-	-
Male	1	-	-	-	-	1	-	-	1	1	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	18	7	13	8	14	13	10	13	4	5	-	-
Male	15	4	8	4	5	5	4	10	-	1	-	-
Female	3	3	5	4	9	8	6	3	4	4	-	-
White, both sexes	15	3	11	7	11	11	9	12	4	4	-	-
Male	12	1	6	3	3	5	4	9	-	1	-	-
Female	3	2	5	4	8	6	5	3	4	3	-	-
Black, both sexes	1	2	1	-	1	1	1	-	-	1	-	-
Male	1	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	1	-	1	-	-	-	-
Male	1	1	-	1	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Colorado												
All races, both sexes	2	3	-	-	2	1	4	4	2	-	-	-
Male	2	2	-	-	2	-	1	2	1	-	-	-
Female	-	1	-	-	-	1	3	2	1	-	-	-
White, both sexes	2	3	-	-	2	1	4	4	2	-	-	-
Male	2	2	-	-	2	-	1	2	1	-	-	-
Female	-	1	-	-	-	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	2	-	1	-	1	2	3	1	-	-
Male	-	-	1	-	-	-	1	2	1	-	-	-
Female	-	-	1	-	1	-	-	-	2	1	-	-
White, both sexes	-	-	2	-	1	-	1	2	3	1	-	-
Male	-	-	1	-	-	-	1	2	1	-	-	-
Female	-	-	1	-	1	-	-	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Delaware												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Florida												
All races, both sexes	3	1	-	10	5	10	10	7	6	1	-	-
Male	2	1	-	4	4	4	4	4	1	-	-	-
Female	1	-	-	6	1	6	6	3	5	1	-	-
White, both sexes	3	1	-	6	4	6	10	7	6	1	-	-
Male	2	1	-	2	3	1	4	4	1	-	-	-
Female	1	-	-	4	1	5	6	3	5	1	-	-
Black, both sexes	-	-	-	4	1	4	-	-	-	-	-	-
Male	-	-	-	2	1	3	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	-	2	6	4	3	2	3	1	1	-	-
Male	5	-	1	4	4	2	2	1	1	-	-	-
Female	-	-	1	2	-	1	-	2	-	1	-	-
White, both sexes	3	-	2	3	2	2	1	3	1	1	-	-
Male	3	-	1	3	2	1	1	1	1	-	-	-
Female	-	-	1	-	-	1	-	2	-	1	-	-
Black, both sexes	2	-	-	3	2	1	1	-	-	-	-	-
Male	2	-	-	1	2	1	1	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Hawaii												
All races, both sexes	3	2	1	1	-	1	1	-	-	1	-	-
Male	3	1	1	1	-	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	-	-	1	1	-	-	1	-	-
Male	3	-	1	-	-	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Illinois												
All races, both sexes	3	4	6	4	4	10	5	11	6	-	-	-
Male	1	2	1	-	3	7	2	6	3	-	-	-
Female	2	2	5	4	1	3	3	5	3	-	-	-
White, both sexes	3	3	4	3	3	9	5	11	6	-	-	-
Male	1	1	1	-	2	7	2	6	3	-	-	-
Female	2	2	3	3	1	2	3	5	3	-	-	-
Black, both sexes	-	1	2	1	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	3	2	2	2	6	4	8	1	1	1	-
Male	1	1	-	2	2	5	2	4	1	-	-	-
Female	-	2	2	-	-	1	2	4	-	1	1	-
White, both sexes	-	3	2	2	1	6	4	8	1	1	1	-
Male	-	1	-	2	1	5	2	4	1	-	-	-
Female	-	2	2	-	-	1	2	4	-	1	1	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Iowa												
All races, both sexes	-	-	-	1	2	1	3	2	1	-	-	-
Male	-	-	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	3	-	1	-	-	-
White, both sexes	-	-	-	1	2	1	3	2	1	-	-	-
Male	-	-	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	2	-	1	-	3	3	1	-	1	-
Male	1	-	2	-	-	-	3	2	1	-	-	-
Female	1	-	-	-	1	-	-	1	-	-	1	-
White, both sexes	2	-	2	-	-	-	3	3	1	-	1	-
Male	1	-	2	-	-	-	3	2	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Kentucky												
All races, both sexes	4	1	-	4	3	4	9	2	2	3	-	-
Male	3	-	-	2	1	3	2	-	2	2	-	-
Female	1	1	-	2	2	1	7	2	-	1	-	-
White, both sexes	2	1	-	3	3	4	9	2	2	2	-	-
Male	2	-	-	2	1	3	2	-	2	2	-	-
Female	-	1	-	1	2	1	7	2	-	-	-	-
Black, both sexes	2	-	-	1	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	2	2	-	7	1	5	2	1	2	-	-
Male	-	1	-	-	5	-	4	1	1	1	-	-
Female	-	1	2	-	2	1	1	1	-	1	-	-
White, both sexes	-	-	2	-	5	-	4	1	1	1	-	-
Male	-	-	-	-	4	-	3	1	1	-	-	-
Female	-	-	2	-	1	-	1	-	-	1	-	-
Black, both sexes	-	2	-	-	2	1	1	1	-	1	-	-
Male	-	1	-	-	1	-	1	-	-	1	-	-
Female	-	1	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Maine												
All races, both sexes	1	-	1	3	1	2	3	-	1	1	1	-
Male	1	-	1	-	1	1	3	-	-	-	-	-
Female	-	-	-	3	-	1	-	-	1	1	1	-
White, both sexes	1	-	1	3	1	2	3	-	1	1	1	-
Male	1	-	1	-	1	1	3	-	-	-	-	-
Female	-	-	-	3	-	1	-	-	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	1	4	1	7	2	1	6	1	-	-	-
Male	3	-	4	-	2	-	-	4	-	-	-	-
Female	1	1	-	1	5	2	1	2	1	-	-	-
White, both sexes	1	1	2	-	6	2	-	3	1	-	-	-
Male	-	-	2	-	2	-	-	1	-	-	-	-
Female	1	1	-	-	4	2	-	2	1	-	-	-
Black, both sexes	3	-	2	1	1	-	1	3	-	-	-	-
Male	3	-	2	-	-	-	-	3	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Massachusetts												
All races, both sexes	-	2	-	-	3	1	6	1	-	1	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	1	-	-	2	-	5	1	-	1	-	-
White, both sexes	-	2	-	-	3	1	6	1	-	1	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	1	-	-	2	-	5	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	3	3	3	1	5	8	5	3	1	-	-
Male	-	3	1	3	-	3	3	2	1	1	-	-
Female	2	-	2	-	1	2	5	3	2	-	-	-
White, both sexes	2	3	2	3	1	4	5	5	1	1	-	-
Male	-	3	1	3	-	2	3	2	1	1	-	-
Female	2	-	1	-	1	2	2	3	-	-	-	-
Black, both sexes	-	-	1	-	-	1	3	-	2	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Minnesota												
All races, both sexes	1	1	1	2	3	3	2	-	1	1	1	-
Male	1	-	-	1	1	3	1	-	-	1	-	-
Female	-	1	1	1	2	-	1	-	1	-	1	-
White, both sexes	1	1	1	1	3	3	2	-	1	1	1	-
Male	1	-	-	-	1	3	1	-	-	1	-	-
Female	-	1	1	1	2	-	1	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	1	1	2	3	2	1	4	-	-	-	-
Male	-	-	-	1	3	-	-	3	-	-	-	-
Female	3	1	1	1	-	2	1	1	-	-	-	-
White, both sexes	1	-	1	1	2	1	1	3	-	-	-	-
Male	-	-	-	-	2	-	-	2	-	-	-	-
Female	1	-	1	1	-	1	1	1	-	-	-	-
Black, both sexes	2	1	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	2	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Missouri												
All races, both sexes	-	1	2	1	3	5	7	2	1	4	-	-
Male	-	1	2	-	3	3	3	-	-	1	-	-
Female	-	-	-	1	-	2	4	2	1	3	-	-
White, both sexes	-	-	2	1	2	4	7	2	1	4	-	-
Male	-	-	2	-	2	3	3	-	-	1	-	-
Female	-	-	-	1	-	1	4	2	1	3	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Nebraska												
All races, both sexes	-	-	-	-	1	1	-	3	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	3	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	1	-	-	1	1	1	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	2	1	-	-	1	1	1	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
New Hampshire												
All races, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	1	2	2	2	6	4	5	1	3	-	-
Male	2	1	2	1	-	2	1	2	1	-	-	-
Female	-	-	-	1	2	4	3	3	-	3	-	-
White, both sexes	1	1	2	2	1	5	4	4	1	3	-	-
Male	1	1	2	1	-	2	1	1	1	-	-	-
Female	-	-	-	1	1	3	3	3	-	3	-	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
New Mexico												
All races, both sexes	1	1	2	-	-	-	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	4	2	4	7	7	15	7	5	2	1	-
Male	2	2	1	3	4	3	9	4	-	1	1	-
Female	3	2	1	1	3	4	6	3	5	1	-	-
White, both sexes	4	4	2	3	6	7	14	7	5	2	1	-
Male	2	2	1	2	3	3	9	4	-	1	1	-
Female	2	2	1	1	3	4	5	3	5	1	-	-
Black, both sexes	1	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
North Carolina												
All races, both sexes	1	1	3	-	1	3	7	5	6	-	-	-
Male	-	1	3	-	-	2	3	4	1	-	-	-
Female	1	-	-	-	1	1	4	1	5	-	-	-
White, both sexes	1	1	3	-	1	3	5	4	4	-	-	-
Male	-	1	3	-	-	2	3	3	1	-	-	-
Female	1	-	-	-	1	1	2	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	2	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Ohio												
All races, both sexes	2	4	3	2	6	9	13	10	2	1	-	-
Male	-	2	2	1	4	3	7	3	1	-	-	-
Female	2	2	1	1	2	6	6	7	1	1	-	-
White, both sexes	2	3	1	2	5	7	13	10	2	1	-	-
Male	-	2	1	1	4	2	7	3	1	-	-	-
Female	2	1	-	1	1	5	6	7	1	1	-	-
Black, both sexes	-	1	2	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	3	2	3	2	5	1	1	2	-	-	-
Male	2	1	1	1	2	2	-	1	-	-	-	-
Female	-	2	1	2	-	3	1	-	2	-	-	-
White, both sexes	1	2	2	2	2	5	1	1	2	-	-	-
Male	1	-	1	-	2	2	-	1	-	-	-	-
Female	-	2	1	2	-	3	1	-	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Oregon												
All races, both sexes	1	2	2	1	1	2	2	1	-	-	-	-
Male	1	1	-	-	1	2	1	-	-	-	-	-
Female	-	1	2	1	-	-	1	1	-	-	-	-
White, both sexes	1	2	2	1	1	2	2	1	-	-	-	-
Male	1	1	-	-	1	2	1	-	-	-	-	-
Female	-	1	2	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	3	8	3	4	11	6	8	7	1	-	-
Male	1	1	3	2	2	8	3	4	4	-	-	-
Female	3	2	5	1	2	3	3	4	3	1	-	-
White, both sexes	4	2	8	3	4	9	6	8	7	1	-	-
Male	1	1	3	2	2	6	3	4	4	-	-	-
Female	3	1	5	1	2	3	3	4	3	1	-	-
Black, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Rhode Island												
All races, both sexes	-	-	-	1	1	1	-	-	2	-	-	-
Male	-	-	-	1	1	-	-	-	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	1	-	-	2	-	-	-
Male	-	-	-	1	1	-	-	-	2	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	2	3	1	3	3	9	2	3	-	-	-
Male	1	2	2	-	2	3	4	1	2	-	-	-
Female	2	-	1	1	1	-	5	1	1	-	-	-
White, both sexes	1	-	3	1	1	2	6	2	3	-	-	-
Male	-	-	2	-	1	2	3	1	2	-	-	-
Female	1	-	1	1	-	-	3	1	1	-	-	-
Black, both sexes	2	2	-	-	2	1	3	-	-	-	-	-
Male	1	2	-	-	1	1	1	-	-	-	-	-
Female	1	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
South Dakota												
All races, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	4	1	3	3	5	3	3	-	-	-	-
Male	4	-	1	2	3	-	3	2	-	-	-	-
Female	2	4	-	1	-	5	-	1	-	-	-	-
White, both sexes	5	3	1	2	3	5	3	3	-	-	-	-
Male	3	-	1	1	3	-	3	2	-	-	-	-
Female	2	3	-	1	-	5	-	1	-	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Texas												
All races, both sexes	3	1	8	9	10	11	7	9	4	2	-	-
Male	1	1	5	4	2	6	3	6	2	1	-	-
Female	2	-	3	5	8	5	4	3	2	1	-	-
White, both sexes	3	1	7	8	7	9	5	9	4	2	-	-
Male	1	1	4	4	1	6	3	6	2	1	-	-
Female	2	-	3	4	6	3	2	3	2	1	-	-
Black, both sexes	-	-	1	1	3	2	2	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	2	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Vermont												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	3	1	6	7	5	4	2	-	-	-
Male	1	1	2	1	1	4	2	1	-	-	-	-
Female	1	-	1	-	5	3	3	3	2	-	-	-
White, both sexes	1	1	2	1	5	7	5	4	2	-	-	-
Male	-	1	1	1	-	4	2	1	-	-	-	-
Female	1	-	1	-	5	3	3	3	2	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Washington												
All races, both sexes	3	-	3	2	1	5	6	6	5	1	-	-
Male	1	-	1	-	-	3	4	3	4	-	-	-
Female	2	-	2	2	1	2	2	3	1	1	-	-
White, both sexes	2	-	3	2	1	5	6	4	5	1	-	-
Male	1	-	1	-	-	3	4	2	4	-	-	-
Female	1	-	2	2	1	2	2	2	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	2	2	2	-	-	3	1	-	1	-	-
Male	-	2	1	2	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	1	-	-
White, both sexes	-	2	2	2	-	-	3	1	-	1	-	-
Male	-	2	1	2	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Wisconsin												
All races, both sexes	1	-	3	1	1	4	2	4	2	2	-	-
Male	1	-	2	1	-	2	1	1	1	-	-	-
Female	-	-	1	-	1	2	1	3	1	2	-	-
White, both sexes	1	-	2	1	1	3	2	4	2	2	-	-
Male	1	-	1	1	-	1	1	1	1	-	-	-
Female	-	-	1	-	1	2	1	3	1	2	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
United States												
All races, both sexes	3,307	-	2	-	-	3	4	1	7	15	23	44
Male	968	-	2	-	-	2	2	-	3	11	15	28
Female	2,339	-	-	-	-	1	2	1	4	4	8	16
White, both sexes	2,971	-	1	-	-	3	4	-	4	12	16	34
Male	845	-	1	-	-	2	2	-	2	9	11	23
Female	2,126	-	-	-	-	1	2	-	2	3	5	11
Black, both sexes	287	-	1	-	-	-	-	1	2	3	7	10
Male	109	-	1	-	-	-	-	-	1	2	4	5
Female	178	-	-	-	-	-	-	1	1	1	3	5
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	50	-	-	-	-	1	1	1	-	1	2	-
Male	14	-	-	-	-	1	1	-	-	1	1	-
Female	36	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	38	-	-	-	-	1	1	-	-	1	1	-
Male	10	-	-	-	-	1	1	-	-	1	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	1	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	36	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Arkansas												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	285	-	-	-	-	-	-	-	1	-	3	2
Male	93	-	-	-	-	-	-	-	1	-	2	1
Female	192	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	254	-	-	-	-	-	-	-	1	-	2	2
Male	77	-	-	-	-	-	-	-	1	-	1	1
Female	177	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Colorado												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Florida												
All races, both sexes	155	-	-	-	-	-	1	-	-	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	111	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	141	-	-	-	-	-	1	-	-	-	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	103	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	67	-	-	-	-	-	-	-	2	1	1	4
Male	23	-	-	-	-	-	-	-	1	1	-	3
Female	44	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	1	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	18	-	-	-	-	-	-	-	2	1	-	2
Male	7	-	-	-	-	-	-	-	1	1	-	2
Female	11	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Hawaii												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Illinois												
All races, both sexes	163	-	-	-	-	-	-	-	-	3	1	-
Male	51	-	-	-	-	-	-	-	-	2	-	-
Female	112	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	147	-	-	-	-	-	-	-	-	3	-	-
Male	43	-	-	-	-	-	-	-	-	2	-	-
Female	104	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Iowa												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	40	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	1	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Kentucky												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Maine												
All races, both sexes	35	-	-	-	-	-	1	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	1	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Massachusetts												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	74	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	104	-	-	-	-	-	-	-	-	1	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	96	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Minnesota												
All races, both sexes	92	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	71	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	92	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Missouri												
All races, both sexes	85	-	1	-	-	-	-	-	-	-	-	1
Male	24	-	1	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	-	1	-	-	-	-	-	-	-	-	1
Male	21	-	1	-	-	-	-	-	-	-	-	1
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Nebraska												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
New Hampshire												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
New Mexico												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	128	-	-	-	-	-	-	-	-	1	4	4
Male	45	-	-	-	-	-	-	-	-	-	3	3
Female	83	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	103	-	-	-	-	-	-	-	-	1	2	3
Male	32	-	-	-	-	-	-	-	-	-	1	2
Female	71	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	25	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	2	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
North Carolina												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	78	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Ohio												
All races, both sexes	162	-	-	-	-	-	-	-	1	-	1	-
Male	51	-	-	-	-	-	-	-	-	-	1	-
Female	111	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	145	-	-	-	-	-	-	-	1	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	101	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Oregon												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	2
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	86	-	-	-	-	-	-	-	-	-	1	2
Male	21	-	-	-	-	-	-	-	-	-	1	2
Female	65	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	156	-	-	-	-	1	-	-	-	1	1	2
Male	50	-	-	-	-	-	-	-	-	1	-	1
Female	106	-	-	-	-	1	-	-	-	-	1	1
White, both sexes	145	-	-	-	-	1	-	-	-	1	1	1
Male	44	-	-	-	-	-	-	-	-	1	-	1
Female	101	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
South Dakota												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	94	-	-	-	-	-	1	-	-	-	3	2
Male	25	-	-	-	-	-	1	-	-	-	3	1
Female	69	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	78	-	-	-	-	-	1	-	-	-	2	1
Male	20	-	-	-	-	-	1	-	-	-	2	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Texas												
All races, both sexes	199	-	-	-	-	1	-	-	1	5	1	7
Male	75	-	-	-	-	1	-	-	-	4	1	5
Female	124	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	174	-	-	-	-	1	-	-	1	4	1	5
Male	65	-	-	-	-	1	-	-	-	3	1	4
Female	109	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	23	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Washington												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	72	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	98	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	26	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Wisconsin												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
United States												
All races, both sexes	58	76	99	113	191	310	595	734	643	305	83	1
Male	36	41	52	52	83	117	204	170	108	33	8	1
Female	22	35	47	61	108	193	391	564	535	272	75	-
White, both sexes	47	58	74	93	169	272	533	678	613	285	75	-
Male	31	31	39	41	74	97	183	161	102	29	7	-
Female	16	27	35	52	95	175	350	517	511	256	68	-
Black, both sexes	10	17	23	18	19	33	52	49	24	12	5	1
Male	4	9	13	10	8	18	18	8	4	2	1	1
Female	6	8	10	8	11	15	34	41	20	10	4	-
American Indian 1/, both sexes	-	1	1	1	2	1	4	1	-	-	-	-
Male	-	1	-	1	1	-	2	-	-	-	-	-
Female	-	-	1	-	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	4	6	6	6	8	3	-
Male	1	-	-	-	-	2	1	1	2	2	-	-
Female	-	-	1	1	1	2	5	5	4	6	3	-
Alabama												
All races, both sexes	3	4	2	2	2	2	9	8	11	1	-	-
Male	1	2	-	-	1	1	4	1	-	-	-	-
Female	2	2	2	2	1	1	5	7	11	1	-	-
White, both sexes	3	1	1	2	2	2	6	7	9	1	-	-
Male	1	1	-	-	1	1	2	-	-	-	-	-
Female	2	-	1	2	1	1	4	7	9	1	-	-
Black, both sexes	-	3	1	-	-	-	3	1	2	-	-	-
Male	-	1	-	-	-	-	2	1	-	-	-	-
Female	-	2	1	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Alaska												
All races, both sexes	-	-	-	1	-	-	1	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	3	2	1	-	6	4	8	7	4	-	-
Male	-	2	1	-	-	2	3	3	1	-	-	-
Female	-	1	1	1	-	4	1	5	6	4	-	-
White, both sexes	-	3	2	1	-	6	4	8	7	4	-	-
Male	-	2	1	-	-	2	3	3	1	-	-	-
Female	-	1	1	1	-	4	1	5	6	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Arkansas												
All races, both sexes	-	1	-	3	2	1	5	3	5	4	1	-
Male	-	1	-	2	1	-	3	1	1	-	-	-
Female	-	-	-	1	1	1	2	2	4	4	1	-
White, both sexes	-	1	-	2	1	1	5	2	5	4	1	-
Male	-	1	-	1	1	-	3	1	1	-	-	-
Female	-	-	-	1	-	1	2	1	4	4	1	-
Black, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	9	12	15	17	31	47	62	51	24	5	-
Male	3	4	7	5	10	13	16	15	11	5	-	-
Female	3	5	5	10	7	18	31	47	40	19	5	-
White, both sexes	5	7	8	12	15	28	42	59	49	19	5	-
Male	2	2	5	4	9	12	14	14	9	3	-	-
Female	3	5	3	8	6	16	28	45	40	16	5	-
Black, both sexes	-	2	3	2	2	1	1	1	-	-	-	-
Male	-	2	2	1	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	2	4	2	2	5	-	-
Male	1	-	-	-	-	-	1	1	2	2	-	-
Female	-	-	1	1	-	2	3	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Colorado												
All races, both sexes	-	-	2	7	4	9	12	18	18	9	1	-
Male	-	-	2	4	1	3	3	4	3	1	-	-
Female	-	-	-	3	3	6	9	14	15	8	1	-
White, both sexes	-	-	2	6	4	8	12	18	18	9	1	-
Male	-	-	2	3	1	2	3	4	3	1	-	-
Female	-	-	-	3	3	6	9	14	15	8	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	-	1	6	8	16	8	6	2	-
Male	1	-	1	-	-	1	1	3	-	1	-	-
Female	-	-	-	-	1	5	7	13	8	5	2	-
White, both sexes	1	-	1	-	1	6	6	16	7	6	2	-
Male	1	-	1	-	-	1	-	3	-	1	-	-
Female	-	-	-	-	1	5	6	13	7	5	2	-
Black, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Delaware												
All races, both sexes	-	-	-	-	-	1	1	-	1	2	1	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	1	1	1	-
White, both sexes	-	-	-	-	-	1	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Florida												
All races, both sexes	2	8	7	5	7	18	27	30	25	19	5	-
Male	2	4	3	2	4	7	6	8	6	1	-	-
Female	-	4	4	3	3	11	21	22	19	18	5	-
White, both sexes	2	7	6	4	6	16	26	28	22	17	5	-
Male	2	3	2	1	4	6	6	6	6	1	-	-
Female	-	4	4	3	2	10	20	22	16	16	5	-
Black, both sexes	-	1	1	1	1	2	1	2	3	2	-	-
Male	-	1	1	1	-	1	-	2	-	-	-	-
Female	-	-	-	-	1	1	1	-	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	1	2	3	2	7	12	14	8	4	2	-
Male	2	1	1	2	1	3	4	2	2	-	-	-
Female	2	-	1	1	1	4	8	12	6	4	2	-
White, both sexes	3	1	2	2	1	6	10	13	5	3	-	-
Male	2	1	1	2	-	3	3	2	1	-	-	-
Female	1	-	1	-	1	3	7	11	4	3	-	-
Black, both sexes	1	-	-	1	1	1	2	1	3	1	2	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	-	1	-	1	1	1	2	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Hawaii												
All races, both sexes	-	-	-	-	1	1	1	3	3	3	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	3	3	3	2	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	3	3	3	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	3	3	3	2	-
Idaho												
All races, both sexes	-	-	-	-	1	5	4	12	5	4	-	-
Male	-	-	-	-	-	3	-	2	2	1	-	-
Female	-	-	-	-	1	2	4	10	3	3	-	-
White, both sexes	-	-	-	-	1	5	4	12	5	4	-	-
Male	-	-	-	-	-	3	-	2	2	1	-	-
Female	-	-	-	-	1	2	4	10	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Illinois												
All races, both sexes	3	3	4	4	10	17	34	35	30	14	5	-
Male	2	2	3	2	5	6	17	6	4	1	1	-
Female	1	1	1	2	5	11	17	29	26	13	4	-
White, both sexes	2	3	4	3	8	13	30	32	30	14	5	-
Male	1	2	3	1	4	3	16	5	4	1	1	-
Female	1	1	1	2	4	10	14	27	26	13	4	-
Black, both sexes	1	-	-	1	2	4	4	3	-	-	-	-
Male	1	-	-	1	1	3	1	1	-	-	-	-
Female	-	-	-	-	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	3	4	2	5	7	13	17	11	5	-	-
Male	3	1	-	1	2	3	5	4	-	1	-	-
Female	-	2	4	1	3	4	8	13	11	4	-	-
White, both sexes	1	3	3	1	5	7	12	16	9	5	-	-
Male	1	1	-	1	2	3	4	4	-	1	-	-
Female	-	2	3	-	3	4	8	12	9	4	-	-
Black, both sexes	2	-	1	1	-	-	1	1	2	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Iowa												
All races, both sexes	-	-	-	-	2	3	9	16	16	3	-	-
Male	-	-	-	-	2	-	8	3	5	-	-	-
Female	-	-	-	-	-	3	1	13	11	3	-	-
White, both sexes	-	-	-	-	2	3	9	16	16	3	-	-
Male	-	-	-	-	2	-	8	3	5	-	-	-
Female	-	-	-	-	-	3	1	13	11	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	1	-	-	3	11	9	10	3	2	-
Male	-	-	-	-	-	1	2	1	2	-	-	-
Female	-	-	1	-	-	2	9	8	8	3	2	-
White, both sexes	-	-	1	-	-	3	10	9	10	3	2	-
Male	-	-	-	-	-	1	2	1	2	-	-	-
Female	-	-	1	-	-	2	8	8	8	3	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Kentucky												
All races, both sexes	2	1	2	3	-	2	8	10	8	1	2	-
Male	1	1	2	2	-	1	1	2	2	-	-	-
Female	1	-	-	1	-	1	7	8	6	1	2	-
White, both sexes	2	1	2	2	-	2	8	10	8	1	2	-
Male	1	1	2	1	-	1	1	2	2	-	-	-
Female	1	-	-	1	-	1	7	8	6	1	2	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	2	1	5	6	10	8	6	3	2	-
Male	-	-	-	1	2	1	1	2	-	1	-	-
Female	-	-	2	-	3	5	9	6	6	2	2	-
White, both sexes	-	-	1	1	3	5	7	6	5	2	2	-
Male	-	-	-	1	2	1	1	2	-	1	-	-
Female	-	-	1	-	1	4	6	4	5	1	2	-
Black, both sexes	-	-	1	-	2	1	3	2	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	1	3	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Maine												
All races, both sexes	2	-	-	1	2	5	8	7	4	3	1	-
Male	2	-	-	-	2	1	5	3	-	1	-	-
Female	-	-	-	1	-	4	3	4	4	2	1	-
White, both sexes	2	-	-	1	2	5	8	7	4	3	1	-
Male	2	-	-	-	2	1	5	3	-	1	-	-
Female	-	-	-	1	-	4	3	4	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	3	3	2	3	6	16	20	8	7	2	-
Male	-	-	-	1	-	2	1	3	1	1	1	-
Female	-	3	3	1	3	4	15	17	7	6	1	-
White, both sexes	-	-	2	1	3	5	13	17	8	7	1	-
Male	-	-	-	1	-	1	1	3	1	1	-	-
Female	-	-	2	-	3	4	12	14	7	6	1	-
Black, both sexes	-	3	1	1	-	1	2	3	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	3	1	1	-	-	2	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Massachusetts												
All races, both sexes	-	2	2	3	4	12	14	12	16	9	2	-
Male	-	-	2	1	2	5	8	4	4	-	-	-
Female	-	2	-	2	2	7	6	8	12	9	2	-
White, both sexes	-	1	2	3	3	12	13	12	16	9	2	-
Male	-	-	2	1	2	5	8	4	4	-	-	-
Female	-	1	-	2	1	7	5	8	12	9	2	-
Black, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	2	5	7	5	23	23	20	11	4	-
Male	-	1	-	1	2	2	11	2	8	1	2	-
Female	-	-	2	4	5	3	12	21	12	10	2	-
White, both sexes	-	1	1	5	7	3	22	22	19	11	4	-
Male	-	1	-	1	2	2	10	2	7	1	2	-
Female	-	-	1	4	5	1	12	20	12	10	2	-
Black, both sexes	-	-	1	-	-	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Minnesota												
All races, both sexes	1	2	-	-	2	9	13	23	29	8	5	-
Male	1	2	-	-	1	3	4	6	4	-	-	-
Female	-	-	-	-	1	6	9	17	25	8	5	-
White, both sexes	1	2	-	-	2	9	13	23	29	8	5	-
Male	1	2	-	-	1	3	4	6	4	-	-	-
Female	-	-	-	-	1	6	9	17	25	8	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	2	1	4	4	2	3	-	-
Male	-	-	1	1	1	-	1	1	-	1	-	-
Female	-	-	-	-	1	1	3	3	2	2	-	-
White, both sexes	-	-	-	1	2	-	2	4	2	1	-	-
Male	-	-	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	3	2	1	-	-
Black, both sexes	-	-	1	-	-	1	2	-	-	2	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Missouri												
All races, both sexes	1	-	2	3	3	10	19	17	18	8	2	-
Male	1	-	2	2	2	3	6	2	1	1	2	-
Female	-	-	-	1	1	7	13	15	17	7	-	-
White, both sexes	1	-	1	1	3	9	18	15	18	8	2	-
Male	1	-	1	1	2	2	6	2	1	1	2	-
Female	-	-	-	-	1	7	12	13	17	7	-	-
Black, both sexes	-	-	1	2	-	1	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	3	4	10	15	2	1	-
Male	-	-	-	-	-	-	1	1	3	1	-	-
Female	-	-	-	-	-	3	3	9	12	1	1	-
White, both sexes	-	-	-	-	-	3	4	10	15	2	1	-
Male	-	-	-	-	-	-	1	1	3	1	-	-
Female	-	-	-	-	-	3	3	9	12	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Nebraska												
All races, both sexes	-	-	-	2	2	3	4	12	13	6	2	-
Male	-	-	-	1	-	-	-	4	3	-	-	-
Female	-	-	-	1	2	3	4	8	10	6	2	-
White, both sexes	-	-	-	2	2	3	4	12	13	6	2	-
Male	-	-	-	1	-	-	-	4	3	-	-	-
Female	-	-	-	1	2	3	4	8	10	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	4	3	3	4	3	-	-	-
Male	-	-	-	1	3	1	2	1	-	-	-	-
Female	-	-	-	-	1	2	1	3	3	-	-	-
White, both sexes	-	-	-	1	3	2	3	4	3	-	-	-
Male	-	-	-	1	2	-	2	1	-	-	-	-
Female	-	-	-	-	1	2	1	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
New Hampshire												
All races, both sexes	1	-	-	2	-	2	3	2	4	1	-	-
Male	1	-	-	1	-	2	2	2	-	-	-	-
Female	-	-	-	1	-	-	1	-	4	1	-	-
White, both sexes	1	-	-	2	-	2	3	2	4	1	-	-
Male	1	-	-	1	-	2	2	2	-	-	-	-
Female	-	-	-	1	-	-	1	-	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	1	4	6	8	10	17	20	13	7	-	-
Male	2	-	2	1	2	4	6	7	2	1	-	-
Female	-	1	2	5	6	6	11	13	11	6	-	-
White, both sexes	2	1	3	4	8	10	14	19	11	7	-	-
Male	2	-	1	-	2	4	6	6	2	1	-	-
Female	-	1	2	4	6	6	8	13	9	6	-	-
Black, both sexes	-	-	1	2	-	-	3	1	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
New Mexico												
All races, both sexes	1	-	-	1	1	-	3	5	4	2	-	-
Male	1	-	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	1	1	-	1	3	3	2	-	-
White, both sexes	1	-	-	1	1	-	3	5	4	2	-	-
Male	1	-	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	1	1	-	1	3	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	6	4	6	9	12	20	17	24	14	1	-
Male	2	3	2	3	5	9	9	1	5	-	-	-
Female	4	3	2	3	4	3	11	16	19	14	1	-
White, both sexes	4	4	4	6	5	10	15	14	22	12	1	-
Male	2	1	2	3	2	8	6	1	4	-	-	-
Female	2	3	2	3	3	2	9	13	18	12	1	-
Black, both sexes	2	2	-	-	4	2	5	3	2	2	-	-
Male	-	2	-	-	3	1	3	-	1	-	-	-
Female	2	-	-	-	1	1	2	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
North Carolina												
All races, both sexes	4	-	3	4	9	4	23	26	21	8	3	-
Male	2	-	2	1	4	-	8	5	3	1	1	-
Female	2	-	1	3	5	4	15	21	18	7	2	-
White, both sexes	3	-	1	1	5	2	19	20	19	8	3	-
Male	2	-	1	-	3	-	6	4	3	1	1	-
Female	1	-	-	1	2	2	13	16	16	7	2	-
Black, both sexes	1	-	2	3	3	2	3	6	2	-	-	-
Male	-	-	1	1	1	-	1	1	-	-	-	-
Female	1	-	1	2	2	2	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	6	2	4	3	-	-
Male	-	-	-	-	-	-	2	1	1	1	-	-
Female	-	-	-	-	-	1	4	1	3	2	-	-
White, both sexes	-	-	-	-	-	1	5	2	4	3	-	-
Male	-	-	-	-	-	-	2	1	1	1	-	-
Female	-	-	-	-	-	1	3	1	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Ohio												
All races, both sexes	2	6	1	6	11	16	28	39	31	16	4	-
Male	-	4	1	4	8	10	6	11	4	2	-	-
Female	2	2	-	2	3	6	22	28	27	14	4	-
White, both sexes	2	5	1	6	9	12	25	34	29	16	4	-
Male	-	3	1	4	7	7	5	11	3	2	-	-
Female	2	2	-	2	2	5	20	23	26	14	4	-
Black, both sexes	-	1	-	-	2	4	3	5	2	-	-	-
Male	-	1	-	-	1	3	1	-	1	-	-	-
Female	-	-	-	-	1	1	2	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	1	2	3	5	7	7	5	3	1	-
Male	-	1	-	1	1	2	2	1	-	-	-	-
Female	-	-	1	1	2	3	5	6	5	3	1	-
White, both sexes	-	1	1	1	3	4	7	7	4	3	-	-
Male	-	1	-	-	1	2	2	1	-	-	-	-
Female	-	-	1	1	2	2	5	6	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Oregon												
All races, both sexes	-	2	2	1	4	10	11	22	23	8	2	1
Male	-	2	1	-	3	2	2	6	3	-	-	1
Female	-	-	1	1	1	8	9	16	20	8	2	-
White, both sexes	-	1	1	1	4	10	11	22	23	8	2	-
Male	-	1	1	-	3	2	2	6	3	-	-	-
Female	-	-	-	1	1	8	9	16	20	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	2	3	3	10	18	31	37	30	12	4	-
Male	1	1	3	3	4	5	12	13	6	-	-	-
Female	-	1	-	-	6	13	19	24	24	12	4	-
White, both sexes	1	-	3	2	10	17	30	34	29	11	4	-
Male	1	-	3	2	4	4	11	11	6	-	-	-
Female	-	-	-	-	6	13	19	23	23	11	4	-
Black, both sexes	-	2	-	1	-	1	1	3	1	1	-	-
Male	-	1	-	1	-	1	1	2	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Rhode Island												
All races, both sexes	-	-	-	1	1	1	5	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	4	-	1	-	1	-
White, both sexes	-	-	-	1	1	1	5	-	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	4	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	1	-	4	3	12	10	9	2	2	-
Male	1	1	-	-	1	1	2	1	1	-	-	-
Female	-	1	1	-	3	2	10	9	8	2	2	-
White, both sexes	1	1	-	-	4	2	11	8	9	2	2	-
Male	1	-	-	-	1	1	2	1	1	-	-	-
Female	-	1	-	-	3	1	9	7	8	2	2	-
Black, both sexes	-	1	1	-	-	1	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
South Dakota												
All races, both sexes	1	-	-	-	1	2	1	3	10	6	1	-
Male	-	-	-	-	-	1	-	2	4	-	-	-
Female	1	-	-	-	1	1	1	1	6	6	1	-
White, both sexes	1	-	-	-	1	2	1	3	10	6	1	-
Male	-	-	-	-	-	1	-	2	4	-	-	-
Female	1	-	-	-	1	1	1	1	6	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	5	3	6	9	10	14	19	10	8	1	-
Male	2	2	1	4	1	3	3	2	1	1	-	-
Female	1	3	2	2	8	7	11	17	9	7	1	-
White, both sexes	1	4	3	6	8	6	11	17	10	7	1	-
Male	1	2	1	4	1	1	2	2	1	1	-	-
Female	-	2	2	2	7	5	9	15	9	6	1	-
Black, both sexes	2	1	-	-	-	4	3	2	-	1	-	-
Male	1	-	-	-	-	2	1	-	-	-	-	-
Female	1	1	-	-	-	2	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Texas												
All races, both sexes	4	4	15	6	14	14	35	39	32	17	4	-
Male	3	4	10	3	7	5	14	10	6	2	-	-
Female	1	-	5	3	7	9	21	29	26	15	4	-
White, both sexes	3	4	9	6	14	12	30	32	32	16	4	-
Male	3	4	6	3	7	4	11	10	6	2	-	-
Female	-	-	3	3	7	8	19	22	26	14	4	-
Black, both sexes	1	-	6	-	-	2	4	6	-	1	-	-
Male	-	-	4	-	-	1	2	-	-	-	-	-
Female	1	-	2	-	-	1	2	6	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	1	-	4	2	12	17	7	3	-	-
Male	1	-	1	-	-	2	5	6	1	-	-	-
Female	1	-	-	-	4	-	7	11	6	3	-	-
White, both sexes	2	-	1	-	4	2	12	17	7	3	-	-
Male	1	-	1	-	-	2	5	6	1	-	-	-
Female	1	-	-	-	4	-	7	11	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	2	6	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	6	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	6	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	2	1	3	3	8	16	15	7	2	-
Male	1	-	-	-	-	3	2	3	1	-	-	-
Female	-	-	2	1	3	-	6	13	14	7	2	-
White, both sexes	1	-	2	1	3	1	6	13	13	7	2	-
Male	1	-	-	-	-	1	2	3	1	-	-	-
Female	-	-	2	1	3	-	4	10	12	7	2	-
Black, both sexes	-	-	-	-	-	2	2	3	2	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Washington												
All races, both sexes	-	6	5	1	8	9	13	23	18	10	6	-
Male	-	2	2	-	3	3	5	7	3	1	1	-
Female	-	4	3	1	5	6	8	16	15	9	5	-
White, both sexes	-	6	4	1	8	9	13	23	18	10	5	-
Male	-	2	1	-	3	3	5	7	3	1	1	-
Female	-	4	3	1	5	6	8	16	15	9	4	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	-	-	1	-	1	2	7	8	4	-	1	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	7	8	4	-	1	-
White, both sexes	-	-	1	-	1	2	7	8	4	-	1	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	7	8	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Wisconsin												
All races, both sexes	1	-	1	2	3	3	13	14	20	9	1	-
Male	-	-	-	1	2	-	7	4	2	4	-	-
Female	1	-	1	1	1	3	6	10	18	5	1	-
White, both sexes	1	-	1	2	3	3	12	13	20	9	1	-
Male	-	-	-	1	2	-	7	4	2	4	-	-
Female	1	-	1	1	1	3	5	9	18	5	1	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	4	1	2	-	-
Male	-	-	-	-	-	-	-	2	1	1	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	4	1	2	-	-
Male	-	-	-	-	-	-	-	2	1	1	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
United States												
All races, both sexes	1,695	-	-	-	-	3	-	-	1	2	-	5
Male	216	-	-	-	-	2	-	-	1	1	-	3
Female	1,479	-	-	-	-	1	-	-	-	1	-	2
White, both sexes	1,640	-	-	-	-	3	-	-	1	2	-	5
Male	211	-	-	-	-	2	-	-	1	1	-	3
Female	1,429	-	-	-	-	1	-	-	-	1	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	24	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Arkansas												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	150	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	127	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	140	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	120	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Colorado												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Florida												
All races, both sexes	75	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	75	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Hawaii												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Illinois												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Iowa												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Kentucky												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Maine												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Massachusetts												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Minnesota												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Missouri												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Nebraska												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
North Carolina												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Ohio												
All races, both sexes	90	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	88	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Oregon												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	69	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	67	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
South Dakota												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Texas												
All races, both sexes	92	-	-	-	-	1	-	-	-	-	-	1
Male	14	-	-	-	-	1	-	-	-	-	-	1
Female	78	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	90	-	-	-	-	1	-	-	-	-	-	1
Male	14	-	-	-	-	1	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Vermont												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Washington												
All races, both sexes	69	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
Wisconsin												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
United States												
All races, both sexes	5	10	11	20	44	117	290	456	430	240	61	-
Male	-	5	3	2	9	16	51	60	41	18	4	-
Female	5	5	8	18	35	101	239	396	389	222	57	-
White, both sexes	5	9	10	19	43	113	276	446	420	230	58	-
Male	-	5	3	2	9	15	50	60	40	16	4	-
Female	5	4	7	17	34	98	226	386	380	214	54	-
Black, both sexes	-	1	-	1	-	1	8	4	5	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	7	4	5	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	4	5	5	8	3	-
Male	-	-	-	-	-	1	-	-	1	2	-	-
Female	-	-	-	-	1	1	4	5	4	6	3	-
Alabama												
All races, both sexes	1	-	1	1	-	1	5	6	7	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	1	4	6	7	1	-	-
White, both sexes	1	-	1	1	-	1	5	6	7	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	1	4	6	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	2	-	3	5	4	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	2	-	2	5	4	-	-
White, both sexes	-	-	-	1	-	2	-	3	5	4	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	2	-	2	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Arkansas												
All races, both sexes	-	-	-	-	-	1	3	1	3	3	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	3	3	1	-
White, both sexes	-	-	-	-	-	1	3	1	3	3	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	3	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	3	2	4	12	29	37	37	21	5	-
Male	-	-	2	1	1	2	6	4	3	4	-	-
Female	-	-	1	1	3	10	23	33	34	17	5	-
White, both sexes	-	-	3	2	4	11	27	36	36	16	5	-
Male	-	-	2	1	1	2	6	4	2	2	-	-
Female	-	-	1	1	3	9	21	32	34	14	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	1	1	5	-	-
Male	-	-	-	-	-	-	-	-	1	2	-	-
Female	-	-	-	-	-	1	2	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Colorado												
All races, both sexes	-	-	-	1	1	4	8	14	10	8	1	-
Male	-	-	-	-	-	-	1	2	1	1	-	-
Female	-	-	-	1	1	4	7	12	9	7	1	-
White, both sexes	-	-	-	1	1	4	8	14	10	8	1	-
Male	-	-	-	-	-	-	1	2	1	1	-	-
Female	-	-	-	1	1	4	7	12	9	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	1	3	3	11	4	3	2	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	3	3	9	4	3	2	-
White, both sexes	-	-	-	-	1	3	3	11	4	3	2	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	3	3	9	4	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Florida												
All races, both sexes	-	3	2	-	2	5	11	20	16	14	2	-
Male	-	2	-	-	-	1	1	2	4	1	-	-
Female	-	1	2	-	2	4	10	18	12	13	2	-
White, both sexes	-	3	2	-	2	5	11	20	16	14	2	-
Male	-	2	-	-	-	1	1	2	4	1	-	-
Female	-	1	2	-	2	4	10	18	12	13	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	4	2	9	5	3	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	1	-	-	-	-	3	1	8	4	3	-	-
White, both sexes	1	-	-	-	-	4	2	9	4	3	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	1	-	-	-	-	3	1	8	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Hawaii												
All races, both sexes	-	-	-	-	1	1	1	3	3	3	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	3	3	3	2	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	3	3	3	2	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	3	3	3	2	-
Idaho												
All races, both sexes	-	-	-	-	1	3	4	10	4	4	-	-
Male	-	-	-	-	-	1	-	1	2	1	-	-
Female	-	-	-	-	1	2	4	9	2	3	-	-
White, both sexes	-	-	-	-	1	3	4	10	4	4	-	-
Male	-	-	-	-	-	1	-	1	2	1	-	-
Female	-	-	-	-	1	2	4	9	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Illinois												
All races, both sexes	-	-	-	1	3	7	16	18	22	13	5	-
Male	-	-	-	-	2	2	5	2	2	1	1	-
Female	-	-	-	1	1	5	11	16	20	12	4	-
White, both sexes	-	-	-	1	3	7	14	18	22	13	5	-
Male	-	-	-	-	2	2	4	2	2	1	1	-
Female	-	-	-	1	1	5	10	16	20	12	4	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	1	-	-	1	2	8	12	7	5	-	-
Male	-	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	1	2	7	11	7	4	-	-
White, both sexes	-	1	-	-	1	2	8	12	7	5	-	-
Male	-	1	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	-	1	2	7	11	7	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Iowa												
All races, both sexes	-	-	-	-	-	2	2	8	12	3	-	-
Male	-	-	-	-	-	-	1	-	4	-	-	-
Female	-	-	-	-	-	2	1	8	8	3	-	-
White, both sexes	-	-	-	-	-	2	2	8	12	3	-	-
Male	-	-	-	-	-	-	1	-	4	-	-	-
Female	-	-	-	-	-	2	1	8	8	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	2	5	8	6	2	2	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	2	5	7	5	2	2	-
White, both sexes	-	-	-	-	-	2	5	8	6	2	2	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	2	5	7	5	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Kentucky												
All races, both sexes	1	-	-	1	-	1	5	7	6	1	2	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	1	4	7	6	1	2	-
White, both sexes	1	-	-	1	-	1	5	7	6	1	2	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	1	4	7	6	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	2	6	6	4	-	1	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	2	5	5	4	-	1	-
White, both sexes	-	-	-	-	-	2	5	5	4	-	1	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	2	4	4	4	-	1	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Maine												
All races, both sexes	-	-	-	1	-	2	4	5	2	2	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	2	3	4	2	2	-	-
White, both sexes	-	-	-	1	-	2	4	5	2	2	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	2	3	4	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	1	1	1	9	12	4	5	1	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	1	1	1	9	10	4	5	1	-
White, both sexes	-	-	1	-	1	1	7	12	4	5	1	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	-	1	1	7	10	4	5	1	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Massachusetts												
All races, both sexes	-	-	-	-	-	4	5	5	8	7	2	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	4	3	4	8	7	2	-
White, both sexes	-	-	-	-	-	4	5	5	8	7	2	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	4	3	4	8	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	3	2	-	8	17	9	9	1	-
Male	-	-	-	-	-	-	2	2	3	1	1	-
Female	-	-	-	3	2	-	6	15	6	8	-	-
White, both sexes	-	-	-	3	2	-	8	17	9	9	1	-
Male	-	-	-	-	-	-	2	2	3	1	1	-
Female	-	-	-	3	2	-	6	15	6	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Minnesota												
All races, both sexes	-	1	-	-	-	2	7	16	18	7	3	-
Male	-	1	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	-	-	2	5	14	17	7	3	-
White, both sexes	-	1	-	-	-	2	7	16	18	7	3	-
Male	-	1	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	-	-	2	5	14	17	7	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	2	-	1	2	2	2	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	2	2	-	-
White, both sexes	-	-	-	-	2	-	1	2	2	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Missouri												
All races, both sexes	-	-	-	-	1	4	9	13	13	6	1	-
Male	-	-	-	-	-	-	1	2	-	-	1	-
Female	-	-	-	-	1	4	8	11	13	6	-	-
White, both sexes	-	-	-	-	1	4	8	12	13	6	1	-
Male	-	-	-	-	-	-	1	2	-	-	1	-
Female	-	-	-	-	1	4	7	10	13	6	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	2	4	9	13	2	1	-
Male	-	-	-	-	-	-	1	1	3	1	-	-
Female	-	-	-	-	-	2	3	8	10	1	1	-
White, both sexes	-	-	-	-	-	2	4	9	13	2	1	-
Male	-	-	-	-	-	-	1	1	3	1	-	-
Female	-	-	-	-	-	2	3	8	10	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Nebraska												
All races, both sexes	-	-	-	-	-	3	3	9	10	5	2	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	-	3	3	8	8	5	2	-
White, both sexes	-	-	-	-	-	3	3	9	10	5	2	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	-	3	3	8	8	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	4	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	1	-	2	3	2	3	9	7	7	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	2	3	2	3	8	7	6	-	-
White, both sexes	-	1	-	2	3	2	3	9	6	7	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	1	-	2	3	2	3	8	6	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	2	2	2	2	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	2	-	-
White, both sexes	-	-	-	-	-	-	2	2	2	2	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	-	2	2	6	9	5	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	-	-	1	1	6	8	5	-	-
White, both sexes	-	-	1	-	-	2	1	6	9	5	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	-	-	1	-	6	8	5	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
North Carolina												
All races, both sexes	-	-	-	1	2	-	13	13	14	6	2	-
Male	-	-	-	-	1	-	1	2	-	1	1	-
Female	-	-	-	1	1	-	12	11	14	5	1	-
White, both sexes	-	-	-	1	2	-	12	12	13	6	2	-
Male	-	-	-	-	1	-	1	2	-	1	1	-
Female	-	-	-	1	1	-	11	10	13	5	1	-
Black, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	5	2	2	2	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	3	1	2	2	-	-
White, both sexes	-	-	-	-	-	1	4	2	2	2	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	2	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Ohio												
All races, both sexes	-	-	-	-	2	6	16	25	21	16	4	-
Male	-	-	-	-	1	2	1	4	1	2	-	-
Female	-	-	-	-	1	4	15	21	20	14	4	-
White, both sexes	-	-	-	-	2	6	16	24	20	16	4	-
Male	-	-	-	-	1	2	1	4	1	2	-	-
Female	-	-	-	-	1	4	15	20	19	14	4	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	2	3	4	5	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	2	4	5	2	-	-
White, both sexes	-	-	-	-	-	1	3	4	4	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	4	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Oregon												
All races, both sexes	-	-	1	-	1	6	8	18	16	7	2	-
Male	-	-	-	-	1	-	-	3	-	-	-	-
Female	-	-	1	-	-	6	8	15	16	7	2	-
White, both sexes	-	-	-	-	1	6	8	18	16	7	2	-
Male	-	-	-	-	1	-	-	3	-	-	-	-
Female	-	-	-	-	-	6	8	15	16	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	1	7	15	15	18	7	4	-
Male	-	-	-	-	-	-	3	2	3	-	-	-
Female	-	1	-	-	1	7	12	13	15	7	4	-
White, both sexes	-	-	-	-	1	7	15	15	17	7	4	-
Male	-	-	-	-	-	-	3	2	3	-	-	-
Female	-	-	-	-	1	7	12	13	14	7	4	-
Black, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	2	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	1	-
White, both sexes	-	-	-	-	-	1	2	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	1	1	8	7	7	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	8	7	7	2	1	-
White, both sexes	-	-	-	-	1	1	8	7	7	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	8	7	7	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
South Dakota												
All races, both sexes	1	-	-	-	-	2	1	1	8	6	1	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	1	-	-	-	-	1	1	-	6	6	1	-
White, both sexes	1	-	-	-	-	2	1	1	8	6	1	-
Male	-	-	-	-	-	1	-	1	2	-	-	-
Female	1	-	-	-	-	1	1	-	6	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	1	6	2	5	10	6	5	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	5	2	5	10	6	5	1	-
White, both sexes	-	-	-	1	5	1	5	10	6	4	1	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	4	1	5	10	6	4	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Texas												
All races, both sexes	-	1	1	2	-	7	15	24	23	14	3	-
Male	-	1	1	1	-	1	2	3	2	1	-	-
Female	-	-	-	1	-	6	13	21	21	13	3	-
White, both sexes	-	1	1	2	-	7	14	23	23	14	3	-
Male	-	1	1	1	-	1	2	3	2	1	-	-
Female	-	-	-	1	-	6	12	20	21	13	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	2	1	11	15	5	3	-	-
Male	-	-	-	-	-	1	4	4	-	-	-	-
Female	1	-	-	-	2	-	7	11	5	3	-	-
White, both sexes	1	-	-	-	2	1	11	15	5	3	-	-
Male	-	-	-	-	-	1	4	4	-	-	-	-
Female	1	-	-	-	2	-	7	11	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	1	5	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	5	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	5	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	1	1	3	9	13	6	2	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	1	-	3	8	12	6	2	-
White, both sexes	-	-	-	-	1	1	3	8	13	6	2	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	1	-	3	7	12	6	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Washington												
All races, both sexes	-	2	1	1	3	5	9	20	14	10	4	-
Male	-	-	-	-	-	1	3	5	1	1	-	-
Female	-	2	1	1	3	4	6	15	13	9	4	-
White, both sexes	-	2	1	1	3	5	9	20	14	10	3	-
Male	-	-	-	-	-	1	3	5	1	1	-	-
Female	-	2	1	1	3	4	6	15	13	9	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	3	2	4	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	2	4	-	1	-
White, both sexes	-	-	-	-	-	-	3	2	4	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	2	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Wisconsin												
All races, both sexes	-	-	-	-	2	1	5	8	14	5	1	-
Male	-	-	-	-	1	-	2	2	1	1	-	-
Female	-	-	-	-	1	1	3	6	13	4	1	-
White, both sexes	-	-	-	-	2	1	5	8	14	5	1	-
Male	-	-	-	-	1	-	2	2	1	1	-	-
Female	-	-	-	-	1	1	3	6	13	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
United States												
All races, both sexes	1,612	-	2	-	-	-	4	1	6	13	23	39
Male	752	-	2	-	-	-	2	-	2	10	15	25
Female	860	-	-	-	-	-	2	1	4	3	8	14
White, both sexes	1,331	-	1	-	-	-	4	-	3	10	16	29
Male	634	-	1	-	-	-	2	-	1	8	11	20
Female	697	-	-	-	-	-	2	-	2	2	5	9
Black, both sexes	265	-	1	-	-	-	-	1	2	3	7	10
Male	108	-	1	-	-	-	-	-	1	2	4	5
Female	157	-	-	-	-	-	-	1	1	1	3	5
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	26	-	-	-	-	-	1	1	-	1	2	-
Male	12	-	-	-	-	-	1	-	-	1	1	-
Female	14	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	14	-	-	-	-	-	1	-	-	1	1	-
Male	8	-	-	-	-	-	1	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	1	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Arkansas												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	135	-	-	-	-	-	-	-	1	-	3	2
Male	70	-	-	-	-	-	-	-	1	-	2	1
Female	65	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	114	-	-	-	-	-	-	-	1	-	2	2
Male	57	-	-	-	-	-	-	-	1	-	1	1
Female	57	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Colorado												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Florida												
All races, both sexes	80	-	-	-	-	-	1	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	66	-	-	-	-	-	1	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	42	-	-	-	-	-	-	-	2	1	1	3
Male	19	-	-	-	-	-	-	-	1	1	-	3
Female	23	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	17	-	-	-	-	-	-	-	2	1	-	2
Male	7	-	-	-	-	-	-	-	1	1	-	2
Female	10	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Illinois												
All races, both sexes	78	-	-	-	-	-	-	-	-	3	1	-
Male	36	-	-	-	-	-	-	-	-	2	-	-
Female	42	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	64	-	-	-	-	-	-	-	-	3	-	-
Male	29	-	-	-	-	-	-	-	-	2	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	15	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	1	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Kentucky												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Maine												
All races, both sexes	19	-	-	-	-	-	1	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	1	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Massachusetts												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	55	-	-	-	-	-	-	-	-	1	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Minnesota												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Missouri												
All races, both sexes	37	-	1	-	-	-	-	-	-	-	-	-
Male	19	-	1	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	1	-	-	-	-	-	-	-	-	-
Male	16	-	1	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Nebraska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
New Hampshire												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
New Mexico												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	102	-	-	-	-	-	-	-	-	-	4	4
Male	42	-	-	-	-	-	-	-	-	-	3	3
Female	60	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	78	-	-	-	-	-	-	-	-	-	2	3
Male	29	-	-	-	-	-	-	-	-	-	1	2
Female	49	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	2	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
North Carolina												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Ohio												
All races, both sexes	72	-	-	-	-	-	-	-	1	-	1	-
Male	40	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	1	-	1	-
Male	33	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Oregon												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	2
Male	19	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	2
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	87	-	-	-	-	-	-	-	-	1	1	2
Male	42	-	-	-	-	-	-	-	-	1	-	1
Female	45	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	78	-	-	-	-	-	-	-	-	1	1	1
Male	36	-	-	-	-	-	-	-	-	1	-	1
Female	42	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	58	-	-	-	-	-	1	-	-	-	3	2
Male	24	-	-	-	-	-	1	-	-	-	3	1
Female	34	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	1	-	-	-	2	1
Male	19	-	-	-	-	-	1	-	-	-	2	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Texas												
All races, both sexes	107	-	-	-	-	-	-	-	1	5	1	6
Male	61	-	-	-	-	-	-	-	-	4	1	4
Female	46	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	84	-	-	-	-	-	-	-	1	4	1	4
Male	51	-	-	-	-	-	-	-	-	3	1	3
Female	33	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Washington												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Wisconsin												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
United States												
All races, both sexes	53	66	88	93	147	193	305	278	213	65	22	1
Male	36	36	49	50	74	101	153	110	67	15	4	1
Female	17	30	39	43	73	92	152	168	146	50	18	-
White, both sexes	42	49	64	74	126	159	257	232	193	55	17	-
Male	31	26	36	39	65	82	133	101	62	13	3	-
Female	11	23	28	35	61	77	124	131	131	42	14	-
Black, both sexes	10	16	23	17	19	32	44	45	19	10	5	1
Male	4	9	13	10	8	18	17	8	4	2	1	1
Female	6	7	10	7	11	14	27	37	15	8	4	-
American Indian 1/, both sexes	-	1	-	1	2	-	2	-	-	-	-	-
Male	-	1	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	2	2	1	1	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-
Alabama												
All races, both sexes	2	4	1	1	2	1	4	2	4	-	-	-
Male	1	2	-	-	1	1	3	1	-	-	-	-
Female	1	2	1	1	1	-	1	1	4	-	-	-
White, both sexes	2	1	-	1	2	1	1	1	2	-	-	-
Male	1	1	-	-	1	1	1	-	-	-	-	-
Female	1	-	-	1	1	-	-	1	2	-	-	-
Black, both sexes	-	3	1	-	-	-	3	1	2	-	-	-
Male	-	1	-	-	-	-	2	1	-	-	-	-
Female	-	2	1	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Alaska												
All races, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	3	2	-	-	4	4	5	2	-	-	-
Male	-	2	1	-	-	2	3	2	1	-	-	-
Female	-	1	1	-	-	2	1	3	1	-	-	-
White, both sexes	-	3	2	-	-	4	4	5	2	-	-	-
Male	-	2	1	-	-	2	3	2	1	-	-	-
Female	-	1	1	-	-	2	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Arkansas												
All races, both sexes	-	1	-	3	2	-	2	2	2	1	-	-
Male	-	1	-	2	1	-	2	1	1	-	-	-
Female	-	-	-	1	1	-	-	1	1	1	-	-
White, both sexes	-	1	-	2	1	-	2	1	2	1	-	-
Male	-	1	-	1	1	-	2	1	1	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	9	9	13	13	19	18	25	14	3	-	-
Male	3	4	5	4	9	11	10	11	8	1	-	-
Female	3	5	4	9	4	8	8	14	6	2	-	-
White, both sexes	5	7	5	10	11	17	15	23	13	3	-	-
Male	2	2	3	3	8	10	8	10	7	1	-	-
Female	3	5	2	7	3	7	7	13	6	2	-	-
Black, both sexes	-	2	3	2	2	1	1	1	-	-	-	-
Male	-	2	2	1	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	1	2	1	1	-	-	-
Male	1	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	1	1	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Colorado												
All races, both sexes	-	-	2	6	3	5	4	4	8	1	-	-
Male	-	-	2	4	1	3	2	2	2	-	-	-
Female	-	-	-	2	2	2	2	2	6	1	-	-
White, both sexes	-	-	2	5	3	4	4	4	8	1	-	-
Male	-	-	2	3	1	2	2	2	2	-	-	-
Female	-	-	-	2	2	2	2	2	6	1	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	-	-	3	5	5	4	3	-	-
Male	1	-	1	-	-	1	1	1	-	1	-	-
Female	-	-	-	-	-	2	4	4	4	2	-	-
White, both sexes	1	-	1	-	-	3	3	5	3	3	-	-
Male	1	-	1	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	2	3	4	3	2	-	-
Black, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Delaware												
All races, both sexes	-	-	-	-	-	1	1	-	1	1	1	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	1	-	1	-
White, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Florida												
All races, both sexes	2	5	5	5	5	13	16	10	9	5	3	-
Male	2	2	3	2	4	6	5	6	2	-	-	-
Female	-	3	2	3	1	7	11	4	7	5	3	-
White, both sexes	2	4	4	4	4	11	15	8	6	3	3	-
Male	2	1	2	1	4	5	5	4	2	-	-	-
Female	-	3	2	3	-	6	10	4	4	3	3	-
Black, both sexes	-	1	1	1	1	2	1	2	3	2	-	-
Male	-	1	1	1	-	1	-	2	-	-	-	-
Female	-	-	-	-	1	1	1	-	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	1	2	3	2	3	10	5	3	1	2	-
Male	2	1	1	2	1	2	3	1	1	-	-	-
Female	1	-	1	1	1	1	7	4	2	1	2	-
White, both sexes	2	1	2	2	1	2	8	4	1	-	-	-
Male	2	1	1	2	-	2	2	1	-	-	-	-
Female	-	-	1	-	1	-	6	3	1	-	-	-
Black, both sexes	1	-	-	1	1	1	2	1	2	1	2	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	-	1	-	1	1	1	1	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	2	-	2	1	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	2	-	2	1	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Illinois												
All races, both sexes	3	3	4	3	7	10	18	17	8	1	-	-
Male	2	2	3	2	3	4	12	4	2	-	-	-
Female	1	1	1	1	4	6	6	13	6	1	-	-
White, both sexes	2	3	4	2	5	6	16	14	8	1	-	-
Male	1	2	3	1	2	1	12	3	2	-	-	-
Female	1	1	1	1	3	5	4	11	6	1	-	-
Black, both sexes	1	-	-	1	2	4	2	3	-	-	-	-
Male	1	-	-	1	1	3	-	1	-	-	-	-
Female	-	-	-	-	1	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	2	4	2	4	5	5	5	4	-	-	-
Male	3	-	-	1	2	3	4	3	-	-	-	-
Female	-	2	4	1	2	2	1	2	4	-	-	-
White, both sexes	1	2	3	1	4	5	4	4	2	-	-	-
Male	1	-	-	1	2	3	3	3	-	-	-	-
Female	-	2	3	-	2	2	1	1	2	-	-	-
Black, both sexes	2	-	1	1	-	-	1	1	2	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Iowa												
All races, both sexes	-	-	-	-	2	1	7	8	4	-	-	-
Male	-	-	-	-	2	-	7	3	1	-	-	-
Female	-	-	-	-	-	1	-	5	3	-	-	-
White, both sexes	-	-	-	-	2	1	7	8	4	-	-	-
Male	-	-	-	-	2	-	7	3	1	-	-	-
Female	-	-	-	-	-	1	-	5	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	1	-	-	1	6	1	4	1	-	-
Male	-	-	-	-	-	1	2	-	1	-	-	-
Female	-	-	1	-	-	-	4	1	3	1	-	-
White, both sexes	-	-	1	-	-	1	5	1	4	1	-	-
Male	-	-	-	-	-	1	2	-	1	-	-	-
Female	-	-	1	-	-	-	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Kentucky												
All races, both sexes	1	1	2	2	-	1	3	3	2	-	-	-
Male	1	1	2	2	-	1	-	2	2	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
White, both sexes	1	1	2	1	-	1	3	3	2	-	-	-
Male	1	1	2	1	-	1	-	2	2	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	2	1	5	4	4	2	2	3	1	-
Male	-	-	-	1	2	1	-	1	-	1	-	-
Female	-	-	2	-	3	3	4	1	2	2	1	-
White, both sexes	-	-	1	1	3	3	2	1	1	2	1	-
Male	-	-	-	1	2	1	-	1	-	1	-	-
Female	-	-	1	-	1	2	2	-	1	1	1	-
Black, both sexes	-	-	1	-	2	1	2	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Maine												
All races, both sexes	2	-	-	-	2	3	4	2	2	1	1	-
Male	2	-	-	-	2	1	4	2	-	1	-	-
Female	-	-	-	-	-	2	-	-	2	-	1	-
White, both sexes	2	-	-	-	2	3	4	2	2	1	1	-
Male	2	-	-	-	2	1	4	2	-	1	-	-
Female	-	-	-	-	-	2	-	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	3	2	1	2	5	7	8	4	2	1	-
Male	-	-	-	1	-	2	1	1	1	1	1	-
Female	-	3	2	-	2	3	6	7	3	1	-	-
White, both sexes	-	-	1	1	2	4	6	5	4	2	-	-
Male	-	-	-	1	-	1	1	1	1	1	-	-
Female	-	-	1	-	2	3	5	4	3	1	-	-
Black, both sexes	-	3	1	-	-	1	1	3	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	3	1	-	-	-	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Massachusetts												
All races, both sexes	-	2	2	3	4	8	9	7	8	2	-	-
Male	-	-	2	1	2	5	6	3	4	-	-	-
Female	-	2	-	2	2	3	3	4	4	2	-	-
White, both sexes	-	1	2	3	3	8	8	7	8	2	-	-
Male	-	-	2	1	2	5	6	3	4	-	-	-
Female	-	1	-	2	1	3	2	4	4	2	-	-
Black, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	2	2	5	5	15	6	11	2	3	-
Male	-	1	-	1	2	2	9	-	5	-	1	-
Female	-	-	2	1	3	3	6	6	6	2	2	-
White, both sexes	-	1	1	2	5	3	14	5	10	2	3	-
Male	-	1	-	1	2	2	8	-	4	-	1	-
Female	-	-	1	1	3	1	6	5	6	2	2	-
Black, both sexes	-	-	1	-	-	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Minnesota												
All races, both sexes	1	1	-	-	2	7	6	7	11	1	2	-
Male	1	1	-	-	1	3	2	4	3	-	-	-
Female	-	-	-	-	1	4	4	3	8	1	2	-
White, both sexes	1	1	-	-	2	7	6	7	11	1	2	-
Male	1	1	-	-	1	3	2	4	3	-	-	-
Female	-	-	-	-	1	4	4	3	8	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	-	1	3	2	-	1	-	-
Male	-	-	1	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	1	-	-	1	2	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Missouri												
All races, both sexes	1	-	2	3	2	6	10	4	5	2	1	-
Male	1	-	2	2	2	3	5	-	1	1	1	-
Female	-	-	-	1	-	3	5	4	4	1	-	-
White, both sexes	1	-	1	1	2	5	10	3	5	2	1	-
Male	1	-	1	1	2	2	5	-	1	1	1	-
Female	-	-	-	-	-	3	5	3	4	1	-	-
Black, both sexes	-	-	1	2	-	1	-	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Nebraska												
All races, both sexes	-	-	-	2	2	-	1	3	3	1	-	-
Male	-	-	-	1	-	-	-	3	1	-	-	-
Female	-	-	-	1	2	-	1	-	2	1	-	-
White, both sexes	-	-	-	2	2	-	1	3	3	1	-	-
Male	-	-	-	1	-	-	-	3	1	-	-	-
Female	-	-	-	1	2	-	1	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	1	4	3	2	2	1	-	-	-
Male	-	-	-	1	3	1	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	1	-	-	-
White, both sexes	-	-	-	1	3	2	2	2	1	-	-	-
Male	-	-	-	1	2	-	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
New Hampshire												
All races, both sexes	1	-	-	1	-	2	3	2	-	1	-	-
Male	1	-	-	1	-	2	2	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	1	-	-	1	-	2	3	2	-	1	-	-
Male	1	-	-	1	-	2	2	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	4	4	5	8	14	11	6	-	-	-
Male	2	-	2	1	2	4	6	6	2	-	-	-
Female	-	-	2	3	3	4	8	5	4	-	-	-
White, both sexes	2	-	3	2	5	8	11	10	5	-	-	-
Male	2	-	1	-	2	4	6	5	2	-	-	-
Female	-	-	2	2	3	4	5	5	3	-	-	-
Black, both sexes	-	-	1	2	-	-	3	1	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
New Mexico												
All races, both sexes	1	-	-	1	1	-	1	3	2	-	-	-
Male	1	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	1	-	-	1	2	-	-	-
White, both sexes	1	-	-	1	1	-	1	3	2	-	-	-
Male	1	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	1	1	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	6	3	6	9	10	18	11	15	9	1	-
Male	2	3	2	3	5	8	8	1	4	-	-	-
Female	4	3	1	3	4	2	10	10	11	9	1	-
White, both sexes	4	4	3	6	5	8	14	8	13	7	1	-
Male	2	1	2	3	2	7	5	1	3	-	-	-
Female	2	3	1	3	3	1	9	7	10	7	1	-
Black, both sexes	2	2	-	-	4	2	4	3	2	2	-	-
Male	-	2	-	-	3	1	3	-	1	-	-	-
Female	2	-	-	-	1	1	1	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
North Carolina												
All races, both sexes	4	-	3	3	7	4	10	13	7	2	1	-
Male	2	-	2	1	3	-	7	3	3	-	-	-
Female	2	-	1	2	4	4	3	10	4	2	1	-
White, both sexes	3	-	1	-	3	2	7	8	6	2	1	-
Male	2	-	1	-	2	-	5	2	3	-	-	-
Female	1	-	-	-	1	2	2	6	3	2	1	-
Black, both sexes	1	-	2	3	3	2	2	5	1	-	-	-
Male	-	-	1	1	1	-	1	1	-	-	-	-
Female	1	-	1	2	2	2	1	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Ohio												
All races, both sexes	2	6	1	6	9	10	12	14	10	-	-	-
Male	-	4	1	4	7	8	5	7	3	-	-	-
Female	2	2	-	2	2	2	7	7	7	-	-	-
White, both sexes	2	5	1	6	7	6	9	10	9	-	-	-
Male	-	3	1	4	6	5	4	7	2	-	-	-
Female	2	2	-	2	1	1	5	3	7	-	-	-
Black, both sexes	-	1	-	-	2	4	3	4	1	-	-	-
Male	-	1	-	-	1	3	1	-	1	-	-	-
Female	-	-	-	-	1	1	2	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	1	2	3	3	4	3	-	1	1	-
Male	-	1	-	1	1	2	1	1	-	-	-	-
Female	-	-	1	1	2	1	3	2	-	1	1	-
White, both sexes	-	1	1	1	3	3	4	3	-	1	-	-
Male	-	1	-	-	1	2	1	1	-	-	-	-
Female	-	-	1	1	2	1	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Oregon												
All races, both sexes	-	2	1	1	3	4	3	4	7	1	-	1
Male	-	2	1	-	2	2	2	3	3	-	-	1
Female	-	-	-	1	1	2	1	1	4	1	-	-
White, both sexes	-	1	1	1	3	4	3	4	7	1	-	-
Male	-	1	1	-	2	2	2	3	3	-	-	-
Female	-	-	-	1	1	2	1	1	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	3	3	9	11	16	22	12	5	-	-
Male	1	1	3	3	4	5	9	11	3	-	-	-
Female	-	-	-	-	5	6	7	11	9	5	-	-
White, both sexes	1	-	3	2	9	10	15	19	12	4	-	-
Male	1	-	3	2	4	4	8	9	3	-	-	-
Female	-	-	-	-	5	6	7	10	9	4	-	-
Black, both sexes	-	1	-	1	-	1	1	3	-	1	-	-
Male	-	1	-	1	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Rhode Island												
All races, both sexes	-	-	-	1	1	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	3	-	1	-	-	-
White, both sexes	-	-	-	1	1	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	1	-	3	2	4	3	2	-	1	-
Male	1	1	-	-	1	1	2	1	1	-	-	-
Female	-	1	1	-	2	1	2	2	1	-	1	-
White, both sexes	1	1	-	-	3	1	3	1	2	-	1	-
Male	1	-	-	-	1	1	2	1	1	-	-	-
Female	-	1	-	-	2	-	1	-	1	-	1	-
Black, both sexes	-	1	1	-	-	1	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
South Dakota												
All races, both sexes	-	-	-	-	1	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	2	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	5	3	5	3	8	9	9	4	3	-	-
Male	2	2	1	4	-	3	3	2	1	1	-	-
Female	1	3	2	1	3	5	6	7	3	2	-	-
White, both sexes	1	4	3	5	3	5	6	7	4	3	-	-
Male	1	2	1	4	-	1	2	2	1	1	-	-
Female	-	2	2	1	3	4	4	5	3	2	-	-
Black, both sexes	2	1	-	-	-	3	3	2	-	-	-	-
Male	1	-	-	-	-	2	1	-	-	-	-	-
Female	1	1	-	-	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Texas												
All races, both sexes	4	3	14	4	14	7	20	15	9	3	1	-
Male	3	3	9	2	7	4	12	7	4	1	-	-
Female	1	-	5	2	7	3	8	8	5	2	1	-
White, both sexes	3	3	8	4	14	5	16	9	9	2	1	-
Male	3	3	5	2	7	3	9	7	4	1	-	-
Female	-	-	3	2	7	2	7	2	5	1	1	-
Black, both sexes	1	-	6	-	-	2	3	6	-	1	-	-
Male	-	-	4	-	-	1	2	-	-	-	-	-
Female	1	-	2	-	-	1	1	6	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	1	-	2	1	1	2	2	-	-	-
Male	1	-	1	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
White, both sexes	1	-	1	-	2	1	1	2	2	-	-	-
Male	1	-	1	-	-	1	1	2	1	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	2	1	2	2	5	7	2	1	-	-
Male	1	-	-	-	-	2	2	2	-	-	-	-
Female	-	-	2	1	2	-	3	5	2	1	-	-
White, both sexes	1	-	2	1	2	-	3	5	-	1	-	-
Male	1	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	2	1	2	-	1	3	-	1	-	-
Black, both sexes	-	-	-	-	-	2	2	2	2	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Washington												
All races, both sexes	-	4	4	-	5	4	4	3	4	-	2	-
Male	-	2	2	-	3	2	2	2	2	-	1	-
Female	-	2	2	-	2	2	2	1	2	-	1	-
White, both sexes	-	4	3	-	5	4	4	3	4	-	2	-
Male	-	2	1	-	3	2	2	2	2	-	1	-
Female	-	2	2	-	2	2	2	1	2	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	1	2	4	6	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	4	6	-	-	-	-
White, both sexes	-	-	1	-	1	2	4	6	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	4	6	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Wisconsin												
All races, both sexes	1	-	1	2	1	2	8	6	6	4	-	-
Male	-	-	-	1	1	-	5	2	1	3	-	-
Female	1	-	1	1	-	2	3	4	5	1	-	-
White, both sexes	1	-	1	2	1	2	7	5	6	4	-	-
Male	-	-	-	1	1	-	5	2	1	3	-	-
Female	1	-	1	1	-	2	2	3	5	1	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-